

Dafang Zhao , Xuexiao You and Delfim F. M. Torres**

*Discrete Opial type inequalities for interval-valued
functions*

MATHEMATICAL INEQUALITIES & APPLICATIONS

AIMS AND SCOPE

Mathematical Inequalities & Applications (MIA, Math. Inequal. Appl.) brings together original research papers in all areas of mathematics, provided they are concerned with inequalities or their role. From time to time MIA will publish invited survey articles. Short notes with interesting results or open problems will also be accepted.

MIA is published quarterly, in January, April, July and October.

SUBMISSION

Manuscripts should be submitted electronically in PostScript or Adobe Acrobat PDF format to the MIA Editorial Office through MIA page: www.ele-math.com or by e-mail mia@ele-math.com

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the MIA regarding the article. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to MIA. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal, and that its submission for publication has been approved by all the authors.

PREPARATION OF MANUSCRIPT

Manuscripts should be written in English.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send final version of file(s) to the Editorial Office.

TITLE PAGE

A title page should include the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author and running head less than 60 characters. Authors should be listed alphabetically.

ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2000) with primary (and secondary) subject classification codes and a list of 4-5 key words must be given.

FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, grey scale only, preferably at resolution of 300 dpi or higher, sized to fit a journal page, and in EPS, TIFF or JPEG format. Figures should have a short descriptive title and should be numbered consecutively in order of appearance in the text. The position of each figure in the text should be indicated.

REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The author should consult Mathematical Reviews for the standard abbreviations of journal names.

PROOFS

The author will receive proofs only once. Late return of the proofs will delay the article to a later issue.

PAGE CHARGES, OFFPRINTS

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from 'MIA' prior to publication.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

MIA is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

Mathematical Inequalities & Applications
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia
www.ele-math.com
e-mail: mia@ele-math.com
Fax: +385 1 6008799

DISCRETE OPIAL TYPE INEQUALITIES FOR INTERVAL-VALUED FUNCTIONS

DAFANG ZHAO*, XUEXIAO YOU AND DELFIM F. M. TORRES*

(Communicated by I. Perić)

Abstract. We introduce the forward (backward) gH -difference operator of interval sequences, and establish some new discrete Opial type inequalities for interval-valued functions. Further, we obtain generalizations of classical discrete Opial type inequalities. Some examples are presented to illustrate our results.

1. Introduction

The theory of inequalities has a long history but, from the applicative point of view, it fell into neglect for hundreds of years because of lack of applications to other branch of mathematics as well as other sciences, such as physics and engineering. Only in 1934 did Hardy, Littlewood and Pólya transformed the field of inequalities from a collection of isolated formulas into a systematic discipline [24]. After that, an enormous amount of effort has been devoted to the discovery of new types of inequalities and to applications of inequalities [1].

It is known that many physical problems in various applications are governed by finite difference equations. Moreover, discrete inequalities play an important role in the continuing development of the theory of difference equations. This importance seems to have increased considerably during the past decades. It has attracted the attention of a large number of researchers, stimulated new research directions, and influenced various aspects of difference equations and applications. Among the many types of inequalities, those associated with the names of Jensen [12, 17, 18], Hilbert [26, 45], Wirtinger [2, 4, 19], Chebyshev [36, 49], Gronwall–Bellman [20, 37] and Opial [5, 7, 23, 27, 34, 35] have deep roots and made a great impact on various branches of mathematics. The development of discrete inequalities resulted in a renewal of interest in the field and has attracted interest from more researchers [6, 10, 11, 16, 22, 28, 29, 30, 31, 39, 40, 41, 42].

More recently, some of classical inequalities have been extended to set-valued functions by Nikodem et al. [32], Štrboja et al. [44], and Zhang et al. [46, 47], especially to interval-valued functions by Chalco-Cano et al. [8, 9], Costa et al. [13, 14, 15], Flores-Franulić et al. [21], Román-Flores et al. [38], and Zhao et al. [48, 49, 50, 51].

Mathematics subject classification (2020): 26D15, 26E50, 65G30.

Keywords and phrases: Discrete Opial type inequalities, gH -difference operators, interval-valued functions.

* Corresponding author.

The present article is, in some sense, a continuation of the previous work [50]. Here, we establish some new discrete inequalities of Opial type involving sequences of intervals and their forward (backward) difference operator. Furthermore, our present results can be considered as tools for further research in interval difference equations and inequalities for interval-valued functions, among others.

The paper is organized as follows. Section 2 contains some necessary preliminaries. In Section 3, we present some new interval Opial type inequalities involving the backward gH-difference operator, and present some examples to illustrate our theorems. In Section 4, some new discrete Opial type inequalities, involving the forward gH-difference operator, are given. Finally, in the concluding Section 5, we summarize our results and outline some possible future work directions.

2. Preliminaries

We begin by recalling some basic notations, definitions, and results of interval analysis. We define an interval u by

$$u = [\underline{u}, \bar{u}] = \{t \in \mathbb{R} \mid \underline{u} \leq t \leq \bar{u}\}.$$

We write $len(u) = \bar{u} - \underline{u}$. If $len(u) = 0$, then u is called a degenerate interval. The set of all intervals of \mathbb{R} is denoted by $\mathbb{R}_{\mathcal{I}}$. For $\lambda \in \mathbb{R}$ and $u \in \mathbb{R}_{\mathcal{I}}$, λu is defined by

$$\lambda [\underline{u}, \bar{u}] = \begin{cases} [\lambda \underline{u}, \lambda \bar{u}] & \text{if } \lambda \geq 0, \\ [\lambda \bar{u}, \lambda \underline{u}] & \text{if } \lambda < 0. \end{cases}$$

For $u = [\underline{u}, \bar{u}]$ and $v = [\underline{v}, \bar{v}]$, the four arithmetic operators $(+, -, \cdot, /)$ are defined by

$$u + v = [\underline{u} + \underline{v}, \bar{u} + \bar{v}],$$

$$u - v = [\underline{u} - \bar{v}, \bar{u} - \underline{v}],$$

$$u \cdot v = [\min\{\underline{u}\underline{v}, \underline{u}\bar{v}, \bar{u}\underline{v}, \bar{u}\bar{v}\}, \max\{\underline{u}\underline{v}, \underline{u}\bar{v}, \bar{u}\underline{v}, \bar{u}\bar{v}\}],$$

$$u/v = [\min\{\underline{u}/\underline{v}, \underline{u}/\bar{v}, \bar{u}/\underline{v}, \bar{u}/\bar{v}\}, \max\{\underline{u}/\underline{v}, \underline{u}/\bar{v}, \bar{u}/\underline{v}, \bar{u}/\bar{v}\}], \text{ where } 0 \notin [\underline{v}, \bar{v}].$$

Note that $\mathbb{R}_{\mathcal{I}}$ with the above operations (i.e., the Minkowski addition and the scalar multiplication) is *not* a linear space since an interval does not have inverse element and, therefore, the subtraction does not have adequate properties. For example, when subtracting two intervals u and v , the width of the result is the sum of the widths of u and v , i.e.,

$$len(u - v) = len(u) + len(v).$$

To partially overcome this situation, Hukuhara [25] introduced the following H-difference:

$$u \ominus v = w \Leftrightarrow u = v + w.$$

Unfortunately, the H-difference does not always exist for any u and v .

In [43], Stefanini introduced the gH-difference as follows:

$$u \ominus_g v = w \Leftrightarrow \begin{cases} (a) u = v + w, \\ \text{or } (b) v = u + (-1)w. \end{cases}$$

The gH-difference always exists for any u and v . We also have

$$u \ominus_g v = [\min\{\underline{u} - \underline{v}, \bar{u} - \bar{v}\}, \max\{\underline{u} - \underline{v}, \bar{u} - \bar{v}\}].$$

The Hausdorff distance between u and v is defined by

$$d(u, v) = \max\{|\underline{u} - \underline{v}|, |\bar{u} - \bar{v}|\}.$$

Then, $(\mathbb{R}_{\mathcal{I}}, d)$ is a complete metric space. Note that $(\mathbb{R}_{\mathcal{I}}, +, \cdot)$ is a quasi-linear space (see [15]) equipped with the quasi-norm $\|\cdot\|$, which is given by

$$\|u\| = d(u, [0, 0]) = d([\underline{u}, \bar{u}], [0, 0]) = \max\{|\underline{u}|, |\bar{u}|\}$$

for all $u \in \mathbb{R}_{\mathcal{I}}$.

On $[a, b]$, u_i is called increasing if and only if \underline{u}_i and \bar{u}_i are increasing; u_i and v_i are synchronous (asynchronous) monotone if they have the same (opposite) monotonicity; u_i is μ -increasing if $len(u_i)$ is increasing. One defines u^λ by

$$u^\lambda = \{t^\lambda \mid t \in [\underline{u}, \bar{u}]\}.$$

For convenience, we now recall the classical Opial’s inequality:

THEOREM 1. (continuous Opial inequality [33]) *Let $F \in C^1[0, h]$, $F(0) = F(h) = 0$ and $F(t) > 0$ for $t \in (0, h)$. Then,*

$$\int_0^h |F(t)F'(t)|dt \leq \frac{h}{4} \int_0^h (F'(t))^2 dt, \tag{1}$$

where $\frac{h}{4}$ is the best possible.

A discrete analogue of Theorem 1 is the following:

THEOREM 2. (discrete Opial inequality [3]) *Let $\{u_i\}_{i=0}^n$ be a sequence of numbers with $u_0 = 0$ and $u_n = 0$. Then,*

$$\sum_{i=1}^{n-1} |u_i \Delta u_i| \leq \frac{1}{2} \left[\frac{n+1}{2} \right] \sum_{i=0}^{n-1} |\Delta u_i|^2, \tag{2}$$

where Δ is the forward difference operator and $[\cdot]$ is the greatest integer function.

Many generalizations of Theorem 2 are available in the literature: see, e.g., [5, 7, 23]. In Sections 3 and 4, we give several extensions of Theorem 2 for sequences of intervals.

3. Opial type inequalities involving the backward/nabla gH-difference operator

DEFINITION 1. Let $\{u_i\}$ be a sequence of intervals. We define the forward (delta) gH-difference operator Δu by

$$\Delta u_i = u_{i+1} \ominus_g u_i.$$

Similarly, we define the backward (nabla) gH-difference operator ∇u by

$$\nabla u_i = u_i \ominus_g u_{i-1}.$$

REMARK 1. Note that if $\{u_i\}$ is a sequence of degenerate intervals, then the forward (backward) gH-difference operator reduces to the classical forward (backward) difference operator.

Lemma 1 has been obtained by Lee in [27]. Here we give a new and more direct proof.

LEMMA 1. (cf. [27]) *Let $\{u_i\}_{i=1}^n$ be a non-decreasing sequence of non-negative real numbers, $u_0 = 0$, and $\lambda_1, \lambda_2 \geq 1$. Then,*

$$\sum_{i=1}^n u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n (\nabla u_i)^{\lambda_1 + \lambda_2}. \tag{3}$$

Proof. Since $\nabla u_i = u_i - u_{i-1}$, we have $u_i = \sum_{j=1}^i \nabla u_j$. We may rewrite (3) as

$$\sum_{i=1}^n \left(\sum_{j=1}^i \nabla u_j \right)^{\lambda_1} (\nabla u_i)^{\lambda_2} \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n (\nabla u_i)^{\lambda_1 + \lambda_2}. \tag{4}$$

We shall prove (4) by induction on n . Clearly, (4) holds with $n = 1$. Assume that it holds for n , so that

$$\begin{aligned} & \sum_{i=1}^{n+1} \left(\sum_{j=1}^i \nabla u_j \right)^{\lambda_1} (\nabla u_i)^{\lambda_2} \\ & \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \left(\sum_{i=1}^n (\nabla u_i)^{\lambda_1 + \lambda_2} + \frac{\lambda_1 + \lambda_2}{\lambda_2} \left(\frac{1}{n+1} \sum_{j=1}^{n+1} \nabla u_j \right)^{\lambda_1} (\nabla u_{n+1})^{\lambda_2} \right) \\ & \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \left(\sum_{i=1}^n (\nabla u_i)^{\lambda_1 + \lambda_2} + \frac{\lambda_1 + \lambda_2}{\lambda_2} A_{n+1}^{\lambda_1} (\nabla u_{n+1})^{\lambda_2} \right), \end{aligned} \tag{5}$$

where $A_{n+1} = \frac{1}{n+1} \sum_{j=1}^{n+1} \nabla u_j$. Using Young's inequality, we have

$$A_{n+1}^{\lambda_1} (\nabla u_{n+1})^{\lambda_2} \leq \frac{\lambda_1}{\lambda_1 + \lambda_2} A_{n+1}^{\lambda_1 + \lambda_2} + \frac{\lambda_2}{\lambda_1 + \lambda_2} (\nabla u_{n+1})^{\lambda_1 + \lambda_2}.$$

Then,

$$\frac{\lambda_1 + \lambda_2}{\lambda_2} A_{n+1}^{\lambda_1} (\nabla u_{n+1})^{\lambda_2} \leq \frac{\lambda_1}{\lambda_2} A_{n+1}^{\lambda_1 + \lambda_2} + (\nabla u_{n+1})^{\lambda_1 + \lambda_2}. \quad (6)$$

Thanks to Hölder's inequality, it follows that

$$\begin{aligned} A_{n+1} &= \frac{1}{n+1} \sum_{j=1}^{n+1} \nabla u_j \\ &\leq \left(\sum_{j=1}^{n+1} \left(\frac{1}{n+1} \right)^{\frac{\lambda_1 + \lambda_2}{\lambda_1 + \lambda_2 - 1}} \right)^{\frac{\lambda_1 + \lambda_2 - 1}{\lambda_1 + \lambda_2}} \left(\sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2} \right)^{\frac{1}{\lambda_1 + \lambda_2}} \\ &\leq \left(\frac{1}{n+1} \right)^{\frac{1}{\lambda_1 + \lambda_2}} \left(\sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2} \right)^{\frac{1}{\lambda_1 + \lambda_2}}. \end{aligned}$$

Consequently, we get

$$A_{n+1}^{\lambda_1 + \lambda_2} \leq \frac{1}{n+1} \sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2}. \quad (7)$$

Thus, combining (5), (6) and (7), we have

$$\begin{aligned} &\sum_{i=1}^{n+1} \left(\sum_{j=1}^i \nabla u_j \right)^{\lambda_1} (\nabla u_i)^{\lambda_2} \\ &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \left(\sum_{i=1}^n (\nabla u_i)^{\lambda_1 + \lambda_2} + \frac{\lambda_1 + \lambda_2}{\lambda_2} A_{n+1}^{\lambda_1} (\nabla u_{n+1})^{\lambda_2} \right) \\ &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \left(\sum_{i=1}^n (\nabla u_i)^{\lambda_1 + \lambda_2} + \frac{\lambda_1}{\lambda_2} A_{n+1}^{\lambda_1 + \lambda_2} + (\nabla u_{n+1})^{\lambda_1 + \lambda_2} \right) \\ &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \left(\sum_{i=1}^{n+1} (\nabla u_i)^{\lambda_1 + \lambda_2} + \frac{\lambda_1}{n+1} \sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2} \right) \\ &= \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \cdot \frac{n+1 + \lambda_1}{n+1} \cdot \sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2} \\ &= \frac{\lambda_2 \left[(n+1)^{\lambda_1} + \lambda_1 (n+1)^{\lambda_1 - 1} \right]}{\lambda_1 + \lambda_2} \sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2} \\ &\leq \frac{\lambda_2(n+2)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{j=1}^{n+1} (\nabla u_j)^{\lambda_1 + \lambda_2}. \end{aligned}$$

The proof is complete. \square

Thanks to Lemma 1, we can easily obtain the following Lemma 2,

LEMMA 2. Let $\{u_i\}_{i=1}^n$ be a sequence of numbers, $u_0 = 0$, and $\lambda_1, \lambda_2 \geq 1$. Then,

$$\sum_{i=1}^n |u_i|^{\lambda_1} |\nabla u_i|^{\lambda_2} \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n |\nabla u_i|^{\lambda_1 + \lambda_2}. \tag{8}$$

Proof. Since $|\nabla u_i| = |u_i - u_{i-1}|$, we have $|u_i| \leq \sum_{j=1}^i |\nabla u_j|$. The rest of proof is similar to that of Lemma 1 and is omitted here. \square

We are now ready to formulate and prove our first original result.

THEOREM 3. Let $\{u_i\}_{i=1}^n$ be a sequence of intervals, $u_0 = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If u_i is monotone and μ -increasing, then

$$\sum_{i=1}^n \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \tag{9}$$

Proof. Suppose that u_i is increasing and μ -increasing. Then,

$$u_i^{\lambda_1} = \left[\underline{u}_i^{\lambda_1}, \overline{u}_i^{\lambda_1} \right], \quad (\nabla u_i)^{\lambda_2} = \left[(\underline{\nabla} u_i)^{\lambda_2}, (\overline{\nabla} u_i)^{\lambda_2} \right].$$

Consequently, we obtain that

$$u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} = \left[\underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2} \right].$$

If u_i is decreasing and μ -increasing, then

$$u_i^{\lambda_1} = \begin{cases} \left[\underline{u}_i^{\lambda_1}, \overline{u}_i^{\lambda_1} \right] & \text{if } \lambda_1 \text{ is odd,} \\ \left[\overline{u}_i^{\lambda_1}, \underline{u}_i^{\lambda_1} \right] & \text{if } \lambda_1 \text{ is even,} \end{cases}$$

$$(\nabla u_i)^{\lambda_2} = \begin{cases} \left[(\underline{\nabla} u_i)^{\lambda_2}, (\overline{\nabla} u_i)^{\lambda_2} \right] & \text{if } \lambda_2 \text{ is odd,} \\ \left[(\overline{\nabla} u_i)^{\lambda_2}, (\underline{\nabla} u_i)^{\lambda_2} \right] & \text{if } \lambda_2 \text{ is even.} \end{cases}$$

Consequently, we obtain

$$u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} = \begin{cases} \left[\overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2}, \underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2} \right] & \text{if } \lambda_1 \text{ and } \lambda_2 \text{ are odd,} \\ \left[\overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2}, \underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2} \right] & \text{if } \lambda_1 \text{ and } \lambda_2 \text{ are even,} \\ \left[\underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2} \right] & \text{if } \lambda_1 \text{ is odd and } \lambda_2 \text{ is even,} \\ \left[\underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2} \right] & \text{if } \lambda_1 \text{ is even and } \lambda_2 \text{ is odd.} \end{cases}$$

By Lemma 2, it follows that

$$\begin{aligned}
 & \sum_{i=1}^n \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \\
 &= \sum_{i=1}^n \left\| \left[\min \{ \underline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \}, \max \{ \underline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \} \right] \right\| \\
 &= \sum_{i=1}^n \max \left\{ \left| \underline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right|, \left| \overline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right| \right\} \\
 &= \max \left\{ \sum_{i=1}^n \left| \underline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right|, \sum_{i=1}^n \left| \overline{u}_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right| \right\} \\
 &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \max \left\{ \sum_{i=1}^n \left| \nabla u_i^{\lambda_1 + \lambda_2} \right|, \sum_{i=1}^n \left| \overline{\nabla u_i}^{\lambda_1 + \lambda_2} \right| \right\} \\
 &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n \max \left\{ \left| \nabla u_i \right|^{\lambda_1 + \lambda_2}, \left| \overline{\nabla u_i} \right|^{\lambda_1 + \lambda_2} \right\} \\
 &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n \|\nabla u_i\|^{\lambda_1 + \lambda_2}.
 \end{aligned}$$

This concludes the proof. \square

Follows an example of application of our Theorem 3.

EXAMPLE 1. Suppose that $\{u_i\}_{i=0}^n = \{[0, 0], [1, 2], [2, 4], \dots, [n, 2n]\}$ and $\lambda_1, \lambda_2 \geq 1$. By Theorem 3, we have

$$\begin{aligned}
 \sum_{i=1}^n \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| &= \sum_{i=1}^n \left\| [i^{\lambda_1}, (2i)^{\lambda_1}] \cdot [1, 2]^{\lambda_2} \right\| \\
 &= 2^{\lambda_1 + \lambda_2} \sum_{i=1}^n i^{\lambda_1} \\
 &\leq \frac{\lambda_2 n(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} 2^{\lambda_1 + \lambda_2} \\
 &= \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n 2^{\lambda_1 + \lambda_2} \\
 &= \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n \|\nabla u_i\|^{\lambda_1 + \lambda_2}.
 \end{aligned}$$

LEMMA 3. Let $\{u_i\}_{i=1}^m$ be a sequence of numbers, $u_m = 0$, and $\lambda_1, \lambda_2 \geq 1$. Then,

$$\sum_{i=n}^{m-1} |u_i|^{\lambda_1} |\nabla u_i|^{\lambda_2} \leq \frac{\lambda_2(m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=n}^m |\nabla u_i|^{\lambda_1 + \lambda_2}. \tag{10}$$

Proof. Since $|\nabla u_i| = |u_i - u_{i-1}|$, we have $|u_i| \leq \sum_{j=i+1}^m |\nabla u_j|$. The rest of proof is similar to that of Lemma 1 and is omitted here. \square

Similarly to Theorem 3, we obtain an analogous result when u_i is monotone but μ -decreasing instead of μ -increasing.

THEOREM 4. *Let $\{u_i\}_{i=0}^m$ be a sequence of intervals, $u_m = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If u_i is monotone and μ -decreasing, then*

$$\sum_{i=n}^{m-1} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \leq \frac{\lambda_2(m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=n}^m \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \tag{11}$$

Proof. Suppose that u_i is increasing and μ -decreasing. Then,

$$u_i^{\lambda_1} = \begin{cases} \left[\underline{u}_i^{\lambda_1}, \overline{u}_i^{\lambda_1} \right] & \text{if } \lambda_1 \text{ is odd,} \\ \left[\overline{u}_i^{\lambda_1}, \underline{u}_i^{\lambda_1} \right] & \text{if } \lambda_1 \text{ is even,} \end{cases}$$

and

$$(\nabla u_i)^{\lambda_2} = \left[(\overline{\nabla u_i})^{\lambda_2}, (\underline{\nabla u_i})^{\lambda_2} \right].$$

Consequently, we obtain

$$u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} = \begin{cases} \left[\underline{u}_i^{\lambda_1} (\underline{\nabla u_i})^{\lambda_2}, \overline{u}_i^{\lambda_1} (\overline{\nabla u_i})^{\lambda_2} \right] & \text{if } \lambda_1 \text{ is odd,} \\ \left[\overline{u}_i^{\lambda_1} (\overline{\nabla u_i})^{\lambda_2}, \underline{u}_i^{\lambda_1} (\underline{\nabla u_i})^{\lambda_2} \right] & \text{if } \lambda_1 \text{ is even.} \end{cases} \tag{12}$$

If u_i is decreasing and μ -decreasing, then

$$u_i^{\lambda_1} = \left[\underline{u}_i^{\lambda_1}, \overline{u}_i^{\lambda_1} \right],$$

$$(\nabla u_i)^{\lambda_2} = \begin{cases} \left[(\overline{\nabla u_i})^{\lambda_2}, (\underline{\nabla u_i})^{\lambda_2} \right] & \text{if } \lambda_2 \text{ is odd,} \\ \left[(\underline{\nabla u_i})^{\lambda_2}, (\overline{\nabla u_i})^{\lambda_2} \right] & \text{if } \lambda_2 \text{ is even.} \end{cases}$$

Consequently, we obtain

$$u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} = \begin{cases} \left[\overline{u}_i^{\lambda_1} (\overline{\nabla u_i})^{\lambda_2}, \underline{u}_i^{\lambda_1} (\underline{\nabla u_i})^{\lambda_2} \right] & \text{if } \lambda_2 \text{ is odd,} \\ \left[\underline{u}_i^{\lambda_1} (\underline{\nabla u_i})^{\lambda_2}, \overline{u}_i^{\lambda_1} (\overline{\nabla u_i})^{\lambda_2} \right] & \text{if } \lambda_2 \text{ is even.} \end{cases} \tag{13}$$

By (12), (13) and Lemma 3, it follows that

$$\begin{aligned}
 & \sum_{i=n}^{m-1} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \\
 &= \sum_{i=n}^{m-1} \left\| \left[\min \{ \underline{u}_i^{\lambda_1} (\nabla \underline{u}_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\nabla \overline{u}_i)^{\lambda_2} \}, \max \{ \underline{u}_i^{\lambda_1} (\nabla \underline{u}_i)^{\lambda_2}, \overline{u}_i^{\lambda_1} (\nabla \overline{u}_i)^{\lambda_2} \} \right] \right\| \\
 &= \sum_{i=n}^{m-1} \max \left\{ \left| \underline{u}_i^{\lambda_1} (\nabla \underline{u}_i)^{\lambda_2} \right|, \left| \overline{u}_i^{\lambda_1} (\nabla \overline{u}_i)^{\lambda_2} \right| \right\} \\
 &= \max \left\{ \sum_{i=n}^{m-1} \left| \underline{u}_i^{\lambda_1} (\nabla \underline{u}_i)^{\lambda_2} \right|, \sum_{i=n}^{m-1} \left| \overline{u}_i^{\lambda_1} (\nabla \overline{u}_i)^{\lambda_2} \right| \right\} \\
 &\leq \frac{\lambda_2(m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \max \left\{ \sum_{i=n}^m \left| \nabla \underline{u}_i \right|^{\lambda_1 + \lambda_2}, \sum_{i=n}^m \left| \nabla \overline{u}_i \right|^{\lambda_1 + \lambda_2} \right\} \\
 &\leq \frac{\lambda_2(m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=n}^m \max \left\{ \left| \nabla \underline{u}_i \right|^{\lambda_1 + \lambda_2}, \left| \nabla \overline{u}_i \right|^{\lambda_1 + \lambda_2} \right\} \\
 &\leq \frac{\lambda_2(m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=n}^m \left\| \nabla u_i \right\|^{\lambda_1 + \lambda_2}.
 \end{aligned}$$

This concludes the proof. \square

EXAMPLE 2. Suppose that

$$\{u_i\}_{i=1}^n = \left\{ [1, 2], \left[\frac{1}{2}, 1 \right], \left[\frac{1}{i}, \frac{2}{i} \right], \dots, \left[\frac{1}{n-1}, \frac{2}{n-1} \right], [0, 0] \right\}$$

and $\lambda_1 = 1$ and $\lambda_2 = 2$. By induction on n , we have

$$\begin{aligned}
 \sum_{i=2}^{n-1} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| &= \sum_{i=2}^{n-1} \left\| \left[\frac{1}{i}, \frac{2}{i} \right] \cdot \left[\frac{-2}{i(i-1)}, \frac{-1}{i(i-1)} \right]^2 \right\| \\
 &= \sum_{i=2}^{n-1} \left\| \left[\frac{1}{i}, \frac{2}{i} \right] \cdot \left[\frac{1}{i^2(i-1)^2}, \frac{4}{i^2(i-1)^2} \right] \right\| \\
 &= \sum_{i=2}^{n-1} \frac{8}{i^3(i-1)^2} \\
 &\leq \frac{2(n-1)}{3} \sum_{i=2}^n \frac{2^3}{i^3(i-1)^3} \\
 &= \frac{\lambda_2(n-1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=2}^n \left\| \nabla u_i \right\|^{\lambda_1 + \lambda_2}.
 \end{aligned}$$

Theorem 3 is a special case of our next Theorem 5.

THEOREM 5. *Let $\{u_i\}_{i=1}^n$ be a sequence of intervals, $u_0 = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If $\{u_i\}_{i=1}^n$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$, then*

$$\sum_{i=1}^n \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \tag{14}$$

Proof. First, suppose that there exists a finite number of points such that

$$1 = i_0 \leq i_1 < i_2 < \dots < i_{k-1} < i_k = n$$

and u_i is piecewise alternate monotone and piecewise alternate μ -monotone. By Lemma 1, we have

$$\begin{aligned} & \sum_{i=1}^n \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \\ &= \sum_{i=i_0}^{i_1} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| + \sum_{i=i_1+1}^{i_2} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| + \dots + \sum_{i=i_{k-1}+1}^{i_k} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \\ &= \sum_{j=0}^{k-1} \sum_{i=i_j}^{i_{j+1}} \max \left\{ \left| \underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2} \right|, \left| \overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2} \right| \right\} \\ &= \sum_{i=1}^n \max \left\{ \left| \underline{u}_i^{\lambda_1} (\underline{\nabla} u_i)^{\lambda_2} \right|, \left| \overline{u}_i^{\lambda_1} (\overline{\nabla} u_i)^{\lambda_2} \right| \right\} \\ &\leq \frac{\lambda_2(n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^n \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \end{aligned}$$

The proof is complete. \square

Similarly, we can also generalize Theorem 4 as follows.

THEOREM 6. *Let $\{u_i\}_{i=0}^m$ be a sequence of intervals, $u_m = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If $\{u_i\}_{i=0}^m$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$, then*

$$\sum_{i=n}^{m-1} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \leq \frac{\lambda_2(m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=n}^m \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \tag{15}$$

Proof. The proof is analogous to the one of Theorem 5. \square

As an application of Theorems 5 and 6, we now obtain the following result.

THEOREM 7. *Let $\{u_i\}_{i=0}^m$ be a sequence of intervals, $u_0 = u_m = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If $\{u_i\}_{i=0}^m$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$, then*

$$\sum_{i=1}^{m-1} \left\| u_i^{\lambda_1} (\nabla u_i)^{\lambda_2} \right\| \leq \frac{\lambda_2 \left(\left[\frac{m}{2} \right] + 1 \right)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^m \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \tag{16}$$

Proof. Let us take $n = \lceil \frac{m}{2} \rceil$. By Theorem 5, we have

$$\sum_{i=1}^{\lceil \frac{m}{2} \rceil} \|u_i^{\lambda_1} (\nabla u_i)^{\lambda_2}\| \leq \frac{\lambda_2 \left(\lceil \frac{m}{2} \rceil + 1\right)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^{\lceil \frac{m}{2} \rceil} \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \tag{17}$$

Similarly, by Theorem 6, we have

$$\begin{aligned} \sum_{i=\lceil \frac{m}{2} \rceil + 1}^{m-1} \|u_i^{\lambda_1} (\nabla u_i)^{\lambda_2}\| &\leq \frac{\lambda_2 \left(m - \lceil \frac{m}{2} \rceil\right)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=\lceil \frac{m}{2} \rceil + 1}^m \|\nabla u_i\|^{\lambda_1 + \lambda_2} \\ &\leq \frac{\lambda_2 \left(\lceil \frac{m}{2} \rceil + 1\right)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=\lceil \frac{m}{2} \rceil + 1}^m \|\nabla u_i\|^{\lambda_1 + \lambda_2}. \end{aligned} \tag{18}$$

The intended relation (16) follows by adding the above two inequalities (17) and (18). \square

EXAMPLE 3. Suppose that

$$\{u_i\}_{i=0}^5 = \{[0, 0], [1, 2], [2, 4], [3, 6], [1, 2], [0, 0]\}, \lambda_1 = 2, \lambda_2 = 3.$$

Then, we have

$$\sum_{i=1}^4 \|u_i^2 (\nabla u_i)^3\| = 704 < 6048 = \frac{3 \cdot 3^2}{5} \sum_{i=1}^4 \|\nabla u_i\|^5.$$

Let $\lambda_1 = 1, \lambda_2 = 2$. Then, we have

$$\sum_{i=1}^4 \|u_i (\nabla u_i)^2\| = 80 < 184 = \frac{2 \cdot 3}{3} \sum_{i=1}^4 \|\nabla u_i\|^3.$$

Now, we give new discrete Opial inequalities involving two interval sequences.

THEOREM 8. Let $\{u_i\}_{i=0}^n$ and $\{v_i\}_{i=0}^n$ be two sequences of intervals, $u_0 = v_0 = [0, 0]$. If u_i and v_i are synchronous monotone and μ -increasing, then

$$\sum_{i=1}^n \|u_{i-1} \nabla v_i + v_i \nabla u_i\| \leq \frac{n}{2} \sum_{i=1}^n \|(\nabla u_i)^2 + (\nabla v_i)^2\|. \tag{19}$$

Proof. Suppose that u_i and v_i are increasing and μ -increasing. Then, $u_i v_i$ is also increasing and μ -increasing. Consequently, we obtain that

$$\begin{aligned} u_{i-1} \nabla v_i + v_i \nabla u_i &= \left[\underline{u}_{i-1}, \overline{u}_{i-1} \right] \cdot \left[\underline{v}_i - \underline{v}_{i-1}, \overline{v}_i - \overline{v}_{i-1} \right] + \left[\underline{v}_i, \overline{v}_i \right] \cdot \left[\underline{u}_i - \underline{u}_{i-1}, \overline{u}_i - \overline{u}_{i-1} \right] \\ &= \left[\underline{u}_{i-1} (\underline{v}_i - \underline{v}_{i-1}) + \underline{v}_i (\underline{u}_i - \underline{u}_{i-1}), \overline{u}_{i-1} (\overline{v}_i - \overline{v}_{i-1}) + \overline{v}_i (\overline{u}_i - \overline{u}_{i-1}) \right] \\ &= \left[\underline{u}_i \cdot \underline{v}_i - \underline{u}_{i-1} \cdot \underline{v}_{i-1}, \overline{u}_i \cdot \overline{v}_i - \overline{u}_{i-1} \cdot \overline{v}_{i-1} \right] \\ &= \nabla (u_i v_i). \end{aligned}$$

If u_i and v_i are decreasing and μ -increasing, then $u_i v_i$ is increasing and μ -increasing. Consequently, we obtain that

$$\begin{aligned} u_{i-1} \nabla v_i + v_i \nabla u_i &= \left[\underline{u}_{i-1}, \overline{u}_{i-1} \right] \cdot \left[\underline{v}_i - \underline{v}_{i-1}, \overline{v}_i - \overline{v}_{i-1} \right] + \left[\underline{v}_i, \overline{v}_i \right] \cdot \left[\underline{u}_i - \underline{u}_{i-1}, \overline{u}_i - \overline{u}_{i-1} \right] \\ &= \left[\overline{u}_{i-1}(\overline{v}_i - \overline{v}_{i-1}) + \overline{v}_i(\overline{u}_i - \overline{u}_{i-1}), \underline{u}_{i-1}(\underline{v}_i - \underline{v}_{i-1}) + \underline{v}_i(\underline{u}_i - \underline{u}_{i-1}) \right] \\ &= \left[\overline{u}_i \cdot \overline{v}_i - \overline{u}_{i-1} \cdot \overline{v}_{i-1}, \underline{u}_i \cdot \underline{v}_i - \underline{u}_{i-1} \cdot \underline{v}_{i-1} \right] \\ &= \nabla(u_i v_i). \end{aligned}$$

Then, by the Cauchy–Schwarz inequality, we have

$$\begin{aligned} \sum_{i=1}^n \|u_{i-1} \nabla v_i + v_i \nabla u_i\| &= \sum_{i=1}^n \max \left\{ \underline{u}_i \cdot \underline{v}_i - \underline{u}_{i-1} \cdot \underline{v}_{i-1}, \overline{u}_i \cdot \overline{v}_i - \overline{u}_{i-1} \cdot \overline{v}_{i-1} \right\} \\ &\leq \max \left\{ \sum_{i=1}^n \left(\underline{u}_i \cdot \underline{v}_i - \underline{u}_{i-1} \cdot \underline{v}_{i-1} \right), \sum_{i=1}^n \left(\overline{u}_i \cdot \overline{v}_i - \overline{u}_{i-1} \cdot \overline{v}_{i-1} \right) \right\} \\ &\leq \max \left\{ \underline{u}_n \cdot \underline{v}_n, \overline{u}_n \cdot \overline{v}_n \right\} \\ &= \|u_n v_n\| \\ &= \left\| \sum_{i=1}^n \nabla u_i \right\| \cdot \left\| \sum_{i=1}^n \nabla v_i \right\| \\ &\leq \frac{n}{2} \sum_{i=1}^n \left(\|\nabla u_i\|^2 + \|\nabla v_i\|^2 \right). \end{aligned}$$

This concludes the proof. \square

The following results are proved similarly to Theorem 8.

THEOREM 9. *Let $\{u_i\}_{i=0}^m$ and $\{v_i\}_{i=0}^m$ be two sequences of intervals, $u_m = v_m = [0, 0]$. If u_i and v_i are synchronous monotone and μ -decreasing, then*

$$\sum_{i=n+1}^m \|u_{i-1} \nabla v_i + v_i \nabla u_i\| \leq \frac{m-n}{2} \sum_{i=n+1}^m \|(\nabla u_i)^2 + (\nabla v_i)^2\|.$$

THEOREM 10. *Let $\{u_i\}_{i=0}^m$ and $\{v_i\}_{i=0}^m$ be two sequences of intervals, $u_0 = v_0 = [0, 0]$. If $\{u_i\}_{i=0}^m$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$ and $v_i = [0, 0]$, then*

$$\sum_{i=1}^n \|u_{i-1} \nabla v_i + v_i \nabla u_i\| \leq \frac{n}{2} \sum_{i=1}^n \|(\nabla u_i)^2 + (\nabla v_i)^2\|.$$

THEOREM 11. *Let $\{u_i\}_{i=0}^m$ and $\{v_i\}_{i=0}^m$ be two sequences of intervals, $u_m = v_m = [0, 0]$. If $\{u_i\}_{i=0}^m$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$ and $v_i = [0, 0]$, then*

$$\sum_{i=n+1}^m \|u_{i-1} \nabla v_i + v_i \nabla u_i\| \leq \frac{m-n}{2} \sum_{i=n+1}^m \|(\nabla u_i)^2 + (\nabla v_i)^2\|.$$

THEOREM 12. Let $\{u_i\}_{i=0}^m$ and $\{v_i\}_{i=0}^m$ be two sequences of intervals, $u_1 = v_1 = [0, 0]$, and $u_m = v_m = [0, 0]$. If $\{u_i\}_{i=0}^m$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$ and $v_i = [0, 0]$, then

$$\sum_{i=1}^m \|u_{i-1} \nabla v_i + v_i \nabla u_i\| \leq \frac{\lceil \frac{m+1}{2} \rceil}{2} \sum_{i=1}^m \|(\nabla u_i)^2 + (\nabla v_i)^2\|.$$

4. Opial type inequalities involving the forward/delta gH-difference operator

In Section 3, we obtained several Opial type inequalities involving the backward gH-difference operator. Similar arguments can be used to establish discrete Opial type inequalities concerning the forward gH-difference operator. The proofs of the results formulated here are left to the interested reader.

THEOREM 13. (delta version of Theorem 3) Let $\{u_i\}_{i=1}^n$ be a sequence of intervals, $u_0 = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If u_i is monotone and μ -increasing, then

$$\sum_{i=0}^{n-1} \|u_i^{\lambda_1} (\Delta u_i)^{\lambda_2}\| \leq \frac{\lambda_2 (n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=0}^{n-1} \|\Delta u_i\|^{\lambda_1 + \lambda_2}.$$

THEOREM 14. (delta version of Theorem 4) Let $\{u_i\}_{i=0}^m$ be a sequence of intervals, $u_m = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If u_i is monotone and μ -decreasing, then

$$\sum_{i=n}^{m-1} \|u_i^{\lambda_1} (\Delta u_i)^{\lambda_2}\| \leq \frac{\lambda_2 (m-n+1)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=n}^m \|\Delta u_i\|^{\lambda_1 + \lambda_2}.$$

THEOREM 15. (delta version of Theorem 7) Let $\{u_i\}_{i=0}^m$ be a sequence of intervals, $u_0 = u_m = [0, 0]$, and $\lambda_1, \lambda_2 \geq 1$. If $\{u_i\}_{i=0}^m$ is piecewise alternate monotone, piecewise alternate μ -monotone, and there is no other point i such that $u_i = [0, 0]$, then

$$\sum_{i=1}^{m-1} \|u_i^{\lambda_1} (\Delta u_i)^{\lambda_2}\| \leq \frac{\lambda_2 \left(\lceil \frac{m}{2} \rceil + 1\right)^{\lambda_1}}{\lambda_1 + \lambda_2} \sum_{i=1}^m \|\Delta u_i\|^{\lambda_1 + \lambda_2}.$$

5. Conclusions

We investigated discrete Opial type inequalities for interval-valued functions, and obtained several new interval discrete Opial type inequalities. Our results generalize many known discrete Opial type inequalities, and will be useful in developing the theory of interval difference inequalities and interval difference equations. As future research directions, we intend to investigate interval discrete Opial type inequalities on time scales, and give some applications to interval difference equations.

Acknowledgements. The authors would like to thank the anonymous reviewers and editors for their truly constructive comments, which helped them to improve the quality and clarity of the manuscript. This research is supported by Natural Science Foundation of Hubei Province (2023AFD013), Philosophy and Social Sciences of Educational Commission of Hubei Province of China (22Y109), and Foundation of Hubei Normal University (2022055). Torres is supported by the Portuguese Foundation for Science and Technology (FCT), project UIDB/04106/2020 (CIDMA).

REFERENCES

- [1] R. P. AGARWAL, M. BOHNER AND A. ÖZBEKLER, *Lyapunov inequalities and applications*, Springer, Cham, 2021.
- [2] R. P. AGARWAL, V. ČULJAK AND J. PEČARIĆ, *On discrete and continuous Wirtinger inequalities*, *Appl. Anal.*, **70** (1998), 195–204.
- [3] R. P. AGARWAL AND P. Y. H. PANG, *Opial inequalities with applications in differential and difference equations*, Kluwer Academic Publishers, Dordrecht, 1995.
- [4] R. P. AGARWAL AND P. Y. H. PANG, *Discrete Opial-type inequalities involving higher order partial differences*, *Nonlinear Anal.*, **27** (1996), 429–454.
- [5] M. ANDRIĆ, J. PEČARIĆ AND I. PERIĆ, *On weighted integral and discrete Opial-type inequalities*, *Math. Inequal. Appl.*, **19** (2016), 1295–1307.
- [6] H. D. BLOCK, *Discrete analogues of certain integral inequalities*, *Proc. Amer. Math. Soc.*, **8** (1957), 852–859.
- [7] M. J. BOHNER, R. R. MAHMOUD AND S. H. SAKER, *Discrete, continuous, delta, nabla, and diamond-alpha Opial inequalities*, *Math. Inequal. Appl.*, **18** (2015), 923–940.
- [8] Y. CHALCO-CANO, A. FLORES-FRANULIĆ AND H. ROMÁN-FLORES, *Ostrowski type inequalities for interval-valued functions using generalized Hukuhara derivative*, *Comput. Appl. Math.*, **31** (2012), 457–472.
- [9] Y. CHALCO-CANO, W. A. LODWICK AND W. CONDORI-EQUICE, *Ostrowski type inequalities and applications in numerical integration for interval-valued functions*, *Soft Comput.*, **19** (2015), 3293–3300.
- [10] W. S. CHEUNG, *Sharp discrete inequalities and applications to discrete variational problems*, *J. Comput. Appl. Math.*, **232** (2009), 176–186.
- [11] W. S. CHEUNG AND J. L. REN, *Discrete non-linear inequalities and applications to boundary value problems*, *J. Math. Anal. Appl.*, **319** (2006), 708–724.
- [12] P. CERONE AND S. S. DRAGOMIR, *Mathematical inequalities. A perspective*, CRC Press, Boca Raton, FL, 2011.
- [13] T. M. COSTA, *Jensen's inequality type integral for fuzzy-interval-valued functions*, *Fuzzy Sets Syst.*, **327** (2017), 31–47.
- [14] T. M. COSTA AND H. ROMÁN-FLORES, *Some integral inequalities for fuzzy-interval-valued functions*, *Inform. Sci.*, **420** (2017), 110–125.
- [15] T. M. COSTA, H. ROMÁN-FLORES AND Y. CHALCO-CANO, *Opial-type inequalities for interval-valued functions*, *Fuzzy Sets Syst.*, (2019), 48–63.
- [16] S. S. DRAGOMIR, *Discrete inequalities of the Cauchy-Bunyakovsky-Schwarz type*, Nova Science Publishers, Inc., Hauppauge, NY, 2004.
- [17] S. S. DRAGOMIR, *Discrete inequalities of Jensen type for λ -convex functions on linear spaces*, *Rend. Istit. Mat. Univ. Trieste*, **47** (2015), 241–265.
- [18] S. S. DRAGOMIR, *Recent developments of discrete inequalities for convex functions defined on linear spaces with applications*, *Modern discrete mathematics and analysis*, 117–172, Springer Optim. Appl., 131, Springer, Cham, 2018.
- [19] K. FAN, O. TAUSSKY AND J. TODD, *Discrete analogs of inequalities of Wirtinger*, *Monatsh. Math.*, **59** (1955), 73–90.
- [20] Q. H. FENG, *Some new generalized Gronwall-Bellman type discrete fractional inequalities*, *Appl. Math. Comput.*, **259** (2015), 403–411.

- [21] A. FLORES-FRANULIĆ, Y. CHALCO-CANO AND H. ROMÁN-FLORES, *An Ostrowski type inequality for interval-valued functions*, IFSA World Congress and NAFIPS Annual Meeting IEEE, **35** (2013), 1459–1462.
- [22] A. HAMIAZ AND W. ABUELELA, *Some new discrete Hilbert's inequalities involving Fenchel-Legendre transform*, J. Inequal. Appl., **39** (2020), 14 pp.
- [23] X. HAN, S. LI AND Q. L. LI, *Some new discrete inequalities of Opial with two sequences*, Ann. Appl. Math., **34** (2018), 376–382.
- [24] G. H. HARDY, J. E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge University Press, 1934.
- [25] M. HUKUHARA, *Integration des applications mesurables dont la valeur est un compact convexe*, Funkcial. Ekvac., **10** (1967), 205–223.
- [26] M. KRNIĆ AND P. VUKOVIĆ, *On a more accurate class of discrete Hilbert-type inequalities*, Appl. Math. Comput., **234** (2014), 543–547.
- [27] C.-M. LEE, *On a discrete analogue of inequalities of Opial and Yang*, Canad. Math. Bull., **11** (1968), 73–77.
- [28] Q. H. MA, *Some new nonlinear Volterra-Fredholm-type discrete inequalities and their applications*, J. Comput. Appl. Math., **216** (2008), 451–466.
- [29] Q. H. MA, *Estimates on some power nonlinear Volterra-Fredholm type discrete inequalities and their applications*, J. Comput. Appl. Math., **233** (2010), 2170–2180.
- [30] F. W. MENG AND D. H. JI, *On some new nonlinear discrete inequalities and their applications*, J. Comput. Appl. Math., **208** (2007), 425–433.
- [31] F. W. MENG AND W. N. LI, *On some new nonlinear discrete inequalities and their applications*, J. Comput. Appl. Math., **158** (2003), 407–417.
- [32] K. NIKODEM, J. L. SÁNCHEZ AND L. SÁNCHEZ, *Jensen and Hermite-Hadamard inequalities for strongly convex set-valued maps*, Math. Aeterna., **4** (2014), 979–987.
- [33] Z. OPIAL, *Sur une inégalité*, Ann. Polon. Math., **8** (1960), 29–32.
- [34] B. G. PACHPATTE, *A note on Opial and Wirtinger type discrete inequalities*, J. Math. Anal. Appl., **127** (1987), 470–474.
- [35] B. G. PACHPATTE, *Opial type discrete inequalities in two variables*, Tamkang J. Math., **22** (1991), 323–328.
- [36] B. G. PACHPATTE, *A note on new discrete Chebyshev-Grüss type inequalities*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.), **54** (2008), 117–121.
- [37] Y. M. QIN, *Integral and discrete inequalities and their applications*, Vol. I. Linear inequalities. Birkhäuser/Springer, [Cham], 2016.
- [38] H. ROMÁN-FLORES, Y. CHALCO-CANO AND W. A. LODWICK, *Some integral inequalities for interval-valued functions*, Comput. Appl. Math., **37** (2018), 1306–1318.
- [39] S. H. SAKER AND R. P. AGARWAL, *Discrete Hardy's type inequalities and structure of discrete class of weights satisfy reverse Hölder's inequality*, Math. Inequal. Appl., **24** (2021), 521–541.
- [40] S. H. SAKER, M. KRNIĆ AND J. PEČARIĆ, *Higher summability theorems from the weighted reverse discrete inequalities*, Appl. Anal. Discrete Math., **13** (2019), 423–439.
- [41] S. H. SAKER AND I. KUBIACZYK, *Higher summability and discrete weighted Muckenhoupt and Gehring type inequalities*, Proc. Edinb. Math. Soc., **62** (2019), 949–973.
- [42] E. SET AND M. Z. SARIKAYA, *On the generalization of Ostrowski and Grüss type discrete inequalities*, Comput. Math. Appl., **62** (2011), 455–461.
- [43] L. STEFANINI, *A generalization of Hukuhara difference and division for interval and fuzzy arithmetic*, Fuzzy Sets Syst., **161** (2010), 1564–1584.
- [44] M. ŠTRBOJA, T. GRBIĆ, I. ŠTAJNER-PAPUGA, G. GRUJIĆ AND S. MEDIĆ, *Jensen and Chebyshev inequalities for pseudo-integrals of set-valued functions*, Fuzzy Sets Syst., **222** (2013), 18–32.
- [45] B. C. YANG, *On more accurate reverse multidimensional half-discrete Hilbert-type inequalities*, Math. Inequal. Appl., **18** (2015), 589–605.
- [46] D. L. ZHANG, C. M. GUO, D. G. CHEN AND G. J. WANG, *Jensen's inequalities for set-valued and fuzzy set-valued functions*, Fuzzy Sets Syst., **404** (2021), 178–204.
- [47] D. L. ZHANG, C. M. GUO, D. G. CHEN AND G. J. WANG, *Choquet integral Jensen's inequalities for set-valued and fuzzy set-valued functions*, Soft Comput., **25** (2021), 903–918.
- [48] D. F. ZHAO, T. Q. AN, G. J. YE AND W. LIU, *New Jensen and Hermite-Hadamard type inequalities for h -convex interval-valued functions*, J. Inequal. Appl., **2018** (2018): 302, 14 pp.

- [49] D. F. ZHAO, T. Q. AN, G. J. YE AND W. LIU, *Chebyshev type inequalities for interval-valued functions*, *Fuzzy Sets Syst.*, **396** (2020), 82–101.
- [50] D. F. ZHAO, T. Q. AN, G. J. YE AND W. LIU, *Some generalizations of Opial type inequalities for interval-valued functions*, *Fuzzy Sets Syst.*, **436** (2022), 128–151.
- [51] D. F. ZHAO, G. J. YE, W. LIU AND D. F. M. TORRES, *Some inequalities for interval-valued functions on time scales*, *Soft Comput.*, **23** (2019), 6005–6015.

(Received May 18, 2022)

Dafang Zhao
School of Mathematics and Statistics
Hubei Normal University
Huangshi 435002, P. R. China
e-mail: dafangzhao@163.com

Xuexiao You
School of Mathematics and Statistics
Hubei Normal University
Huangshi 435002, P. R. China
e-mail: youxuexiao@126.com

Delfim F. M. Torres
Center for Research and Development in Mathematics and
Applications (CIDMA)
Department of Mathematics, University of Aveiro
3810-193 Aveiro, Portugal
e-mail: delfim@ua.pt