

Whittaker differential equation associated to the initial heat problem*

M. M. Rodrigues and S. Saitoh

Abstract. In this paper, by using the theory of reproducing kernels, we investigate integral transforms with kernels related to the solutions of the initial Whittaker heat problem.

Mathematics Subject Classification (2010). Primary 42A38; Secondary 32A30, 4E22.

Keywords. Integral transform, isometric mapping, inversion formula, initial value problem, linear integral equation, regularization, eigenfunction, eigenvalue, reproducing kernel, Whittaker function.

1. Introduction

The Whittaker functions are closely related to the confluent hypergeometric functions which play an important role in various branches of applied mathematics and theoretical physics, for instance, fluid mechanics scalar and electromagnetic diffraction theory, atomic structure theory, input-output situations and storage consumption situations in economic problems, and so on. Moreover, they have acquired an ever increasing significance due to their frequent use in applications of mathematics to physical and technical problems [2] [3]. This justifies the continuous effort in studying properties of these functions and in gathering information about them.

Let consider the general method given in [5] for the existence and construction of the solution of the following initial problem

$$(\partial_t + L_x)u_f(t, x) = 0, \quad t > 0 \quad (1.1)$$

satisfying the initial value condition

$$u_f(0, x) = f(x), \quad (1.2)$$

*The final version is published is published in chapter IX in the book *Current Trends in Analysis and Its Applications*, 523–530, 2015. It as available via the website http://link.springer.com/chapter/10.1007%2F978-3-319-12577-0_58#page-1

for some general linear operator L_x on some function space, and on some domain, by using the theory of reproducing kernels.

We shall investigate the integral transforms with kernels related to the solutions of the equations by using the theory of reproducing kernels, for the Whittaker heat equations.

The Whittaker functions denoted by $M_{\mu,\nu}(x)$, $W_{\mu,\nu}(x)$ arise as solutions to the Whittaker differential equation, i.e., are solutions of the linear homogeneous ordinary differential equation of the second order

$$\frac{d^2W}{dx^2} + \left(-\frac{1}{4} + \frac{\mu}{x} + \frac{\frac{1}{4} - \nu^2}{x^2} \right) W = 0. \quad (1.3)$$

Here, we will deal with the Whittaker function $W_{\mu,\nu}$ which is an eigenfunction of a second order differential operator

$$A_x W_{\mu,\nu}(x) = 4\nu^2 W_{\mu,\nu}(x) \quad (1.4)$$

with a eigenvalue $4\nu^2$, and where

$$A_x = 4x^2 \frac{d^2}{dx^2} - x^2 + 4\mu x + 1. \quad (1.5)$$

Further, we consider the case that ν are reals, and the functions

$$\exp(-4\nu^2 t) W_{\mu,\nu}(x) \quad (1.6)$$

are the solutions of the operator equation

$$(\partial_t + A_x) u(t, x) = 0. \quad (1.7)$$

We shall consider some general solutions of (1.7) by a suitable sum of the solutions (1.6). In order to consider a fully general sum, we shall consider the kernel form for a nonnegative continuous function ρ ,

$$\mathcal{K}_t(x, y; \rho) = \int_0^{+\infty} \exp\{-4\nu^2 t\} W_{i\nu,\nu}(x) \overline{W_{i\nu,\nu}(y)} \rho(\tau) d\tau. \quad (1.8)$$

Of course, here, we are considering the integral with absolutely convergence for the kernel form.

The fully general solutions of the equation (1.7) may be represented in the integral form

$$u(t, x) = \int_0^{+\infty} \exp\{-4\nu^2 t\} W_{i\nu,\nu}(x) F(\tau) \rho(\tau) d\tau. \quad (1.9)$$

for the functions F satisfying

$$\int_0^{+\infty} \exp\{-4\nu^2 t\} |F(\tau)|^2 \rho(\tau) d\tau < \infty. \quad (1.10)$$

Then, the solution $u(t, x)$ of (1.7) satisfying the initial condition

$$u(0, x) = F(x) \quad (1.11)$$

will be obtained by $t \rightarrow 0$ in (1.9) with a natural meaning. However, this point will be very delicate and we will need to consider some deep and beautiful structure. Here, (1.8) is a reproducing kernel and in order to analyze the

logic above, we will need the theory of reproducing kernels, essentially and beautiful ways. Indeed, in order to construct natural solutions (1.8) we will need a new framework and function space.

2. Main results

In order to analyze the integral transform (1.11), we will need the essence of the theory of reproducing kernels. We are interesting in the integral transforms (1.11) in the framework of Hilbert spaces. Of course, we are interesting in the characterization of the image functions, the isometric identity like the Parseval identity and the inversion formula, basically. For these general and fundamental problems, we have a unified and fundamental method and concept in the general situation as follows in [6], [7] and [8], where we can find the general theory for linear mappings in the framework of Hilbert spaces.

Moreover, recently, we obtained a very general inversion formula based on the Aveiro Discretization Method in Mathematics ([4]) by using the ultimate realization of reproducing kernel Hilbert spaces. Following that the general theory, we shall build our result.

For this purpose, we need to recall that for $k \neq \frac{1}{2}$ the Whittaker function $W_{i\tau, k-\frac{1}{2}}$ is an eigenfunction of a second order differential, i.e.

$$A_x W_{i\tau, k-\frac{1}{2}}(x) = 4 \left(k - \frac{1}{2} \right)^2 W_{i\tau, k-\frac{1}{2}}(x), \quad x > 0 \quad (2.1)$$

where

$$A_x = 4x^2 \frac{d^2}{dx^2} - x^2 + 4i\tau x + 1. \quad (2.2)$$

Then, we form the reproducing kernel

$$\mathcal{K}(x, y; \rho) = \int_0^{+\infty} W_{i\tau, k-\frac{1}{2}}(x) \overline{W_{i\tau, k-\frac{1}{2}}(y)} \rho(\tau) d\tau, \quad t > 0, \quad (2.3)$$

and consider the reproducing kernel Hilbert space $H_{\mathcal{K}(\rho)}(\mathbb{R}^+)$ admitting the kernel $\mathcal{K}(x, y; \rho)$. In particular, note that

$$\mathcal{K}_t(x, y; \rho) \in H_{\mathcal{K}(\rho)}(\mathbb{R}^+), \quad y > 0.$$

Then, we obtain the main theorem in this paper:

Theorem 2.1 (Main Theorem). *For any member $f \in H_{\mathcal{K}(\rho)}(\mathbb{R}^+)$, the solution $u_f(t, x)$ of the initial value problem, for $t > 0$*

$$(\partial_t - A_x)u_f(t, x) = 0 \quad (2.4)$$

satisfying the initial value condition

$$u_f(0, x) = f(x), \quad (2.5)$$

exists and it is given by

$$u_f(t, x) = (f(\cdot), \mathcal{K}_t(\cdot, x; \rho))_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)}. \quad (2.6)$$

Here, the meaning of the initial value (2.5) is given by

$$\begin{aligned} \lim_{t \rightarrow +0} u_f(t, x) &= \lim_{t \rightarrow +0} (f(\cdot), \mathcal{K}_t(\cdot, x\rho))_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)} \\ &= (f(\cdot), \mathcal{K}(\cdot, x\rho))_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)} \\ &= f(x), \end{aligned} \quad (2.7)$$

whose existence is, in general, ensured and the limit is the uniformly convergence on any subset of \mathbb{R}^+ such that $\mathcal{K}(x, x; \rho)$ is bounded.

The uniqueness property of the initial value problem is depending on the completeness of the family of functions

$$\{\mathcal{K}_t(\cdot, x; \rho) : x \in \mathbb{R}^+\} \quad (2.8)$$

in $H_{\mathcal{K}(\rho)}(\mathbb{R}^+)$.

In our theorem, the complete property of the solutions $u_f(t, x)$ of (2.4) and (2.5) satisfying the initial value f may be derived by the reproducing kernel Hilbert space admitting the kernel

$$k(x, t; y, \tau; \rho) := (\mathcal{K}_\tau(\cdot, y; \rho), \mathcal{K}_t(\cdot, x; \rho))_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)}. \quad (2.9)$$

In our method, we see that the existence problem of the initial value problem is based on the eigenfunctions and we are constructing the desired solution satisfying the desired initial condition. For a larger knowledge for the eigenfunctions we can consider a more general initial value problem.

Furthermore, by considering the linear mapping of (2.6) with various situations, we will be able to obtain various inverse problems looking for the initial values f from the various out put data of $u_f(t, x)$.

3. Proof of the Theorem

The first, note that the kernel $\mathcal{K}_t(x, y; \rho)$ satisfies the operator equation (2.4) for any fixed y , because the functions

$$\exp \left\{ -4 \left(k - \frac{1}{2} \right)^2 t \right\} \overline{W_{-i\tau, k - \frac{1}{2}}(x)}$$

satisfy the operator equation and it is the summation. Similarly, the function $u_f(t, x)$ defined by (2.6) is the solution of the operator equation (2.4).

In order to see the initial value problem, we note the important general properties

$$\mathcal{K}_t(x, y; \rho) \ll \mathcal{K}(x, y; \rho); \quad (3.1)$$

that is, $\mathcal{K}(x, y; \rho) - \mathcal{K}_t(x, y; \rho)$ is a positive definite quadratic form function and we have

$$H_{\mathcal{K}_t(\rho)} \subset H_{\mathcal{K}(\rho)}(\mathbb{R}^+)$$

and for any function $f \in H_{\mathcal{K}_t(\rho)}$

$$\|f\|_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)} = \lim_{t \rightarrow +0} \|f\|_{H_{\mathcal{K}_t(\rho)}}$$

in the sense of non-decreasing norm convergence ([1]). In order to see the crucial point in (2.7), note that

$$\begin{aligned} & \|\mathcal{K}(x, y; \rho) - \mathcal{K}_t(x, y; \rho)\|_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)}^2 \\ &= \mathcal{K}(y, y; \rho) - 2\mathcal{K}_t(y, y; \rho) + \|\mathcal{K}_t(x, y; \rho)\|_{H_{\mathcal{K}(\rho)}(\mathbb{R}^+)}^2 \\ &\leq \mathcal{K}(y, y; \rho) - 2\mathcal{K}_t(y, y; \rho) + \|\mathcal{K}_t(x, y; \rho)\|_{H_{\mathcal{K}_t(\rho)}(\mathbb{R}^+)}^2 \\ &= \mathcal{K}(y, y; \rho) - \mathcal{K}_t(y, y; \rho), \end{aligned}$$

that converges to zero as $t \rightarrow +0$. We thus obtain the desired limit property in the theorem.

The uniqueness property of the initial value problem follows from (2.6) easily.

In the main theorem, the realization of the reproducing kernel Hilbert space $H_{\mathcal{K}(\rho)}(\mathbb{R}^+)$ is a crucial point, for this purpose, we are interested in the calculation of the kernel $\mathcal{K}(x, y; \rho)$.

4. Concrete realization of the reproducing kernel Hilbert spaces

As the theory of reproducing kernels, their realizations will give interesting research topics that are requested separate papers. So, here, we shall discuss the following concrete form of the reproducing kernels. From [2], we have

$$\begin{aligned} & \int_0^{+\infty} |\Gamma(k + i\tau)|^2 W_{i\tau, k - \frac{1}{2}}(x) \overline{W_{i\tau, k - \frac{1}{2}}(y)} d\tau \\ &= \sqrt{\pi} \Gamma(2k) (xy)^k (x + y)^{-2k+1} K_{2k - \frac{1}{2}}\left(\frac{x + y}{2}\right), \end{aligned} \quad (4.1)$$

where $K_{2k - \frac{1}{2}}(z)$ denotes the modified Bessel function. Furthermore, since k, x are reals then $\overline{W_{i\tau, k - \frac{1}{2}}(x)} = W_{-i\tau, k - \frac{1}{2}}(x)$.

Moreover, note that:

$$W_{\mu, \nu}(0) = 0, \quad -\frac{1}{2} < \operatorname{Re}(\nu) < \frac{1}{2},$$

$$W_{\mu, \nu}(z) \sim z^\mu e^{-\frac{z}{2}} \left(1 + O\left(\frac{1}{z}\right)\right), \quad |z| \rightarrow +\infty$$

We were able to realize the important reproducing kernel Hilbert space concretely and analytically. Meanwhile, we are also interested in the kernel forms \mathcal{K}_t and K . These calculations will create a new and large field in integral formulas.

We have to analyze and realize the corresponding reproducing kernel Hilbert spaces, however, we can apply quite general formula by the Aveiro discretization method in ([4]).

Proposition 4.1. (Ultimate realization of reproducing kernel Hilbert spaces). *In our general situation and for a uniqueness set $\{p_j\}$ for the reproducing*

kernel Hilbert space H_K of the set E satisfying the linearly independence of $K(\cdot, p_j)$ for any finite number of the points p_j , we obtain

$$\|f\|_{H_K}^2 = \|\mathbf{f}^*\|_{\mathcal{H}}^2 = \lim_{n \rightarrow \infty} \sum_j \sum_{j'} f(p_j) \overline{a_{jj'}} \overline{f(p_{j'})}. \quad (4.2)$$

In this Proposition, for the uniqueness set of the space, if the reproducing kernel is analytical, then, the criteria will be very simple by the *identity theorem of analytic functions*. For the Sovelev space cases, we have to consider some dense subset of E for the uniqueness set. Meanwhile, the linearly independence will be easily derived from the integral representations of the kernels.

Acknowledgment

The authors were supported by *FEDER* funds through *COMPETE*–Operational Programme Factors of Competitiveness (“Programa Operacional Factores de Competitividade”) and by Portuguese funds through the *Center for Research and Development in Mathematics and Applications* (University of Aveiro) and the Portuguese Foundation for Science and Technology (“FCT–Fundação para a Ciência e a Tecnologia”), within project PEst-C/MAT/UI4106/2011 with *COMPETE* number FCOMP-01-0124-FEDER-022690. The second author is supported in part by the Grant-in-Aid for the Scientific Research (C)(2)(No. 24540113)

References

- [1] N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc., **68** (1950), 337–404.
- [2] H. Buchholz, *The confluent hypergeometric function with special emphasis on its applications*, Springer-Verlag, Berlin, Heidelberg, New York, 1969.
- [3] Erdélyi A, Magnus W, Oberhettinger F, Tricomi FG. 1954. *Tables of Integral Transforms-Vol I, II*. Bateman Manuscript Project. California Institute of Technology, McGraw Hill, New York.
- [4] L. P. Castro, H. Fujiwara, M.M. Rodrigues, S. Saitoh and V.K. Tuan, *Aveiro Discretization Method in Mathematics: A New Discretization Principle*, MATHEMATICS WITHOUT BOUNDARIES : SURVEYS IN PURE MATHEMATICS, Edited by Panos Pardalos and Themistocles M. Rassias (to appear)(Springer). 52pages.
- [5] L. P. Castro, M.M. Rodrigues, S. Saitoh, *A fundamental theorem on linear operator equations using the theory of reproducing kernels*, submitted.
- [6] S. Saitoh, *Hilbert spaces induced by Hilbert space valued functions*, Proc. Amer. Math. Soc., 89 (1983), 74–78.
- [7] S. Saitoh, *Integral transforms, reproducing kernels and their applications*, Pitman Research Notes in Mathematics Series 369, Addison Wesley Longman, Harlow, 1997.
- [8] S. Saitoh, *Theory of reproducing kernels: Applications to approximate solutions of bounded linear operator functions on Hilbert spaces*, Amer. Math. Soc. Transl. Ser., 230, Amer. Math. Soc., Providence, RI, 2010.

- [9] E.C. Titchmarsh, *Eigenfunction Expansions Associated With Second-Order Differential Equations*, Part I Clarendon Press, Oxford, 1946.
- [10] J. Wimp, *A class of integral transforms*, Proc. Edinburgh Math Soc., **14(1)**, (1964) 33-44.

M. M. Rodrigues

Center for Research and Development in Mathematics and Applications
Department of Mathematics, University of Aveiro
3810-193 Aveiro
Portugal
e-mail: mrodrigues@ua.pt

S. Saitoh

Center for Research and Development in Mathematics and Applications
Department of Mathematics, University of Aveiro
3810-193 Aveiro
Portugal
e-mail: saburou.saitoh@gmail.com