

Accepted Manuscript

On the multiplicity of α as an eigenvalue of $A_\alpha(G)$ of graphs with pendant vertices

Domingos M. Cardoso, Germain Pastén, Oscar Rojo

PII: S0024-3795(18)30199-X
DOI: <https://doi.org/10.1016/j.laa.2018.04.013>
Reference: LAA 14553

To appear in: *Linear Algebra and its Applications*

Received date: 2 January 2017
Accepted date: 12 April 2018

Please cite this article in press as: D.M. Cardoso et al., On the multiplicity of α as an eigenvalue of $A_\alpha(G)$ of graphs with pendant vertices, *Linear Algebra Appl.* (2018), <https://doi.org/10.1016/j.laa.2018.04.013>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



On the multiplicity of α as an eigenvalue of $A_\alpha(G)$ of graphs with pendant vertices

Domingos M. Cardoso

Department of Mathematics, Universidade de Aveiro, Aveiro, Portugal

Germain Pastén, Oscar Rojo

Department of Mathematics, Universidad Católica del Norte, Antofagasta, Chile.

Abstract

Let G be a simple undirected graph. Let $0 \leq \alpha \leq 1$. Let

$$A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$$

where $D(G)$ and $A(G)$ are the diagonal matrix of the vertex degrees of G and the adjacency matrix of G , respectively. Let $p(G) > 0$ and $q(G)$ be the number of pendant vertices and quasi-pendant vertices of G , respectively. Let $m_G(\alpha)$ be the multiplicity of α as an eigenvalue of $A_\alpha(G)$. It is proved that

$$m_G(\alpha) \geq p(G) - q(G)$$

with equality if each internal vertex is a quasi-pendant vertex. If there is at least one internal vertex which is not a quasi-pendant vertex, the equality

$$m_G(\alpha) = p(G) - q(G) + m_N(\alpha)$$

is determined in which $m_N(\alpha)$ is the multiplicity of α as an eigenvalue of the matrix N . This matrix is obtained from $A_\alpha(G)$ taking the entries corresponding to the internal vertices which are non quasi-pendant vertices. These results are applied to search for the multiplicity of α as an eigenvalue of $A_\alpha(G)$ when G is a path, a caterpillar, a circular caterpillar, a generalized Bethe tree or a Bethe tree. For the Bethe tree case, a simple formula for the nullity is given.

Keywords: Adjacency matrix, signless Laplacian matrix, Laplacian matrix, convex combination of matrices, eigenvalues, nullity.

2000 MSC: 05E30, 15A18, 05C50

1. Introduction

Let $G = (V(G), E(G))$ be a simple undirected graph on n vertices with vertex set $V(G)$ and edge set $E(G)$. Let $D(G)$ be the diagonal matrix of order n whose (i, i) -entry is the degree of the i -th vertex of G and let $A(G)$ be the adjacency matrix of G . The matrices $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ are the Laplacian and signless Laplacian matrix of G , respectively. The matrices $L(G)$ and $Q(G)$ are both positive semidefinite and $(0, \mathbf{1})$ is an eigenpair of $L(G)$ where $\mathbf{1}$ is the all ones vector. For a connected graph G , the smallest eigenvalue of $Q(G)$ is positive if and only if G is non-bipartite.

In [9], the family of matrices $A_\alpha(G)$,

$$A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$$

with $\alpha \in [0, 1]$, is introduced together with a number of some basic results and several open problems.

Observe that $A_0(G) = A(G)$ and $A_{1/2}(G) = \frac{1}{2}Q(G)$.

A pendant vertex is a vertex of degree 1 and a quasi-pendant vertex is a vertex adjacent to a pendant vertex. Let $p(G)$ be the number of pendant vertices and $q(G)$ be the number of quasi-pendant vertices. An internal vertex is a vertex of degree at least 2. Throughout this paper, we assume that G is a graph with pendant vertices.

The multiplicity of μ as an eigenvalue of the matrix M is denoted by $m_M(\mu)$.

The following results are due to I. Faria [6].

Lemma 1. [6] For any graph G ,

$$m_{L(G)}(1) \geq p(G) - q(G) \tag{1}$$

and

$$m_{Q(G)}(1) \geq p(G) - q(G). \tag{2}$$

In [1], it is proved that the equalities in (1) and (2) occur if each internal vertex is a quasi-pendant vertex. Moreover, if there is at least one internal vertex which is not a quasi-pendant vertex then the equalities

$$m_{L(G)}(1) = p(G) - q(G) + m_{N-}(1)$$

and

$$m_{Q(G)}(1) = p(G) - q(G) + m_{N^+}(1)$$

are determined, where the matrices N^- and N^+ are obtained from the Laplacian matrix and signless Laplacian matrix, respectively, taking the entries corresponding to the internal vertices which are non-quasi-pendant vertices.

We already observed that $A_{1/2}(G) = \frac{1}{2}Q(G)$. Then the above mentioned results can be used to find the multiplicity of $1/2$ as an eigenvalue of $A_{1/2}(G)$.

In this paper, we search for the multiplicity of $\alpha \in [0, 1]$ as an eigenvalue of $A_\alpha(G)$. The multiplicity of $\alpha = 0$ as eigenvalue of $A(G)$ is known as the nullity of G and it has been extensively studied (see [12, 13, 15]) and plays an important role in Chemistry. Some of its applications are described in [2, 5, 7, 8, 14, 16].

Since $A_1(G) = D(G)$, from now on, we consider $\alpha \in [0, 1)$.

For simplicity, we write $m_G(\alpha)$ instead $m_{A_\alpha(G)}(\alpha)$. More precisely, we prove that

$$m_G(\alpha) \geq p(G) - q(G). \quad (3)$$

If each internal vertex of G is a quasi-pendant vertex, it is proved that the equality in (3) holds. If there is at least one internal vertex which is not a quasi-pendant vertex, the equality

$$m_G(\alpha) = p(G) - q(G) + m_N(\alpha)$$

is determined in which $m_N(\alpha)$ is the multiplicity of α as an eigenvalue of the matrix N . This matrix is obtained from $A_\alpha(G)$ taking the entries corresponding to the internal vertices which are non quasi-pendant vertices.

Let $r(G)$ be the number of internal vertices of G . From now on, G is a connected graph on n vertices. Let

$$V_P = \{v \in V(G) : v \text{ is a pendant vertex}\},$$

$$V_Q = \{v \in V(G) : v \text{ is a quasi-pendant vertex}\}$$

and

$$C(G) = V(G) \setminus (V_P \cup V_Q).$$

Then $C(G)$ is the set of the internal vertices of G which are not quasi-pendant vertices.

Throughout the text, $|S|$ denotes the cardinality of the set S . Clearly $|V_P| = p(G)$, $|V_Q| = q(G)$ and $|C(G)| = n - p(G) - q(G)$.

We introduce some additional notation.

- The identity matrix is denoted by I and the zero matrix by 0 .
- If M is a matrix of order $m \times m$ with $m \geq 2$ then \widetilde{M} is the matrix obtained from M by deleting its last row and its last column.
- The determinant of a square matrix M is denoted by $|M|$ and the transpose of M by M^T .
- E denotes a matrix whose entries are zeros except the entry in the last row and last column which is 1.
- $G[F]$ denotes the subgraph of G induced by $F \subseteq V(G)$.

The orders of the matrices I , 0 and E will be clear from the context.

We recall Lemma 2.2 of [11] that will play an important role in this paper.

Lemma 2. [11] For $i = 1, 2, \dots, m$, let B_i be a matrix of order $k_i \times k_i$ and $\mu_{i,j}$ be arbitrary scalars. Then

$$\begin{aligned}
& \begin{vmatrix} B_1 & \mu_{1,2}E & \cdots & \mu_{1,m-1}E & \mu_{1,m}E \\ \mu_{2,1}E^T & B_2 & \cdots & \cdots & \mu_{2,m}E \\ \mu_{3,1}E^T & \mu_{3,2}E^T & \cdots & \cdots & \vdots \\ \vdots & \vdots & \vdots & B_{m-1} & \mu_{m-1,m}E \\ \mu_{m,1}E^T & \mu_{m,2}E^T & \cdots & \mu_{m,m-1}E^T & B_m \end{vmatrix} \\
= & \begin{vmatrix} |B_1| & \mu_{1,2}|\widetilde{B}_2| & \cdots & \mu_{1,m-1}|\widetilde{B}_{m-1}| & \mu_{1,m}|\widetilde{B}_m| \\ \mu_{2,1}|\widetilde{B}_1| & |B_2| & \cdots & \cdots & \mu_{2,m}|\widetilde{B}_m| \\ \mu_{3,1}|\widetilde{B}_1| & \mu_{3,2}|\widetilde{B}_2| & \cdots & \cdots & \vdots \\ \vdots & \vdots & \vdots & |B_{m-1}| & \mu_{m-1,m}|\widetilde{B}_m| \\ \mu_{m,1}|\widetilde{B}_1| & \mu_{m,2}|\widetilde{B}_2| & \cdots & \mu_{m,m-1}|\widetilde{B}_{m-1}| & |B_m| \end{vmatrix}.
\end{aligned}$$

The following result is Corollary 1.3 in [1].

Corollary 1. If $k_i = 1$, for some $1 \leq i \leq m$, then defining $\widetilde{B}_i = 1$ the equality in Lemma 2 also holds.

Another known result that we need is the following.

Lemma 3. Consider the square matrix of order $s + 1$

$$S(\alpha) = \begin{bmatrix} \alpha & 0 & \dots & 0 & 1 - \alpha \\ 0 & \alpha & \dots & 0 & 1 - \alpha \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & \alpha & 1 - \alpha \\ 1 - \alpha & 1 - \alpha & \dots & 1 - \alpha & \alpha d \end{bmatrix}.$$

Then the characteristic polynomial of $S(\alpha)$ is

$$|xI - S(\alpha)| = (x - \alpha)^{s-1}((x - \alpha d)(x - \alpha) - s(1 - \alpha)^2).$$

Throughout this paper, unless otherwise stated, $v_1, v_2, \dots, v_{r(G)}$ are the internal vertices of G . As usual, $u \sim v$ means that the vertices u and v are adjacent. For the internal vertices, d_i is the degree of the vertex v_i as a vertex of G and $\varepsilon_{i,j} = 1$ if $v_i \sim v_j$ and $\varepsilon_{i,j} = 0$, otherwise. For instance, $d_7 = 6$, $\varepsilon_{3,7} = 1$ and $\varepsilon_{7,10} = 0$ (see Figure 1 and Figure 2).

We label the vertices of G with the numbers $1, 2, \dots, n$, starting with the vertices of the stars $K_{1,s_1}, \dots, K_{1,s_{q(G)}}$, at each star the vertices are labeled beginning with the pendant vertices, and finishing with the internal vertices which are not quasi-pendants. This labeling of the vertices of G is herein called the global labeling of the vertices of G . Notice that, since the internal vertices of G are also denoted by $v_1, \dots, v_{q(G)}, \dots, v_{r(G)}$, for $j = 1, \dots, q(G)$, each of these vertices v_j corresponds, in the global labeling of the vertices of G , to the vertex $\sum_{i=1}^j s_i + j$ and the vertices $v_{q(G)+1}, \dots, v_{r(G)}$ correspond to the vertices $q(G) + p(G) + 1, \dots, n$, respectively.

2. Each internal vertex is a quasi-pendant vertex

In this section we consider the case in which each internal vertex of G is a quasi-pendant vertex. Then $r(G) = q(G)$ and there are stars $K_{1,s_1}, \dots, K_{1,s_{q(G)}}$ such that G is obtained by identifying the root of K_{1,s_i} with the i -th vertex of the graph induced by the quasi-pendant vertices. Moreover, $s_1 + s_2 + \dots + s_{q(G)} + q(G) = n$ and $s_i \geq 1$ for all i . We denote this graph G by $G(s_1, s_2, \dots, s_{q(G)})$. In Figure 1, we have an example of such a graph.




Figure 1: A example where each internal vertex is quasi-pendant.




Figure 2: The subgraph induced by the internal vertices $v_1 = 3, v_2 = 7, v_3 = 11, v_4 = 14, v_5 = 19$ and $v_6 = 23$ of the graph in Figure 1.

For $i = 1, 2, \dots, q(G)$, let

$$S_i(\alpha) = \begin{bmatrix} \alpha & 0 & \dots & 0 & 1 - \alpha \\ 0 & \alpha & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \ddots & \ddots & \alpha & 1 - \alpha \\ 1 - \alpha & \dots & \dots & 1 - \alpha & \alpha d_{v_i} \end{bmatrix}$$

of order $(s_i + 1) \times (s_i + 1)$ and

$$C_i(\alpha) = \begin{bmatrix} \alpha & (1 - \alpha)\sqrt{s_i} \\ (1 - \alpha)\sqrt{s_i} & \alpha d_{v_i} \end{bmatrix}.$$

From the definition of the matrices S_i and C_i , and Lemma 3, we have the following corollary.

Corollary 2. For $i = 1, 2, \dots, q(G)$,

1.

$$|xI - S_i(\alpha)| = (x - \alpha)^{s_i-1} |xI - C_i(\alpha)|.$$

and

2.

$$|\widetilde{xI - S_i(\alpha)}| = (x - \alpha)^{s_i}.$$

From now on, let $1 - \alpha = \beta$. Using the above mentioned labeling the matrix $A_\alpha(G)$, $G = G(s_1, s_2, \dots, s_{q(G)})$, becomes

$$A_\alpha(G) = \begin{bmatrix} S_1(\alpha) & \varepsilon_{1,2}\beta E & \dots & \varepsilon_{1,q(G)}\beta E \\ \varepsilon_{1,2}\beta E^T & S_2(\alpha) & \ddots & \vdots \\ \vdots & \ddots & \ddots & \varepsilon_{q(G)-1,q(G)}\beta E \\ \varepsilon_{1,q(G)}\beta E^T & \dots & \varepsilon_{q(G)-1,q(G)}\beta E^T & S_{q(G)}(\alpha) \end{bmatrix}$$

The next theorem gives the spectrum of $A_\alpha(G)$ if $G = G(s_1, s_2, \dots, s_{q(G)})$.

Theorem 1. If $G = G(s_1, s_2, \dots, s_{q(G)})$, the eigenvalues of $A_\alpha(G)$ are α with multiplicity at least $p(G) - q(G)$ and the eigenvalues of the $2q(G) \times 2q(G)$ matrix

$$X = \begin{bmatrix} C_1(\alpha) & \varepsilon_{1,2}\beta E & \dots & \dots & \varepsilon_{1,q(G)}\beta E \\ \varepsilon_{1,2}\beta E & C_2(\alpha) & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & C_{q(G)-1}(\alpha) & \varepsilon_{q(G)-1,q(G)}\beta E \\ \varepsilon_{1,q(G)}\beta E & \dots & \dots & \varepsilon_{q(G)-1,q(G)}\beta E & C_{q(G)}(\alpha) \end{bmatrix}.$$

Proof. Applying Lemma 2 and Corollary 2, together with a factoring in each column, we have $|xI - A_\alpha(G)| =$

$$\begin{vmatrix} (xI - S_1(\alpha)) & -\varepsilon_{1,2}\beta E & \dots & -\varepsilon_{1,q(G)}\beta E \\ -\varepsilon_{1,2}\beta E^T & (xI - S_2(\alpha)) & \ddots & \vdots \\ \vdots & \ddots & \ddots & -\varepsilon_{q(G)-1,q(G)}\beta E \\ -\varepsilon_{1,q(G)}\beta E^T & \dots & -\varepsilon_{q(G)-1,q(G)}\beta E^T & (xI - S_{q(G)}(\alpha)) \end{vmatrix} =$$

$$\prod_{j=1}^{q(G)} (x - \alpha)^{s_j - 1} \begin{vmatrix} |xI - C_1(\alpha)| & -\varepsilon_{1,2}\beta(x - \alpha) & \dots & -\varepsilon_{1,q(G)}\beta(x - \alpha) \\ -\varepsilon_{1,2}\beta(x - \alpha) & |xI - C_2(\alpha)| & \dots & \vdots \\ \vdots & \vdots & \ddots & -\varepsilon_{q(G)-1,q(G)}\beta(x - \alpha) \\ -\varepsilon_{1,q(G)}\beta(x - \alpha) & -\varepsilon_{2,q(G)}\beta(x - \alpha) & \dots & |xI - C_{q(G)}(\alpha)| \end{vmatrix}.$$

Applying again Lemma 2, it follows that $|xI - X| =$

$$\begin{vmatrix} |xI - C_1(\alpha)| & -\varepsilon_{1,2}\beta(x - \alpha) & \dots & -\varepsilon_{1,q(G)-1}\beta(x - \alpha) & -\varepsilon_{1,q(G)}\beta(x - \alpha) \\ -\varepsilon_{1,2}\beta(x - \alpha) & |xI - C_2(\alpha)| & \dots & -\varepsilon_{2,q(G)-1}\beta(x - \alpha) & -\varepsilon_{2,q(G)}\beta(x - \alpha) \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -\varepsilon_{1,q(G)-1}\beta(x - \alpha) & -\varepsilon_{2,q(G)-1}\beta(x - \alpha) & \dots & |xI - C_{q(G)-1}(\alpha)| & -\varepsilon_{q(G)-1,q(G)}\beta(x - \alpha) \\ -\varepsilon_{1,q(G)}\beta(x - \alpha) & -\varepsilon_{2,q(G)}\beta(x - \alpha) & \dots & -\varepsilon_{q(G)-1,q(G)}\beta(x - \alpha) & |xI - C_{q(G)}(\alpha)| \end{vmatrix}.$$

Finally, observe that $\prod_{j=1}^{q(G)} (x - \alpha)^{s_j - 1} = (x - \alpha)^{p(G) - q(G)}$. ■

Corollary 3. *The multiplicity of α as an eigenvalue of $A_\alpha(G)$, where $G = G(s_1, s_2, \dots, s_{q(G)})$, is exactly $p(G) - q(G)$.*

Proof. It is sufficient to prove that $|\alpha I - X| \neq 0$. From the above expression for $|xI - X|$, we have $|\alpha I - X| =$

$$\begin{vmatrix} |\alpha I - C_1(\alpha)| & 0 & \dots & 0 \\ 0 & |\alpha I - C_2(\alpha)| & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & |\alpha I - C_{q(G)}(\alpha)| \end{vmatrix}$$

and for $i = 1, 2, \dots, q(G)$, $|\alpha I - C_i(\alpha)| = -(1 - \alpha)^2 s_i \neq 0$. Hence $|\alpha I - X| \neq 0$. ■

3. Graphs having internal vertices which are non quasi-pendant

Let \mathbf{e} be a column vector of zeros except its last entry which is 1. As before, the dimension of \mathbf{e} will be clear from the context.

Suppose $r(G) > q(G)$. Then there are $r(G) - q(G)$ internal vertices which are non quasi-pendant vertices and $q(G)$ internal vertices which are the roots of the stars $K_{1,s_1}, K_{1,s_2}, \dots, K_{1,s_{q(G)}}$. Let us denote such a graph G by $G(s_1, \dots, s_{q(G)}, \mathbf{0})$, where $\mathbf{0}$ indicates a vector of zeros with $r(G) - q(G)$ entries. Without loss of generality, we assume that $V_Q = \{v_1, v_2, \dots, v_{q(G)}\}$ and $C(G) = \{v_{q(G)+1}, v_{q(G)+2}, \dots, v_r(G)\}$. We recall that the global labeling

for the vertices of $G(s_1, \dots, s_{q(G)}, \mathbf{0})$ is such that the labels $1, 2, \dots, p(G) + q(G)$ are used for the vertices of the stars $K_{1,s_1}, K_{1,s_2}, \dots, K_{1,s_{q(G)}}$, and the labels $p(G) + q(G) + 1, \dots, n$ are used for the internal vertices which are non quasi-pendant, as illustrated in Figure 3. Using this global labeling jointly with the labels $v_1, \dots, v_{r(G)}$ for the internal vertices (as before) the matrix $A_\alpha(G)$, where $G = G(s_1, \dots, s_{q(G)}, \mathbf{0})$, is

$$A_\alpha(G) = \begin{bmatrix} U & V \\ V^T & N \end{bmatrix}$$

where

$$U = \begin{bmatrix} S_1(\alpha) & \varepsilon_{1,2}\beta E & \dots & \varepsilon_{1,q(G)-1}\beta E & \varepsilon_{1,q(G)}\beta E \\ \varepsilon_{1,2}\beta E^T & S_2(\alpha) & \dots & \varepsilon_{2,q(G)-1}\beta E & \varepsilon_{2,q(G)}\beta E \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \varepsilon_{1,q(G)-1}\beta E^T & \varepsilon_{2,q(G)-1}\beta E & \dots & S_{q(G)-1}(\alpha) & \varepsilon_{q(G)-1,q(G)}\beta E \\ \varepsilon_{1,q(G)}\beta E^T & \varepsilon_{2,q(G)}\beta E & \dots & \varepsilon_{q(G)-1,q(G)}\beta E^T & S_{q(G)}(\alpha) \end{bmatrix},$$

$$V = \beta \begin{bmatrix} \varepsilon_{1,q(G)+1}\mathbf{e} & \varepsilon_{1,q(G)+2}\mathbf{e} & \dots & \varepsilon_{1,r(G)-1}\mathbf{e} & \varepsilon_{1,r(G)}\mathbf{e} \\ \varepsilon_{2,q(G)+1}\mathbf{e} & \varepsilon_{2,q(G)+2}\mathbf{e} & \dots & \varepsilon_{1,r(G)-1}\mathbf{e} & \varepsilon_{2,r(G)}\mathbf{e} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \varepsilon_{q(G)-1,q(G)+1}\mathbf{e} & \varepsilon_{q(G)-1,q(G)+2}\mathbf{e} & \dots & \varepsilon_{q(G)-1,r(G)-1}\mathbf{e} & \varepsilon_{q(G)-1,r(G)}\mathbf{e} \\ \varepsilon_{q(G),q(G)+1}\mathbf{e} & \varepsilon_{q(G),q(G)+2}\mathbf{e} & \dots & \varepsilon_{q(G),r(G)-1}\mathbf{e} & \varepsilon_{q(G),r(G)}\mathbf{e} \end{bmatrix}$$

and

$$N = \begin{bmatrix} \alpha d_{q(G)+1} & \varepsilon_{q(G)+1,q(G)+2}\beta & \dots & \varepsilon_{q(G)+1,r(G)-1}\beta & \varepsilon_{q(G)+1,r(G)}\beta \\ \varepsilon_{q(G)+1,q(G)+2}\beta & \alpha d_{q(G)+2} & \dots & \varepsilon_{q(G)+2,r(G)-1}\beta & \varepsilon_{q(G)+2,r(G)}\beta \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \varepsilon_{q(G)+1,r(G)-1}\beta & \varepsilon_{q(G)+2,r(G)-1}\beta & \dots & \alpha d_{r(G)-1} & \varepsilon_{r(G)-1,r(G)}\beta \\ \varepsilon_{q(G)+1,r(G)}\beta & \varepsilon_{q(G)+2,r(G)}\beta & \dots & \varepsilon_{r(G)-1,r(G)}\beta & \alpha d_{r(G)} \end{bmatrix}$$

where $d_{q(G)+1}, d_{q(G)+2}, \dots, d_{r(G)-1}, d_{r(G)}$ are the degrees of the vertices $v_{q(G)+1}, v_{q(G)+2}, \dots, v_{r(G)-1}, v_{r(G)}$, respectively.

Applying Lemma 2, Corollary 1 and Corollary 2, together with a factoring in each of the first $q(G)$ columns of the resulting determinant, one can prove the following theorem.

Theorem 2. If $G = G(s_1, s_2, \dots, s_{q(G)}, \mathbf{0})$, the eigenvalues of $A_\alpha(G)$ are α with multiplicity at least $p(G) - q(G)$ and the eigenvalues of the $(n + q(G) - p(G)) \times (n + q(G) - p(G))$ matrix

$$X = \begin{bmatrix} Q & R \\ R^T & N \end{bmatrix}$$

where

$$Q = \begin{bmatrix} C_1(\alpha) & \varepsilon_{1,2}\beta E & \cdots & \varepsilon_{1,q(G)-1}\beta E & \varepsilon_{1,q(G)}\beta E \\ \varepsilon_{1,2}\beta E & C_2(\alpha) & \cdots & \varepsilon_{2,q(G)-1}\beta E & \varepsilon_{2,q(G)}\beta E \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \varepsilon_{1,q(G)-1}\beta E & \varepsilon_{2,q(G)-1}\beta E & \cdots & C_{q(G)-1}(\alpha) & \varepsilon_{q(G)-1,q(G)}\beta E \\ \varepsilon_{1,q(G)}\beta E & \varepsilon_{2,q(G)}\beta E & \cdots & \varepsilon_{q(G)-1,q(G)}\beta E & C_{q(G)}(\alpha) \end{bmatrix},$$

$$N = \begin{bmatrix} \alpha d_{q(G)+1} & \varepsilon_{q(G)+1,q(G)+2}\beta & \cdots & \varepsilon_{q(G)+1,r(G)-1}\beta & \varepsilon_{q(G)+1,r(G)}\beta \\ \varepsilon_{q(G)+1,q(G)+2}\beta & \alpha d_{q(G)+2} & \cdots & \varepsilon_{q(G)+2,r(G)-1}\beta & \varepsilon_{q(G)+2,r(G)}\beta \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \varepsilon_{q(G)+1,r(G)-1}\beta & \varepsilon_{q(G)+2,r(G)-1}\beta & \cdots & \alpha d_{r(G)-1} & \varepsilon_{r(G)-1,r(G)}\beta \\ \varepsilon_{q(G)+1,r(G)}\beta & \varepsilon_{q(G)+2,r(G)}\beta & \cdots & \varepsilon_{r(G)-1,r(G)}\beta & \alpha d_{r(G)} \end{bmatrix}$$

and

$$R = \beta \begin{bmatrix} \varepsilon_{1,q(G)+1}\mathbf{e} & \varepsilon_{1,q(G)+2}\mathbf{e} & \cdots & \varepsilon_{1,r(G)-1}\mathbf{e} & \varepsilon_{1,r(G)}\mathbf{e} \\ \varepsilon_{2,q(G)+1}\mathbf{e} & \varepsilon_{2,q(G)+2}\mathbf{e} & \cdots & \varepsilon_{2,r(G)-1}\mathbf{e} & \varepsilon_{2,r(G)}\mathbf{e} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \varepsilon_{q(G)-1,q(G)+1}\mathbf{e} & \varepsilon_{q(G)-1,q(G)+2}\mathbf{e} & \cdots & \varepsilon_{q(G)-1,r(G)-1}\mathbf{e} & \varepsilon_{q(G)-1,r(G)}\mathbf{e} \\ \varepsilon_{q(G),q(G)+1}\mathbf{e} & \varepsilon_{q(G),q(G)+2}\mathbf{e} & \cdots & \varepsilon_{q(G),r(G)-1}\mathbf{e} & \varepsilon_{q(G),r(G)}\mathbf{e} \end{bmatrix}.$$

Theorem 3. Let $G = G(s_1, s_2, \dots, s_{q(G)}, \mathbf{0})$. Let X, Q and N be the matrices in Theorem 2. Then

1. $m_X(\alpha) = m_N(\alpha)$.
2. $m_G(\alpha) = p(G) - q(G) + m_N(\alpha)$.

Proof.

1. We have $|\alpha I - X| = \begin{vmatrix} \alpha I - Q & -R \\ -R^T & \alpha I - N \end{vmatrix}$. Hence, applying Corollary 1,




Figure 3: The graph $G(2, 3, 3, 4, \mathbf{0})$.

we obtain $|\alpha I - X| =$

$$\begin{vmatrix} -(1-\alpha)^2 s_1 & 0 & \cdots & 0 & * \\ 0 & -(1-\alpha)^2 s_2 & \vdots & \vdots & \vdots \\ \vdots & 0 & \ddots & 0 & \vdots \\ \vdots & \cdots & \cdots & -(1-\alpha)^2 s_{q(G)} & * \\ 0 & 0 & \cdots & 0 & |\alpha I - N| \end{vmatrix} = (-1)^{q(G)} (1-\alpha)^{2q(G)} s_1 s_2 \cdots s_{q(G)} |\alpha I - N|.$$

Since $1 - \alpha \neq 0$ and $s_i \geq 1$, for $1 \leq i \leq q(G)$, we obtain $|\alpha I - X| = 0$ if and only if $|\alpha I - N| = 0$. Hence $m_X(\alpha) = m_N(\alpha)$.

2. From Theorem 2, $m_G(\alpha) = p(G) - q(G) + m_X(\alpha)$. Since $m_X(\alpha) = m_N(\alpha)$, the result follows. ■

We see that the matrix N in Theorem 2 is obtained from $A_\alpha(G)$ by taking the entries corresponding to the internal vertices which are not quasi-pendant vertices.

We recall that $G[F]$ denotes the subgraph of G induced by $F \subseteq V(G)$.

Theorem 4. *Let $G = G(s_1, s_2, \dots, s_{q(G)}, \mathbf{0})$. Let X, Q and N be the matrices in Theorem 3. Let C_1, \dots, C_t be the components of the subgraph induced by $C(G)$. Then*

$$m_G(\alpha) = p(G) - q(G) + \sum_{i=1}^t m_{N_i}(\alpha),$$

where, for $1 \leq i \leq t$, $N_i = (1 - \alpha)A(G[C_i]) + \alpha D_i$, where D_i is a diagonal matrix of order $|C_i|$ in which each diagonal entry is the degree in G of the corresponding vertex.

Proof. From the hypothesis, there is a labeling of the vertices of $C(G)$ such that $N = \bigoplus_{i=1}^t N_i$ (the direct sum of the matrices N_i , $i = 1, \dots, t$). Therefore $m_N(\alpha) = \sum_{i=1}^t m_{N_i}(\alpha)$. Now, the result is immediate by Theorem 3. ■

Example 1. For the graph G in Figure 3, we have $p(G) = 12$, $q(G) = 4$. We see that $N = \begin{bmatrix} 3\alpha & 0 \\ 0 & 3\alpha \end{bmatrix}$. Hence $|\alpha I - N| = \begin{vmatrix} \alpha - 3\alpha & 0 \\ 0 & \alpha - 3\alpha \end{vmatrix} = 0$ if and only if $\alpha = 0$. Therefore, α is not an eigenvalue of N , when $\alpha \neq 0$. Then, from Theorem 3, the multiplicity of α as an eigenvalue of $A_\alpha(G)$ is

$$\begin{cases} p(G) - q(G) = 8, & \text{when } \alpha \neq 0; \\ p(G) - q(G) + m_N(0) = 10, & \text{otherwise.} \end{cases}$$

We recall that the nullity of a graph G , denoted by $\eta(G)$, is the multiplicity of 0 as an eigenvalue of $A(G)$. From Theorem 4, we obtain

Corollary 4. Let $G = G(s_1, s_2, \dots, s_{q(G)}, \mathbf{0})$. Let C_1, \dots, C_t be the components of the subgraph induced by $C(G)$. Then

$$\eta(G) = p(G) - q(G) + \sum_{i=1}^t \eta(N_i), \quad (4)$$

where, for $1 \leq i \leq t$, $N_i = A(G[C_i])$ and $\eta(N_i)$ is the multiplicity of 0 as an eigenvalue of N_i .

Corollary 5. Let $G = G(s_1, s_2, \dots, s_{q(G)}, \mathbf{0})$. Let C_1, \dots, C_t be the components of the subgraph induced by $C(G)$. Let H be the induced subgraph of G obtained by deleting one pendant vertex together with the vertex adjacent to it. Then $\eta(G) = \eta(H)$.

Proof. Clearly $C(H) = C(G)$ and the components of the subgraphs induced by $C(H)$ and $C(G)$ are the same. In addition, $p(H) - q(H) = p(G) - 1 - (q(G) - 1) = p(G) - q(G)$. Applying (4), we obtain $\eta(H) = \eta(G)$. ■

A version of Corollary 5 is proved in [3] assuming that G is a bipartite graph with at least one pendant vertex.

4. Applications on some particular graphs

In this section, the above results are applied to search for the multiplicity of $\alpha \in [0, 1)$ as an eigenvalue of $A_\alpha(G)$ when G is a path, a caterpillar, a circular caterpillar, a generalized Bethe tree or a Bethe tree.

4.1. The multiplicity of α as an eigenvalue of $A_\alpha(P_n)$

Let P_n be the path of n vertices. It is well known that 0 is an eigenvalue of $A(P_n)$ if and only if n is odd. Moreover, if n is odd then 0 is a simple eigenvalue of $A(P_n)$. We begin considering the cases $2 \leq n \leq 4$.

- If $n = 2$ then $A_\alpha(P_2) = \begin{bmatrix} \alpha & 1 - \alpha \\ 1 - \alpha & \alpha \end{bmatrix}$. We have $|A_\alpha(P_2) - \alpha I_2| = \begin{vmatrix} 0 & 1 - \alpha \\ 1 - \alpha & 0 \end{vmatrix} = 0$ if and only if $\alpha = 1$. Hence $\alpha \in (0, 1)$ is not an eigenvalue of $A_\alpha(P_2)$.
- If $n = 3$ then P_3 is a graph with 2 pendant vertices and 1 quasi-pendant vertex. From Corollary 3, $m_{P_3}(\alpha) = 2 - 1 = 1$.
- If $n = 4$ then P_4 is a graph with 2 pendant vertices and 2 quasi-pendant vertices. From Corollary 3, $m_{P_4}(\alpha) = 2 - 2 = 0$. Hence $\alpha \in (0, 1)$ is not an eigenvalue of $A_\alpha(P_4)$.

From now on let us assume that P_n is such that $n \geq 5$. Since $\alpha \neq 1$, the vertices of P_n can be labeled such that $A_\alpha(P_n)$ is a symmetric tridiagonal matrix with nonzero codiagonal entries. Hence the eigenvalues of $A_\alpha(P_n)$ are simple and, in particular, if $\alpha \neq 1$ is an eigenvalue of $A_\alpha(P_n)$ then it will be a simple eigenvalue. We recall the following lemma (see [4]).

Lemma 4. *The eigenvalues of the symmetric tridiagonal matrix*

$$N(\alpha) = \begin{bmatrix} 2\alpha & 1 - \alpha & & & & & \\ 1 - \alpha & 2\alpha & 1 - \alpha & & & & \\ & & \ddots & \ddots & \ddots & & \\ & & & 1 - \alpha & 2\alpha & 1 - \alpha & \\ & & & & 1 - \alpha & 2\alpha & \\ & & & & & & \end{bmatrix} \quad (5)$$

of order $s \times s$ are

$$2\alpha + 2(1 - \alpha) \cos\left(\frac{\pi j}{s + 1}\right)$$

for $j = 1, 2, \dots, s$.

Corollary 6. Let $N(\alpha)$ as in (5).

1. If $2/3 \leq \alpha < 1$ then α is not an eigenvalue of $N(\alpha)$.
2. γ is an eigenvalue of $N(\gamma)$ if and only if

$$\gamma = \frac{-2 \cos(\frac{\pi j}{s+1})}{1 - 2 \cos(\frac{\pi j}{s+1})}$$

for some $j = 1, \dots, s$.

3. Let

$$\alpha_j = \frac{-2 \cos(\frac{\pi j}{s+1})}{1 - 2 \cos(\frac{\pi j}{s+1})}. \quad (6)$$

Then α_j is an eigenvalue of $N(\alpha_j)$ and $0 < \alpha_j < 1$ if and only if $j \in \{\lfloor \frac{s+3}{2} \rfloor, \dots, s\}$.

Proof.

1. Assume $2/3 \leq \alpha < 1$. Using this hypothesis, the matrix $N(\alpha) - \alpha I$ is irreducible and diagonally dominant with strict inequality in at least one row. Then, by Theorem 1.21 in [17], $N(\alpha) - \alpha I$ is an invertible matrix. Hence α is not an eigenvalue of $N(\alpha)$.
2. It is immediate from Lemma 4.
3. Let α_j as in (6). Then α_j is an eigenvalue of $N(\alpha_j)$. The following fact is immediate:

$$0 < \frac{x}{1+x} < 1 \text{ if and only if } x > 0.$$

Let $j \in \{\lfloor \frac{s+3}{2} \rfloor, \dots, s\}$. Then $-2 \cos(\frac{\pi j}{s+1}) > 0$. From the above fact, $0 < \alpha_j = \frac{-2 \cos(\frac{\pi j}{s+1})}{1 - 2 \cos(\frac{\pi j}{s+1})} < 1$. Conversely, suppose that $0 < \alpha_j < 1$. We use again the above mentioned fact, to obtain that $-2 \cos(\frac{\pi j}{s+1}) > 0$. Hence $j \in \{\lfloor \frac{s+3}{2} \rfloor, \dots, s\}$.

■

Since $n \geq 5$, P_n has 2 pendant vertices, 2 quasi-pendant vertices and $n-4$ internal vertices which are not quasi-pendants. By Theorem 3, $m_{P_n}(\alpha) =$

$2 - 2 + m_N(\alpha) = m_N(\alpha)$, where

$$N = N(\alpha) = \begin{bmatrix} 2\alpha & 1 - \alpha & & & & \\ 1 - \alpha & 2\alpha & 1 - \alpha & & & \\ & & \ddots & \ddots & \ddots & \\ & & & 1 - \alpha & 2\alpha & 1 - \alpha \\ & & & & 1 - \alpha & 2\alpha \end{bmatrix}$$

of order $(n - 4) \times (n - 4)$.

In particular, for $n = 5$ and $n = 6$, we have

- $m_{P_5}(\alpha) = m_N(\alpha)$ where $N = N(\alpha) = [2\alpha]$. Since $\alpha \neq 0$ is not an eigenvalue of $[2\alpha]$, we have that $\alpha \neq 0$ is not an eigenvalue of $A_\alpha(P_5)$.

- $m_{P_6}(\alpha) = m_N(\alpha)$ where $N = N(\alpha) = \begin{bmatrix} 2\alpha & 1 - \alpha \\ 1 - \alpha & 2\alpha \end{bmatrix}$. We have

$$|N(\alpha) - \alpha I_2| = \begin{vmatrix} \alpha & 1 - \alpha \\ 1 - \alpha & \alpha \end{vmatrix} = 0 \text{ if and only if } \alpha = 1/2. \text{ Hence } \alpha \in (0, 1) \text{ is an eigenvalue of } A_\alpha(P_6) \text{ if and only if } \alpha = 1/2.$$

Applying Theorem 4 and Corollary 6, we get

Corollary 7. *Let $n \geq 7$. Then $\alpha \in (0, 1)$ is an eigenvalue of $A_\alpha(P_n)$ if and only if*

$$\alpha = \frac{-2 \cos(\frac{\pi j}{n-3})}{1 - 2 \cos(\frac{\pi j}{n-3})}$$

for some $j \in \{\lfloor \frac{n-1}{2} \rfloor, \dots, n-4\}$.

4.2. Multiplicity of α as an eigenvalue of $A_\alpha(G)$ when G is a caterpillar or a circular caterpillar

We recall that a graph G is a caterpillar (respectively, a circular caterpillar) if its internal vertices induce a path (respectively, a cycle). We say that a caterpillar or a circular caterpillar is complete if each internal vertex is a quasi-pendant vertex. From Corollary 3, we get

Corollary 8. *If G is a complete caterpillar or a complete circular caterpillar then $m_G(\alpha) = p(G) - q(G)$.*




Figure 4: An example of a circular caterpillar.

If G is a caterpillar or a circular caterpillar having internal vertices which are not quasi-pendant vertices, we may use Theorem 4 together with equation 6 to find $m_G(\alpha)$.

Example 2. Let us find the exact multiplicity of $\alpha \in [0, 1)$ as an eigenvalue of $A_\alpha(G)$ where G is the circular caterpillar in Figure 4. Applying Theorem 4, we obtain

$$m_G(\alpha) = p(G) - q(G) + m_{N_1}(\alpha) + m_{N_2}(\alpha) = 14 - 6 + m_{N_1}(\alpha) + m_{N_2}(\alpha)$$

where $N_1 = N_1(\alpha) = \begin{bmatrix} 2\alpha & 1-\alpha & 0 \\ 1-\alpha & 2\alpha & 1-\alpha \\ 0 & 1-\alpha & 2\alpha \end{bmatrix}$ and $N_2 = N_2(\alpha) = [2\alpha]$. By equation (6) in Corollary 6, $\alpha \in (0, 1)$ is an eigenvalue of $N_1(\alpha)$ if and only if $\alpha = \alpha_3 = \frac{-2\cos(\frac{3\pi}{4})}{1-2\cos(\frac{3\pi}{4})} = \frac{\sqrt{2}}{1+\sqrt{2}}$.

Moreover, $\alpha \in [0, 1)$ is an eigenvalue of N_2 if and only if $\alpha = 0$. Then, from Theorem 4, the multiplicity of α as an eigenvalue of $A_\alpha(G)$ is

$$\begin{cases} 8 + m_{N_1}(\alpha) + m_{N_2}(\alpha) = 8 + 1 + 0 = 9, & \text{when } \alpha = \alpha_3; \\ 8 + m_{N_1}(0) + m_{N_2}(0) = 8 + 0 + 1 = 9, & \text{when } \alpha = 0; \\ 8 + m_{N_1}(\alpha) + m_{N_2}(\alpha) = 8, & \text{when } \alpha \in (0, 1) \setminus \{\alpha_3\}. \end{cases}$$

4.3. Multiplicity of α as an eigenvalue of $A_\alpha(T)$ when T is a generalized Bethe tree

Given a rooted graph, the level of a vertex is equal to its distance to the root vertex increased by one. A generalized Bethe tree is a rooted tree in which vertices at the same level have the same degree.




Figure 5: An example of a generalized Bethe tree of 4 levels.

In this section, B_k is a generalized Bethe tree of $k > 1$ levels. Given a B_k and an integer $1 \leq j \leq k$, n_{k-j+1} is the number of vertices at level j and d_{k-j+1} is the degree of them. In particular, $d_1 = 1$, $n_k = 1$, $n_1 = p(B_k)$ and $n_2 = q(B_k)$.

Definition 1. For $j = 1, 2, \dots, k-1$, let T_j be the $j \times j$ leading principal submatrix of the $k \times k$ symmetric tridiagonal matrix

$$T_k = \begin{bmatrix} \alpha & \beta\sqrt{d_2-1} & 0 & \cdots & 0 & 0 & 0 \\ \beta\sqrt{d_2-1} & \alpha d_2 & \beta\sqrt{d_3-1} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha d_{k-2} & \beta\sqrt{d_{k-1}-1} & 0 \\ 0 & 0 & 0 & \cdots & \beta\sqrt{d_{k-1}-1} & \alpha d_{k-1} & \beta\sqrt{d_k} \\ 0 & 0 & 0 & \cdots & 0 & \beta\sqrt{d_k} & \alpha d_k \end{bmatrix}$$

where $\beta = 1 - \alpha$.

Let $\sigma(M)$ be the multiset of eigenvalues of the matrix M .

Since $d_s > 1$ for all $s = 2, 3, \dots, j$, each matrix T_j has nonzero codiagonal entries. Then the eigenvalues of each T_j are simple and $\sigma(T_j) \cap \sigma(T_{j+1}) = \emptyset$ for $j = 1, \dots, k-1$.

The following theorem was proved in [10].

Theorem 5. Let B_k be a generalized Bethe tree.

1. The multiset of the eigenvalues of $A_\alpha(B_k)$ is

$$\sigma(A_\alpha(B_k)) = \sigma(T_1) \cup \cdots \cup \sigma(T_k). \quad (7)$$

2. The multiplicity of each eigenvalue of T_j as an eigenvalue of $A_\alpha(B_k)$ is $n_j - n_{j+1}$ if $1 \leq j \leq k-1$, and is 1 if $j = k$. If some eigenvalues obtained in different matrices are equal, their multiplicities are added together.
3. The largest eigenvalue of T_k is the largest eigenvalue of $A_\alpha(B_k)$.

We observe that for $k \geq 3$,

$$T_2 = \begin{bmatrix} \alpha & (1-\alpha)\sqrt{d_2-1} \\ (1-\alpha)\sqrt{d_2-1} & \alpha d_2 \end{bmatrix}.$$

Hence $|T_2 - \alpha I_2| = \begin{vmatrix} 0 & (1-\alpha)\sqrt{d_2-1} \\ (1-\alpha)\sqrt{d_2-1} & \alpha(d_2-1) \end{vmatrix} = -(1-\alpha)^2(d_2-1) \neq 0$ because $\alpha \neq 1$. The next corollary follows easily from Theorem 5.

Corollary 9. *Let $\alpha \in [0, 1)$. Then*

1. $m_{B_2}(\alpha) = n_1 - 1$.
2. $m_{B_3}(\alpha) = n_1 - n_2$ if $\alpha \neq 0$ and $m_{B_3}(0) = n_1 - n_2 + 1$.
3. For $k \geq 4$,

$$m_{B_k}(\alpha) = n_1 - n_2 + \sum_{j=3}^{k-1} (n_j - n_{j+1})m_{T_j}(\alpha) + m_{T_k}(\alpha) \quad (8)$$

in which, for $j = 3, \dots, k$, $m_{T_j}(\alpha) = 1$ if $\alpha \in \sigma(T_j)$ and $m_{T_j}(\alpha) = 0$ otherwise.

We already observed that $\sigma(T_j) \cap \sigma(T_{j+1}) = \emptyset$ for $j = 1, \dots, k-1$.

Example 3. *Let us find $m_{B_4}(\alpha)$. Using (8),*

$$m_{B_4}(\alpha) = n_1 - n_2 + (n_3 - n_4)m_{T_3}(\alpha) + m_{T_4}(\alpha)$$

where $n_4 = 1$, $T_3 = \begin{bmatrix} \alpha & (1-\alpha)\sqrt{d_2-1} & 0 \\ (1-\alpha)\sqrt{d_2-1} & \alpha d_2 & (1-\alpha)\sqrt{d_3-1} \\ 0 & (1-\alpha)\sqrt{d_3-1} & \alpha d_3 \end{bmatrix}$

and

$$T_4 = \begin{bmatrix} \alpha & (1-\alpha)\sqrt{d_2-1} & 0 & 0 \\ (1-\alpha)\sqrt{d_2-1} & \alpha d_2 & (1-\alpha)\sqrt{d_3-1} & 0 \\ 0 & (1-\alpha)\sqrt{d_3-1} & \alpha d_3 & (1-\alpha)\sqrt{d_4} \\ 0 & 0 & (1-\alpha)\sqrt{d_4} & \alpha d_4 \end{bmatrix}.$$

It is easily to find that

$$|T_3 - \alpha I_3| = -\alpha(1 - \alpha)^2(d_2 - 1)(d_3 - 1)$$

and

$$|T_4 - \alpha I_4| = -(1 - \alpha)^2(d_2 - 1)(\alpha^2(d_3 - 1)(d_4 - 1) - (1 - \alpha)^2 d_4).$$

Hence $\alpha \in [0, 1)$ is an eigenvalue of T_3 if and only if $\alpha = 0$ and $\alpha \in [0, 1)$ is an eigenvalue of T_4 if and only if $\alpha = \alpha_0 = \frac{\sqrt{d_4}}{\sqrt{d_4} + \sqrt{(d_3 - 1)(d_4 - 1)}}$. Therefore the multiplicity of α as an eigenvalue of $A_\alpha(B_4)$ is

$$\begin{cases} n_1 - n_2 + n_3 - 1, & \text{when } \alpha = 0; \\ n_1 - n_2 + 1, & \text{when } \alpha = \alpha_0; \\ n_1 - n_2, & \text{when } \alpha \in (0, 1) \setminus \{\alpha_0\}. \end{cases}$$

4.4. Multiplicity of α as an eigenvalue of $A_\alpha(T)$ and the nullity of T when T is a Bethe tree

A Bethe tree $B(d, k)$ is a rooted tree of k levels in which the root has degree d , the vertices in level j ($2 \leq j \leq k - 1$) have degree equal to $d + 1$ and the vertices in level k have degree equal to 1 (pendant vertices). Clearly, any Bethe tree $B(d, k)$ is a generalized Bethe tree in which the matrix T_j is the $j \times j$ leading principal submatrix of the $k \times k$ symmetric tridiagonal matrix

$$T_k = \begin{bmatrix} \alpha & \beta\sqrt{d} & 0 & \cdots & 0 & 0 & 0 \\ \beta\sqrt{d} & \alpha d & \beta\sqrt{d} & \cdots & 0 & 0 & 0 \\ 0 & \beta\sqrt{d} & \alpha d & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha d & \beta\sqrt{d} & 0 \\ 0 & 0 & 0 & \cdots & \beta\sqrt{d} & \alpha d & \beta\sqrt{d} \\ 0 & 0 & 0 & \cdots & 0 & \beta\sqrt{d} & \alpha d \end{bmatrix} \quad (9)$$

where $\beta = 1 - \alpha$. Moreover, $n_j = d^{k-j}$ for $j = 1, \dots, k$. From Corollary 9, we get

Corollary 10. *Let $\alpha \in [0, 1)$. Then*

1. $m_{B(d,2)}(\alpha) = d - 1$.
2. $m_{B(d,3)}(\alpha) = (d - 1)d$ if $\alpha \neq 0$ and $\eta(B(d, 3)) = m_{B(d,3)}(0) = (d - 1)d + 1$.

3. For $k \geq 4$,

$$m_{B(d,k)}(\alpha) = (d-1)(d^{k-2} + \sum_{j=3}^{k-1} d^{k-j-1} m_{T_j}(\alpha)) + m_{T_k}(\alpha) \quad (10)$$

in which the matrix T_j , for $j = 3, \dots, k$, is the $j \times j$ principal submatrix of T_k as in (9) and $m_{T_j}(\alpha) = 1$ if $\alpha \in \sigma(T_j)$ and $m_{T_j}(\alpha) = 0$ otherwise.

Finally, we find the nullity of $B(d, k)$. Let $\alpha = 0$. Then, $\beta = 1$ and the matrix T_k becomes

$$T_k = \begin{bmatrix} 0 & \sqrt{d} & 0 & \cdots & 0 & 0 & 0 \\ \sqrt{d} & 0 & \sqrt{d} & \cdots & 0 & 0 & 0 \\ 0 & \sqrt{d} & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & \sqrt{d} & 0 \\ 0 & 0 & 0 & \cdots & \sqrt{d} & 0 & \sqrt{d} \\ 0 & 0 & 0 & \cdots & 0 & \sqrt{d} & 0 \end{bmatrix} \quad (11)$$

Corollary 11. *The nullity of $B(d, k)$ is given by*

$$\eta(B(d, k)) = \begin{cases} \frac{d^k - 1}{d + 1}, & \text{if } k \text{ is even;} \\ \frac{d^k + 1}{d + 1}, & \text{otherwise.} \end{cases}$$

Proof. We recall that, for $j = 1, \dots, k-1$, T_j is the $j \times j$ principal submatrix of T_k as in (11). One can easily see that 0 is an eigenvalue of T_j if and only if j is odd.

1. Let k be an even integer.

For $k = 2$, we have $m_{B(d,2)}(0) = \eta(B(d, 2)) = d - 1 = \frac{d^2 - 1}{d + 1}$. Let $k \geq 4$. From (10), we obtain

$$\eta(B(d, k)) = (d-1)(d^{k-2} + d^{k-4} + \dots + d^2 + 1) = \frac{d^k - 1}{d + 1}.$$

2. Let k be an odd integer.

For $k = 3$, we have $m_{B(d,3)}(0) = \eta(B(d, 3)) = (d-1)d + 1 = \frac{d^3 + 1}{d + 1}$. Let $k \geq 5$. From (10), we obtain

$$\eta(B(d, k)) = (d-1)(d^{k-2} + d^{k-4} + \dots + d^3 + d) + 1 = \frac{d^k + 1}{d + 1}.$$

■

Acknowledgments

The authors express their thanks to the referee for the valuable comments.

The research of D. M. Cardoso author is partially supported by Portuguese funds through the CIDMA - Center for Research and Development in Mathematics and Applications, and the Portuguese Foundation for Science and Technology (“FCT-Fundação para a Ciência e a Tecnologia”), within project PEst-OE/MAT/UI4106/2013. The research of G. Pastén is supported by CONICYT-PCHA/MagisterNacional/2015-22151590. O. Rojo thanks the hospitality of the Department of Mathematics, Universidade de Aveiro, Aveiro, Portugal, where part of this research was conducted.

References

- [1] E. Andrade, D. M. Cardoso, G. Pastén, O. Rojo, On the Faria’s inequality for the Laplacian and signless Laplacian spectra: A unified approach, *Linear Algebra Appl.* **472** (2015), 81–96.
- [2] B. Borovićanin, I. Gutman, Nullity of graphs, in: D. Cvetković, I. Gutman, Eds. Applications of Graph Spectra, Math. Inst., Belgrade, 2009, 107–122.
- [3] D. Cvetković, I. Gutman, The algebraic multiplicity of the number zero in the spectrum of a bipartite graph, *Matematički Vesnik (Beograd)* 9 (1972), 141–150.
- [4] S. Kouachi, Eigenvalues and eigenvectors of tridiagonal matrix, *Electronic Journal of Linear Algebra*, Volume 15, pp. 115-133, April 2006.
- [5] J. R. Dias, *Molecular Orbital Calculations Using Chemical Graph Theory*, Springer–Verlag, Berlin, 1993.
- [6] I. Faria, Permanent roots and the star degree of a graph. *Linear Algebra Appl.* **64** (1985), 255-265.
- [7] A. Graovac, I. Gutman, N. Trinajstić, *Topological Approach to the Chemistry of Conjugated Molecules*, Springer–Verlag, Berlin, 1977.
- [8] I. Gutman, O.E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer–Verlag, Berlin, 1986.

- [9] V. Nikiforov, Merging the A - and Q -spectral theories, *Applicable Analysis and Discrete Math.* **11** (2017), 81–107.
- [10] V. Nikiforov, G. Pastén, O. Rojo, R. L. Soto, On the A_α -spectra of trees, *Linear Algebra Appl.* **520** (2017), 286–305.
- [11] O. Rojo, L. Medina, Spectra of weighted compound graphs of generalized Bethe trees, *Electronic Journal of Linear Algebra* **18** (2009), 30–57.
- [12] I. Sciriha, On the coefficient of λ in the characteristic polynomial of singular graphs, *Utilitas Mathematica* **52** (1997), 97–111.
- [13] I. Sciriha, On the construction of graphs of nullity one, *Discrete Mathematics* **181**(1-3) (1998), 193–211.
- [14] I. Sciriha, P. W. Fowler, Nonbonding Orbitals in Fullerenes: Nuts and Cores in Singular Polyhedral Graphs, *J.Chem.Inf.Model.* **47** (2007), 1763–1775.
- [15] I. Sciriha, Maximal core size in singular graphs, *Ars Mathematica Contemporanea* **2**(2) (2009), 217–229.
- [16] N. Trinajstić, *Chemical Graph Theory*, Vols. I and II, CRC Press, Boca Raton, 1983.
- [17] R. Varga, *Matrix Iterative Analysis*, Springer–Verlag, Berlin, 2000.