



An analytic effective model for hairy black holes

Yves Brihaye^{a,*}, Thomas Delplace^a, Carlos Herdeiro^b, Eugen Radu^b

^a Physique-Mathématique, Université de Mons-Hainaut, Mons, Belgium

^b Departamento de Física da Universidade de Aveiro and Centre for Research and Development in Mathematics and Applications (CIDMA), Campus de Santiago, 3810-183 Aveiro, Portugal



ARTICLE INFO

Article history:

Received 10 April 2018

Accepted 8 May 2018

Available online xxxx

Editor: M. Cvetič

ABSTRACT

Hairy black holes (BHs) have macroscopic degrees of freedom which are not associated with a Gauss law. As such, these degrees of freedom are not manifest as quasi-local quantities computed at the horizon. This suggests conceiving hairy BHs as an interacting system with two components: a “bald” horizon coupled to a “hairy” environment. Based on this idea we suggest an effective model for hairy BHs – typically described by numerical solutions – that allows computing *analytically* thermodynamic and other quantities of the hairy BH in terms of a fiducial bald BH. The effective model is *universal* in the sense that it is only sensitive to the fiducial BH, but not to the details of the hairy BH. Consequently, it is only valid in the vicinity of the fiducial BH limit. We discuss, quantitatively, the accuracy of the effective model for asymptotically flat BHs with synchronised hair, both in $D = 4$ (including self-interactions) and $D = 5$ spacetime dimensions. We also discuss the applicability of the model to synchronised BHs in $D = 5$ asymptotically AdS and static $D = 4$ coloured BHs, exhibiting its limitations.

© 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The 1970s represent a golden era for the theoretical study of black holes (BHs). At the classical level it was understood that electrovacuum BHs are remarkably featureless, being completely classified by a small number of *macroscopic* independent degrees of freedom: mass, angular momentum and electric (possibly also magnetic) charge – see [1] for a review. At the quantum level, on the other hand, the visionary works of Bekenstein [2] and Hawking [3] heralded BHs as gateways into the realm of quantum gravity, by showing they are thermodynamical objects, and, in particular that they have an entropy, geometrically computed as the horizon area. Understanding and counting the *microscopic* degrees of freedom associated to this entropy became a primary challenge for any quantum gravity candidate theory. Two decades later, some remarkable success was obtained within String Theory, wherein, starting with [4,5], it was possible to identify and count the microscopic degrees of freedom that explain the classical geometric entropy, defined by the BHs macroscopic degrees of freedom, albeit only in some particular classes of BHs.

The macroscopic simplicity of electrovacuum BHs suggested the “no-hair” conjecture [6]: that the endpoint of gravitational collapse

is a stationary BH completely described by a small set of macroscopic degrees of freedom, all of which should be associated to Gauss laws. In particular, this means such degrees of freedom are manifest at the horizon and can be computed therein as quasi-local quantities (e.g. the Komar integrals [7] associated to the mass and angular momentum of stationary axisymmetric BHs and the Gauss law associated to the electric charge). This, in turn, ties up nicely with the microscopic picture, and the view that the horizon contains all relevant BH information.

The discovery of “hairy” BHs in a variety of models (see e.g. [8–10] for reviews) has overshadowed this conceptually simple picture. These BHs have extra macroscopic degrees of freedom not associated to a Gauss law. Therefore they do not seem to be associated to any quasi-local conserved quantity computable at the horizon level. This raises interesting questions, on how the microscopic description of the BH captures these extra macroscopic degrees of freedom, but it also suggests an effective model for obtaining an (in general) analytic approximation for physical and thermodynamical quantities of the hairy BHs associated to the horizon [11].

The basic idea of the effective model sketched in [11] (and suggested by the numerical evolutions in [12]), therein called *quasi-Kerr horizon model*, is that due to the absence of further local charges at the horizon, the horizon of the hairy BH is well approximated by the horizon of a fiducial bald BH but with different parameters. In a sense, the hairy BH can be conceived as a cou-

* Corresponding author.

E-mail address: Yves.Brihaye@umons.ac.be (Y. Brihaye).

pled system of a bald horizon with an external “hair” environment. Naturally, the system is interacting and the non-linearities of the underlying gravity–matter system introduce a non-trivial deformation of the “bald” horizon. But in the feeble hair regime, when only a small percentage of the overall spacetime energy is contained in the matter field, these non-linearities are expected to be small, and the horizon should still behave as that of the bald fiducial BH, but with shifted parameters to take into account the mass and angular momenta that is no longer inside the horizon but rather in the matter environment. One could expect such simple model to yield errors in the thermodynamics quantities of the order of the deviation from the fiducial bald BH. The findings in [11], however, revealed that this effective model gives an unexpectedly good approximation, sometimes with deviations of $\sim \mathcal{O}(1\%)$ even for fairly large deviations from the bald BH, e.g., when $\sim \mathcal{O}(30\%)$ of the spacetime energy is stored in the matter field.

The purpose of this paper is to investigate the applicability and accuracy of the model by considering further examples of hairy BHs. Thus, after reviewing the assumptions, basic statement and corollaries of the effective model in Section 2, we consider two applications in Section 3: we apply it to Kerr BHs with synchronised hair and self-interactions [13] in Section 3.1 and to five dimensional ($D = 5$) Myers–Perry BHs with synchronised hair [14] in Section 3.2. In both these examples the effective model performs well. In the $D = 4$ case the accuracies are comparable to those described at the end of the last paragraph. In the $D = 5$ case, there is a mass gap between the hairy BHs and the fiducial BH. This means that the fiducial BH geometry is never approached globally, but only locally. In this case we find that, even for very hairy BHs, for which $\sim \mathcal{O}(90\%)$ of the spacetime energy is stored in the matter field, the model can yield errors of $\sim \mathcal{O}(1\%)$ for some physical quantities. To exhibit also the limitations of the effective model, we consider in Section 4 two further applications: to the $D = 5$ AdS Myers–Perry BHs with synchronised hair [15] and to the coloured BHs in Einstein–Yang–Mills theory [16]. With these applications, we illustrate either difficulties in the formalism, or unimpressive accuracies. In Section 5 we present some final remarks, in particular speculating about the underlying reason for the good accuracy of the model in the case of asymptotically flat BHs with synchronised hair.

2. The general framework

2.1. Komar integrals and Smarr relation

We consider a general model in $D \geq 4$ spacetime dimensions, consisting of Einstein’s gravity minimally coupled to some matter fields ψ described by a Lagrangian density \mathcal{L}_m

$$S = \int d^D x \sqrt{-g} \left[\frac{R}{16\pi} + \mathcal{L}_m \right], \quad (1)$$

where R is the spacetime Ricci scalar. Here and below we use geometrised units, setting Newton’s constant and the speed of light to unity: $G = 1 = c$.

In this work we shall be interested in stationary space-times with N -azimuthal symmetries, where $N = 1, 2$, for $D = 4, 5$. This implies the existence of $N + 1$ commuting Killing vectors, $\xi \equiv \partial_t$, and $\eta^{(k)} \equiv \partial_{\varphi_k}$, for $k = 1, \dots, N$.

Assuming asymptotic flatness, the total (or ADM) mass M and total angular momenta $J_{(k)}$ of the configurations are obtained from Komar integrals [7] (see also, e.g. [17]), at spatial infinity, associated with the corresponding Killing vector fields

$$M = -\frac{1}{16\pi} \frac{D-2}{D-3} \int_{S_{\infty}^{D-2}} \alpha, \quad J^{(k)} = \frac{1}{16\pi} \int_{S_{\infty}^{D-2}} \beta^{(k)}, \quad (2)$$

with

$$\begin{aligned} \alpha_{\mu_1 \dots \mu_{D-2}} &\equiv \epsilon_{\mu_1 \dots \mu_{D-2} \rho \sigma} \nabla^\rho \xi^\sigma, \\ \beta^{(k)}_{\mu_1 \dots \mu_{D-2}} &\equiv \epsilon_{\mu_1 \dots \mu_{D-2} \rho \sigma} \nabla^\rho \eta^{(k)\sigma}. \end{aligned} \quad (3)$$

We are mainly interested in BH solutions with a regular event horizon geometry (without any restrictions on its topology, which for $D > 4$ can be non-spherical [18–20]). This horizon \mathcal{H} has an associated (hyper)area of its spatial sections, A_H , and a temperature T_H ; there are also N horizon angular velocities $\Omega_{H(k)}$ associated with the N -azimuthal symmetries.

Using Komar integrals computed at the event horizon, one also defines a horizon mass M_H and a set of N horizon angular momenta $J_H^{(k)}$,

$$M_H = -\frac{1}{16\pi} \frac{D-2}{D-3} \int_{\mathcal{H}} \alpha, \quad J_H^{(k)} = \frac{1}{16\pi} \int_{\mathcal{H}} \beta^{(k)}. \quad (4)$$

Then the following Smarr type mass formulae [21] hold: for the horizon quantities we have

$$\frac{D-3}{D-2} M_H = \frac{1}{4} T_H A_H + \sum_{(k)} \Omega_{H(k)} J_H^{(k)}, \quad (5)$$

whereas for the bulk quantities

$$M = \frac{D-2}{D-3} \left[\frac{1}{4} T_H A_H + \sum_{(k)} \Omega_{H(k)} (J^{(k)} - J_{(\psi)}^{(k)}) \right] + M_{(\psi)}. \quad (6)$$

In the above relations, $M_{(\psi)}$, $J_{(\psi)}^{(k)}$ are the energy and angular momenta stored in the matter fields, with

$$M = M_H + M_{(\psi)}, \quad J^{(k)} = J_H^{(k)} + J_{(\psi)}^{(k)}. \quad (7)$$

Via the Einstein equations, $M_{(\psi)}$ and $J_{(\psi)}^{(k)}$ can be expressed as volume integrals for the appropriate components of the energy–momentum tensor (see e.g. [17]).

In addition to the above Smarr relations, the configuration should satisfy the first law of BH thermodynamics [22],

$$dM = \frac{1}{4G} T_H dA_H + \sum_{(k)} \Omega_{H(k)} dJ^{(k)} + \mathcal{W}, \quad (8)$$

where \mathcal{W} denotes the work term(s) associated with the matter fields. In particular, for vacuum solutions, the following relation holds

$$dM_H = \frac{1}{4} T_H dA_H + \sum_{(k)} \Omega_{H(k)} dJ_H^{(k)}. \quad (9)$$

2.2. The effective model

We now turn into the assumptions of the effective model [11], its statement and its corollaries.

Assumption 1 (Fiducial “bald” BH). One defines a vacuum *fiducial* BH solution¹ which is approached smoothly as $M_{(\psi)} \rightarrow 0$, $J_{(\psi)}^{(k)} \rightarrow 0$ (i.e. with the same symmetries and horizon structure as the non-vacuum solution). Moreover, at least in all cases discussed in this work, the horizon quantities of the fiducial BH have known (in

¹ In $D = 4$ the fiducial solution is obviously Kerr. But in higher dimensions, there can be different solutions for the same global charges and the horizon topology [18–20], thus requiring the definition of the fiducial solution.

closed form) expressions in terms of the global charges (macroscopic degrees of freedom); schematically these are:

$$\begin{aligned} A_H &= A_H(M, J^{(k)}), & T_H &= T_H(M, J^{(k)}), \\ \Omega_H^{(k)} &= A_H(M, J^{(k)}). \end{aligned} \quad (10)$$

It will be useful to define a set of $N + 1$ “hairiness” parameters $\mathfrak{h} \equiv (p, q^{(k)})$, which measure the deviation of the hairy BH from the fiducial solution

$$p \equiv \frac{M_{(\psi)}}{M}, \quad q^{(k)} \equiv \frac{J_{(\psi)}^{(k)}}{J^{(k)}}. \quad (11)$$

Assumption 2 (“Hair” matter field). We assume that there is no work term associated with the matter fields in the 1st law (8). This assumption guarantees the matter is adding “hair”, without a Gauss law associated, and it will not introduce a different global charge (another macroscopic degree of freedom), that can be computed at the horizon, as, for instance, an electromagnetic “matter” field would (electric charge). This still allows the “hair” matter field to have a conserved Noether charge, see e.g. [26], which is associated to a global, rather than gauge, symmetry.

Statement of the model The horizon quantities $\mathcal{Q} = (A_H, T_H, \Omega_H^{(k)})$ of the non-vacuum BH are still given by those of the corresponding fiducial BH, relation (10), expressed, however, in term of the horizon mass and angular momenta of the non-vacuum solution. That is, one considers the substitution

$$\mathcal{Q}^{(\text{fid})}(M, J^{(k)}) \longrightarrow \mathcal{Q}^{(\text{HBH})}(M_H, J_H^{(k)}). \quad (12)$$

Corollary 1 (Analytic formulas for horizon quantities). From (7) and (11)

$$M_H = (1 - p)M, \quad J_H^{(k)} = (1 - q^{(k)})J^{(k)}; \quad (13)$$

then, the horizon quantities of the hairy BH (12) can be expressed as

$$\mathcal{Q}^{(\text{HBH})}(M, J^{(k)}; \mathfrak{h}). \quad (14)$$

If the horizon quantities of the fiducial BH are expressed by analytic formulas $\mathcal{Q}^{(\text{fid})}(M, J^{(k)})$ so will be the horizon quantities of the hairy BH, eqs. (14).

Corollary 2 (Analytic relation between “hairiness parameters”). The assumption that the horizon is still described by the vacuum reference BH, together with the 1st law (8) (without a work term \mathcal{W}) implies that the matter fields satisfy the relation, for rotating BHs,²

$$dM_{(\psi)} = \sum_{(k)} \Omega_{H(k)} dJ_{(\psi)}^{(k)}. \quad (15)$$

Then we formally integrate the above relation treating $\Omega_{H(k)}$ as a set of input parameter (which is justified, since it belong to a different subsystem). This results in

$$M_{(\psi)} = \sum_{(k)} \Omega_{H(k)} J_{(\psi)}^{(k)}, \quad \text{or} \quad M - M_H = \sum_{(k)} \Omega_{H(k)} (J^{(k)} - J_H^{(k)}). \quad (16)$$

Using $M_{(\psi)} = pM$, $J_{(\psi)}^{(k)} = q^{(k)} J^{(k)}$, (16) becomes

$$p = \sum_{(k)} \Omega_{H(k)} q^{(k)} \frac{J^{(k)}}{M}. \quad (17)$$

Summary Relations (14) and (17) are the central results of the proposed effective model. After considering them together, one can eliminate p and arrive at the following set of relations

$$A_H(M, J^{(k)}; q^{(k)}), \quad T_H(M, J^{(k)}; q^{(k)}), \quad \Omega_H^{(k)}(M, J^{(k)}; q^{(k)}). \quad (18)$$

The explicit form of these relations is case dependent; the approach, however, is general. These analytic equations can be compared with the numerical solution of the hairy BH to check the domain of validity of the effective model.

As a final remark, we observe that instead of eliminating p , eq. (17) can be used to eliminate instead one of the parameters $q^{(k)}$, which results in an equivalent form of (18). In practise, as usual in BH physics, it is natural to work in units set by the BH mass (i.e. with normalised quantities).

3. Applications of the effective model

In this section we shall consider specific applications of the effective model. The simplest such application is found for the following scalar matter content:

$$\mathcal{L}_m = -(\partial_a \Phi)^\dagger (\partial^a \Phi) - U(|\Phi|), \quad (19)$$

where Φ is in general a scalar multiplet and $U(|\Phi|)$ is a self-interactions potential. The Einstein–Klein–Gordon equations possesses both solitons and hairy BH solutions. When the interaction potential includes a mass term, there are everywhere regular, asymptotically flat, stationary solitonic solutions known as boson stars [23]. Rotating boson stars [24], in particular, arise as a particular limit of Kerr BHs with (scalar) synchronised hair [25,26]. Both the solitonic rotating boson stars [27] and the hairy BHs [14,28] can be generalised to $D = 5$.

In [11] the effective model was already applied to the simplest scalar [25] and vector BHs [29] with synchronised hair. In section 3.1 we shall apply the effective model to the $D = 4$ hairy BHs with self-interactions obtained in [13]. In section 3.2 we shall apply it to the $D = 5$ hairy BHs [14].

3.1. $D = 4$ BHs with synchronised scalar hair

3.1.1. Predictions of the effective model

Consider the $D = 4$ Kerr BHs with synchronised hair [13,25,26] the reference solution is the Kerr metric [30], for which $N = 1$. Then the statement of the model, cf. (12), is that the horizon quantities of the hairy BHs obey:

$$\begin{aligned} \Omega_H &= \frac{M_H}{2J_H} (1 - \chi), & A_H &= 8\pi M_H^2 (1 + \chi), \\ T_H &= \frac{\chi}{4\pi M_H (1 + \chi)}, & \chi &= \sqrt{1 - \frac{J_H^2}{M_H^4}}. \end{aligned} \quad (20)$$

Next, we use (13), which in this case is simply,

$$M_H = (1 - p)M, \quad J_H = J(1 - q), \quad (21)$$

to obtain the specific form of (17), which reads

$$pM = \Omega_H J q. \quad (22)$$

We now introduce a set of reduced parameters, normalising the corresponding physical parameter by the ADM mass,

² The case of static BHs is simpler and will be illustrated in Section 4.2.

$$j \equiv \frac{J}{M^2}, \quad a_H \equiv \frac{A_H}{16\pi M^2}, \quad w_H \equiv \Omega_H M, \quad t_H \equiv 8\pi M T_H. \quad (23)$$

Then (22) takes the compact form

$$p = w_H j q. \quad (24)$$

Replacing (20) in (23), making use of (21) and choosing, via (24), (p, w_H) as independent parameters, one arrives at the following expressions

$$q = p \frac{1 + 4(1-p)^2 w_H^2}{p + 4(1-p)^2 w_H^2}, \quad j = \frac{p + 4(1-p)^2 w_H^2}{w_H [p + 4(1-p)^2 w_H^2]}, \quad (25)$$

$$a_H = \frac{(1-p)^2}{1 + 4(1-p)^2 w_H^2}, \quad t_H = \frac{1 - 4(1-p)^2 w_H^2}{1-p}, \quad (26)$$

which can be taken as the predictions of the effective model, in the spirit of eq. (18). It is also possible to show that

$$\frac{p}{q} = 1 - \frac{a_H}{1-p}, \quad (27)$$

which implies, according to the effective model, that $p < q$ for a hairy BH.

Finally, we observe that it is possible to express all quantities solely in terms of the hairiness parameters (p, q) :

$$w_H = \frac{1}{2(1-p)} \sqrt{\frac{p(1-q)}{q-p}}, \quad j = \frac{2(1-p)\frac{p}{q}}{\sqrt{\frac{p}{q} \frac{1-q}{1-\frac{p}{q}}}}, \quad (28)$$

$$a_H = (1-p) \left(1 - \frac{p}{q}\right), \quad t_H = \frac{1 - \frac{p}{q}(2-q)}{(1-p) \left(1 - \frac{p}{q}\right)}.$$

3.1.2. Validating the effective model

For $D = 4$, we restrict our study here to the simplest case of a scalar singlet (single complex field), with

$$\Phi = \phi(r, \theta) e^{i(m\varphi - \omega t)}. \quad (29)$$

We will also focus on the scalar field potential [13]

$$U(|\Phi|) = \mu^2 |\phi|^2 + \lambda |\phi|^4, \quad (30)$$

where μ is the scalar field mass and the self-coupling is positive, $\lambda > 0$. This complements the study in [11] which addresses the case $\lambda = 0$.

In Fig. 1 we exhibit the relative errors $|1 - \mathcal{Q}^{(\text{model})}/\mathcal{Q}^{(\text{num})}|$ for the quantities $\mathcal{Q} = (j, q, t_H)$ in terms of the parameters (p, w_H) . Observe that the self-interactions only affect $\mathcal{Q}^{(\text{num})}$; the effective model prediction $\mathcal{Q}^{(\text{model})}$ is insensitive to the self-interactions. The overall errors in this domain are comparable to those in [11], which is somewhat expected because the effect of the self interactions is small (for the numerical solutions) in the region studied, which is in the vicinity of the existence line ($p = 0 = q$), wherein the hairy BHs reduce to Kerr BHs. Remarkably, even for fairly large values of p , such as $p \sim 0.3$, the effective models gives an error of only a few percent, for low ω_H . For the reduced angular momentum the error is of $\sim 1\%$ within the whole domain $p \in [0, 0.3]$, $\omega_H \in [0, 0.5]$!

3.2. $D = 5$ BHs with synchronised scalar hair

We now consider Myers–Perry BHs with synchronised hair [14,28]. $D = 5$ rotating BHs with scalar hair possess, generically, two independent angular momenta and may even have a more general topology of the event horizon [31]. Here, we shall focus on the case with two equal angular momenta [14]. In $D = 5$ a qualitative difference with respect to the $D = 4$ case is that there is a mass gap between the hairy BHs and the fiducial model, which is taken to be the Myers–Perry vacuum solution [32], with two equal angular momenta.

3.2.1. Predictions of the effective model

$D = 5$ BHs with two equal-magnitude angular momenta have

$$J^1 = J^2 \equiv J, \quad \Omega_{H(1)} = \Omega_{H(2)} \equiv \Omega_H. \quad (31)$$

We also consider only the case with a spherical horizon topology. For such solutions the isometry group is enhanced from $\mathbb{R}_t \times U(1)^2$ to $\mathbb{R}_t \times U(2)$, where \mathbb{R}_t refers to time translations. This symmetry enhancement allows to factorise the angular dependence and thus leads to ordinary differential equations (not partial differential equations).

The fiducial solution is, in this case, the double spinning Myers–Perry BH. Then the statement of the model, cf. (12), is that the horizon quantities of the hairy BHs obey:

$$\Omega_H = \frac{M_H}{3J_H} (1 - \bar{\chi}), \quad A_H = \frac{16}{3} \sqrt{\frac{2\pi}{3}} M_H^{3/2} (1 + \bar{\chi}),$$

$$T_H = \sqrt{\frac{3}{8\pi M_H}} \frac{\bar{\chi}}{1 + \bar{\chi}}, \quad \bar{\chi} \equiv \sqrt{1 - \frac{27\pi J_H^2}{8M_H^3}}. \quad (32)$$

Again, it is convenient to define quantities normalised w.r.t. the ADM mass of the BHs. The $D = 5$ usual conventions in the literature are

$$j = \frac{3}{2} \sqrt{\frac{3\pi}{2}} \frac{J}{M^{3/2}}, \quad a_H = \frac{3}{32} \sqrt{\frac{3}{2\pi}} \frac{A_H}{M^{3/2}}, \quad w_H = \sqrt{\frac{8}{3\pi}} \Omega_H \sqrt{M},$$

$$t_H = 4 \sqrt{\frac{2\pi}{3}} T_H \sqrt{M}. \quad (33)$$

In the absence of hair, i.e. for $p = q = 0$, corresponding to a Myers–Perry BH, the following relations hold

$$j = \frac{2w_H}{1 + w_H^2}, \quad a_H = \frac{1}{1 + w_H^2}, \quad t_H = 1 - w_H^2, \quad (34)$$

with $0 \leq w_H \leq 1$; the limits correspond, respectively, to the Schwarzschild–Tangherlini [33] and extremal Myers–Perry BHs.

Repeating the procedure described in the previous section, choosing again (p, w_H) as the independent parameters, yields the following simple expressions

$$q = 3p \frac{1 + (1-p)w_H^2}{3p + (4-p)(1-p)w_H^2}, \quad j = \frac{3p + (4-p)(1-p)w_H^2}{2w_H(1 + (1-p)w_H^2)}, \quad (35)$$

$$a_H = \frac{(1-p)^{3/2}}{1 + (1-p)w_H^2}, \quad t_H = \frac{1 - (1-p)w_H^2}{\sqrt{1-p}}. \quad (36)$$

Again, these can be regarded as the predictions of the effective model, in the spirit of eq. (18).




Fig. 1. The relative errors are shown in a strip on the (p, w_H) -plane for reduced angular momentum j (left panel), hairiness parameter q (center) and reduced temperature t_H (right panel) for $D = 4$ BHs with self-interacting scalar hair, with $\lambda = 100$. The Kerr limit ($p = 0$) is the left vertical axis.




Fig. 2. The relative errors are shown in a strip on the (p, w_H) -plane for reduced angular momentum j (left panel), hairiness parameter q (center) and reduced temperature t_H (right panel) for $D = 5$ BHs with scalar hair.

3.2.2. Validating the effective model

The matter content in this case, consistent with the aforementioned symmetries is found by taking Φ a complex doublet scalar field [27], with

$$\Phi = \phi(r)e^{-i\omega t} \begin{pmatrix} \sin\theta e^{i\varphi_1} \\ \cos\theta e^{i\varphi_2} \end{pmatrix}, \quad (37)$$

where $\theta \in [0, \pi/2]$, $(\varphi_1, \varphi_2) \in [0, 2\pi]$, and t denotes the time coordinate. Concerning the scalar field potential, we restrict our study to the simplest case with

$$U(|\Phi|) = \mu^2 \Phi^\dagger \Phi = \mu^2 \phi(r)^2 \quad (38)$$

where μ corresponds to the scalar field mass.

The corresponding $D = 5$ hairy BHs are discussed in [14]. Likewise their four dimensional counterparts, the solutions are supported by rotation and have no static limit. The main difference with respect to the $D = 4$ case is the absence of the existence line. That is, in the limit of vanishing Noether charge density, the scalar field becomes point-wise arbitrarily small and the geometry becomes, locally, arbitrarily close to that of a specific set of Myers–Perry BHs. There remains, however, a global difference with respect to the latter, manifest in a finite mass gap. Thus the hair of these $D = 5$ hairy BHs is intrinsically non-linear.

We have found that the effective model provides an accurate description of the hairy BHs in the vicinity of the “marginally bound set”. This is the natural $D = 5$ counterpart of the $D = 4$ existence line, wherein the matter field becomes point-wise arbitrarily small, even though the global charges do not vanish. This line is approached for $w \rightarrow \mu$.

In Fig. 2 we display the relative errors of the effective model for Myers–Perry BHs with synchronised hair. As before, these relative errors are $|1 - Q^{(\text{model})}/Q^{(\text{num})}|$, and are exhibited for the quantities $Q = (j, q, t_H)$ in terms of the parameters (p, w_H) . Impressively, even for extremely large values of p , such as $p \sim 0.98$, the effective model gives an error of less than one percent for a_H ,

for the ω_H range plotted! For j the error reaches $\sim \mathcal{O}(2\%)$ whereas for t_H is one order of magnitude larger $\sim \mathcal{O}(20\%)$. But even in the latter case it is considerably smaller than the naive expectation that the error should be of order of the deviation from the fiducial BH. As already mentioned, these hairy BHs do not continuously connect globally to the fiducial BH – there is always a mass gap [14], the minimum value of p for the hairy solutions being roughly $p \sim 0.93$.

4. Limitations of the effective model

4.1. $D = 5$ AdS BHs with synchronised scalar hair

The first example of BHs with synchronised hair in the literature are obtained in $D = 5$ AdS asymptotics [15]. It is interesting to inquire if the effective model may still provide a useful description in that case as well, which is qualitatively different due to the AdS asymptotics.

To address this question, we have performed an independent investigation of the hairy BHs in [15]. These solutions can be studied within the same framework in [14], already mentioned in Section 3.2.2. The presence of a mass term, however, is not necessary in this case, since the AdS asymptotics provides the necessary confining mechanism. Thus, following [15], we set $\mu = 0$ in what follows.

The domain of existence of the solutions is shown in Fig. 3. This plot manifests the striking analogy with the $D = 4$ asymptotically flat case. The domain of hairy BHs is bounded by a solitonic limit (red solid line – corresponding to AdS rotating boson stars, with equal angular momenta [27]), by the “bald limit”, defining the existence line (blue dotted line – corresponding to equal angular momenta Myers–Perry AdS BHs [34]) and a set of extremal (*i.e.* zero temperature, extremal hairy BHs).

Concerning the validity of the effective model, we shall now discuss how it holds only partially. Following the procedure in the previous sections, we first identify the fiducial BH has the equal




Fig. 3. The domain of existence in (w, M) -plane is shown for BHs with synchronised hairy with equal angular momenta, in AdS_5 . (For interpretation of the colours in the figure(s), the reader is referred to the web version of this article.)

angular momentum, Myers–Perry–AdS BH [34]. Then, the statement of the model, cf. (12), is that the horizon quantities of the hairy BHs obey the thermodynamical relations of the fiducial BH, in terms of the macroscopic degrees of freedom measured at the horizon. Recall that the horizon quantities (A_H, T_H, Ω_H) of the fiducial BH are

$$A_H = \frac{2\pi^2 L^3 x^4}{U}, \quad T_H = \frac{1}{2\pi LU} \left[1 + 2x^2 - \frac{2a^2}{L^2 x^2} (1 + x^2)^2 \right],$$

$$\Omega_H = \frac{a}{L^2} \left(1 + \frac{1}{x^2} \right), \quad (39)$$

where L is the AdS radius and

$$U \equiv \sqrt{x^2 \left(1 - \frac{a^2}{L^2} \right) - \frac{a^2}{L^2}}. \quad (40)$$

x and a are the two parameters of the solution which determine the horizon mass and angular momentum, according to:

$$M_H = \frac{3\pi L^2 x^4 (1 + 2x^2)}{8 U^2}, \quad J_H = \frac{\pi a L^2 x^4 (1 + x^2)}{4 U^2} \quad (41)$$

Our study reveals that, in a region close to the existence line, the hairy BHs are still well described by the effective model. This can be confirmed in Fig. 4 (left panel) where we show the relative errors for the horizon area for several values of the event horizon radius r_H , comparing the value computed from the effective model, $A_H^{(\text{model})}$, with the one computed from the numerical solutions $A_H^{(\text{num})}$. Similar results were found for T_H and Ω_H .

This case, however, makes clear a limitation in the applicability of the model. In fact, a crucial ingredient of the formalism is missing in the AdS case. Although one can formally define the hairiness parameters p, q (where M_ψ, J_ψ are computed as volume integrals), the splitting of the total mass and angular momentum as the sum of the horizon charges plus the contribution from the matter fields is not possible in the presence of a cosmological constant³

$$M \neq M_H + M_\psi, \quad J \neq J_H + J_\psi. \quad (42)$$

Thus A_H, T_H, Ω_H cannot be further re-expressed in terms of (p, q) and the effective description holds only *locally*, at the horizon level.

³ One can easily verify that $M \neq M_H$ even for a vacuum Schwarzschild AdS BH.




Fig. 4. The relative errors are shown for the event horizon area as a function of the hairiness parameter p , for several values of the event horizon radius.

4.2. $D = 4$ coloured BHs

The first clear counterexamples to the no-hair conjecture was found in 1989 in Einstein–Yang–Mills (EYM) theory [16], dubbed *coloured BHs*, in the wake of the discovery, by Bartnik and McKinnon, that solitonic horizonless configurations exist in the same model [35]. This contrasts with the case of Einstein–Maxwell theory, wherein the absence of solitons is rigorously established [37]. Coloured BHs are hairy since they have a non-trivial matter configuration outside their regular event horizon, and the solutions are no longer completely determined by their global charges. These BHs are unstable [38] already in the spherically symmetric case. A (rather old) review of these solutions can be found in [39].

The EYM-SU(2) “matter” Lagrangian density is

$$\mathcal{L}_m = -\frac{1}{2} \text{Tr} \{ F_{\mu\nu} F^{\mu\nu} \}, \quad (43)$$

where $F_{\mu\nu}$ is the field strength tensor $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + ie[A_\mu, A_\nu]$, e is the gauge coupling constant, the gauge potential is $A_\mu = \tau_a A_\mu^a/2$, and τ_a are the Pauli matrices. Here, μ, ν are space-time indices running from 1 to 4 and the gauge index a is running from 1 to 3.

The static, spherically symmetric EYM hairy BHs possess a purely magnetic YM field, with a single gauge potential $w(r)$. The magnetic flux at infinity vanishes and, as a result, there is a single global charge – the ADM mass M . These BHs consist of a 1-discrete parameter family of solutions, labelled by the integer $k \geq 1$, which denotes the node number of the function $w(r)$. In the following discussion we focus on the solutions with $k = 1$.

The domain of existence of these solutions can be qualitatively described as follows. Firstly, no upper bound exists for the horizon size. In the large horizon size limit, the Einstein equations decouple from the gauge sector, yielding a Yang–Mills system on a fixed Schwarzschild BH background. In this decoupling limit there is a known exact solution (in closed form) [36]. As the horizon size shrinks to zero, the Bartnik–McKinnon family of solitons is recovered. Further details can be found in [39].

For applying the effective model, the obvious fiducial metric is that of a Schwarzschild BH. Then, applying the formalism in Section 2 leads to the following predictions

$$a_H = (1 - p)^2, \quad t_H = \frac{1}{1 - p}. \quad (44)$$

Computing the relative errors, like before, for instance of the temperature one concludes the error is of the order of p – Fig. 5. This




Fig. 5. The (reduced) temperature-area diagram for EYM BHs is exhibited together with the relative errors for the temperature.

the unremarkable result one expects in general: that the error in the effective model is of the order of the deviation from the bald BH. Thus, in this case, the effective model is not particularly accurate.

5. Discussion and final remarks

In this paper we have discussed an effective model for computing, analytically, several quantities for hairy BHs, in terms of the corresponding quantities of a fiducial bald BH. This model, already considered in [11] for Kerr BHs with synchronised scalar [25] and vector [29] hair, was applied to two other examples of BHs with synchronised hair: the $D = 4$ BHs with scalar self-interactions [13] and to the $D = 5$ “hairy” Myers–Perry BHs [14] in section 3.2. In all these cases one finds that the relative errors in the quantities provided by the model can be considerably smaller than those one would naively expect, namely that these errors should be of the order of deviation from the fiducial solution.

To illustrate the limitations of the model, we have also considered two other examples in Section 4: the $D = 5$ AdS Myers–Perry BHs with synchronised hair [15] and to the coloured BHs in Einstein–Yang–Mills theory [16]. The former case shows that one step of the model, namely the splitting between horizon and ADM quantities may be subtle in non-asymptotically flat spacetimes; still one may use the model and find small errors, as in the asymptotically flat case. The latter case illustrates that the model may give errors of the order of the deviations from the fiducial BH, thus unimpressive.

In this set of applications, the example of coloured BHs is the exceptional one. All the remaining examples rely on the synchronisation mechanism to endow rotating BHs with hair. So, why is this effective model performing better for this sort of “hairy” BHs? A possible answer is that, in these cases, there is a separation of scales. The hair field has its largest amplitude not at the horizon but at some distance thereof (see e.g. [26]) – defining a different scale from that of the horizon –, unlike the coloured case, where it decreases away from the horizon. This suggests a more efficient decoupling between the two sub-systems (the “bald” horizon and the matter “hair”) occurs in this case, allowing the horizon to remain fiducial BH-like even for larger amplitudes of the matter field. It would be interesting to further explore this suggestion.

Acknowledgements

C.H. and E.R. acknowledges funding from the FCT-IF programme. This work was supported by the European Union’s Horizon 2020

research and innovation programme under the H2020-MSCA-RISE-2015 Grant No. StronGrHEP-690904, the H2020-MSCA-RISE-2017 Grant No. FunFiCo-777740 and by the CIDMA project UID/MAT/04106/2013. The authors would also like to acknowledge networking support by the COST Action GWverse CA16104.

References

- [1] P.T. Chrusciel, J. Lopes Costa, M. Heusler, *Living Rev. Relativ.* 15 (2012) 7, arXiv:1205.6112 [gr-qc].
- [2] J.D. Bekenstein, *Phys. Rev. D* 7 (1973) 2333.
- [3] S.W. Hawking, *Commun. Math. Phys.* 43 (1975) 199, Erratum: *Commun. Math. Phys.* 46 (1976) 206.
- [4] A. Strominger, C. Vafa, *Phys. Lett. B* 379 (1996) 99, arXiv:hep-th/9601029.
- [5] C.G. Callan, J.M. Maldacena, *Nucl. Phys. B* 472 (1996) 591, [https://doi.org/10.1016/0550-3213\(96\)00225-8](https://doi.org/10.1016/0550-3213(96)00225-8).
- [6] R. Ruffini, J.A. Wheeler, *Phys. Today* 24 (1) (1971) 30.
- [7] A. Komar, *Phys. Rev.* 113 (1959) 934.
- [8] J.D. Bekenstein, in: 2nd International A.D. Sakharov Conference on Physics, Moscow, 1996, pp. 216–219, arXiv:gr-qc/9605059.
- [9] C.A.R. Herdeiro, E. Radu, *Int. J. Mod. Phys. D* 24 (09) (2015) 1542014, arXiv:1504.08209 [gr-qc].
- [10] M.S. Volkov, arXiv:1601.08230 [gr-qc].
- [11] C.A.R. Herdeiro, E. Radu, *Phys. Rev. Lett.* 119 (26) (2017) 261101, arXiv:1706.06597 [gr-qc].
- [12] W.E. East, F. Pretorius, *Phys. Rev. Lett.* 119 (4) (2017) 041101, arXiv:1704.04791 [gr-qc].
- [13] C.A.R. Herdeiro, E. Radu, H. Rúnarsson, *Phys. Rev. D* 92 (8) (2015) 084059, arXiv:1509.02923 [gr-qc].
- [14] Y. Brihaye, C. Herdeiro, E. Radu, *Phys. Lett. B* 739 (2014) 1, arXiv:1408.5581 [gr-qc].
- [15] O.J.C. Dias, G.T. Horowitz, J.E. Santos, *J. High Energy Phys.* 1107 (2011) 115, arXiv:1105.4167 [hep-th].
- [16] M.S. Volkov, D.V. Galtsov, *JETP Lett.* 50 (1989) 346, *Pis'ma Zh. Eksp. Teor. Fiz.* 50 (1989) 312; H.P. Kuenzle, A.K.M. Masood-ul-Alam, *J. Math. Phys.* 31 (1990) 928; P. Bizon, *Phys. Rev. Lett.* 64 (1990) 2844.
- [17] P.K. Townsend, arXiv:gr-qc/9707012.
- [18] R. Emparan, H.S. Reall, *Phys. Rev. Lett.* 88 (2002) 101101, arXiv:hep-th/0110260.
- [19] R. Emparan, H.S. Reall, *Living Rev. Relativ.* 11 (2008) 6, arXiv:0801.3471 [hep-th].
- [20] B. Kleihaus, J. Kunz, E. Radu, *Phys. Lett. B* 718 (2013) 1073, arXiv:1205.5437 [hep-th].
- [21] L. Smarr, *Phys. Rev. Lett.* 30 (1973) 71, Erratum: *Phys. Rev. Lett.* 30 (1973) 521.
- [22] J.M. Bardeen, B. Carter, S.W. Hawking, *Commun. Math. Phys.* 31 (1973) 161.
- [23] F.E. Schunck, E.W. Mielke, *Class. Quantum Gravity* 20 (2003) R301, arXiv:0801.0307 [astro-ph].
- [24] S. Yoshida, Y. Eriguchi, *Phys. Rev. D* 56 (1997) 762.
- [25] C.A.R. Herdeiro, E. Radu, *Phys. Rev. Lett.* 112 (2014) 221101, arXiv:1403.2757 [gr-qc].
- [26] C. Herdeiro, E. Radu, *Class. Quantum Gravity* 32 (14) (2015) 144001, arXiv:1501.04319 [gr-qc].
- [27] B. Hartmann, B. Kleihaus, J. Kunz, M. List, *Phys. Rev. D* 82 (2010) 084022, arXiv:1008.3137 [gr-qc].
- [28] C. Herdeiro, J. Kunz, E. Radu, B. Subagyo, *Phys. Lett. B* 748 (2015) 30, arXiv:1505.02407 [gr-qc].
- [29] C. Herdeiro, E. Radu, H. Rúnarsson, *Class. Quantum Gravity* 33 (15) (2016) 154001, arXiv:1603.02687 [gr-qc].
- [30] R.P. Kerr, *Phys. Rev. Lett.* 11 (1963) 237.
- [31] C. Herdeiro, J. Kunz, E. Radu, B. Subagyo, *Phys. Lett. B* 779 (2018) 151, arXiv:1712.04286 [gr-qc].
- [32] R.C. Myers, M.J. Perry, *Ann. Phys.* 172 (1986) 304.
- [33] F.R. Tangherlini, *Nuovo Cimento* 27 (1963) 636.
- [34] S.W. Hawking, C.J. Hunter, M. Taylor, *Phys. Rev. D* 59 (1999) 064005, arXiv:hep-th/9811056.
- [35] R. Bartnik, J. McKinnon, *Phys. Rev. Lett.* 61 (1988) 141.
- [36] H. Boutaleb-Joutei, A. Chakrabarti, A. Comtet, *Phys. Rev. D* 20 (1979) 1884.
- [37] T. Shiromizu, S. Ohashi, R. Suzuki, *Phys. Rev. D* 86 (2012) 064041, arXiv:1207.7250 [gr-qc].
- [38] Z.h. Zhou, N. Straumann, *Nucl. Phys. B* 360 (1991) 180.
- [39] M.S. Volkov, D.V. Gal'tsov, *Phys. Rep.* 319 (1999) 1, arXiv:hep-th/9810070.