



# Relationship between distress and physical activity in informal carers of patients with COPD

Nádia Hipólito, Adriana Ruivo, Sara Martins, Sofia Flora, Alda Marques, Dina Brooks, Cândida G. Silva, Filipa Januário, Joana Cruz  
European Respiratory Journal 2019 54: PA1245; DOI: 10.1183/13993003.congress-2019.PA1245

[Article](#)[Info & Metrics](#)

## Abstract

**Background:** Chronic obstructive pulmonary disease (COPD) can lead to an increase in patient's dependence on the informal carer and consequently to higher levels of distress. In the general population, higher levels of physical activity (PA) have been found to contribute to lower levels of distress. However, this relationship has been scarcely studied in carers of patients with COPD.

**Aim:** This study aimed to explore the relationship between distress and PA in informal carers of patients with COPD and the influence of caregiving duration.

**Methods:** Forty-one carers ( $62.4 \pm 10.1$  years, 90.2% female; 41.5% caring for patients  $>40$ h/week; patients'  $\text{FEV}_1 = 43.7 \pm 19.7\%_{\text{pred}}$ ) completed the Portuguese tool to assess distress related to caregiving (Informal Caregiver Burden Assessment Questionnaire [QASCI]; higher score meaning higher distress; 7 subscales); the Habitual Physical Activity Questionnaire (HPAQ) to assess PA; and questions related to the caregiving duration (h/week, years). Pearson's correlations and linear regressions were used.

**Results:** There was a negative moderate correlation between the QASCI ( $30.3 \pm 20.7$ ) and the HPAQ ( $5.1 \pm 1.2$ ) ( $r = -.517$ ;  $p = .01$ ). Correlations were also found between PA and some of the QASCI subscales (emotional burden  $r = -.500$ ; implications for personal life  $r = -.652$ ; financial burden  $r = -.471$ ; perception of efficacy and control

WE USE COOKIES ON THIS SITE TO ENHANCE YOUR USER EXPERIENCE

By clicking any link on this page you are giving your consent for us to set cookies.

[OK, I agree](#)[No, give me more info](#)

**Conclusions:** Higher PA levels may be related to decreased levels of distress in this population. Further research is needed.

Mental health    COPD - management    Physical activity

---

## Footnotes

Cite this article as: European Respiratory Journal 2019; 54: Suppl. 63, PA1245.

This is an ERS International Congress abstract. No full-text version is available. Further material to accompany this abstract may be available at [www.ers-education.org](http://www.ers-education.org) (ERS member access only).

Copyright ©the authors 2019

---

## We recommend

Is psychological stress associated with smoking and physical activity levels in patients in the CanCOLD Study?

Brandie Walker et al., European Respiratory Journal, 2017

Symptom experience in patients with COPD and their level of physical activity. Is there a good association?

Heleen Demeyer et al., European Respiratory Journal, 2012

Relationship between caregivers' burden, functional disability and quality of life in COPD patients

Ramón Romero Fernández et al., European Respiratory Journal, 2019

Living with COPD: Psychological distress among family carers

Daniela Figueiredo et al., European Respiratory Journal, 2012

Physical activity and its relation with exercise capacity in patients with early COPD

Hans Van Remoortel et al., European Respiratory Journal, 2011

Providing Care to Relatives With Mental Illness  
Sherilyn Chang et al., Medscape

Prospective study on lung adenocarcinoma - nintedanib plus docetaxel after progression on immune checkpoint inhibitor therapy

Future Oncology

Co-Morbid Depression in Patients With Acne Inversa

PracticeUpdate, 2013

MEDLINE Abstracts: Caregiver Burden and the Geriatric Patient  
Medscape

ERS 2017: Factors Leading to Thrombotic Events in COPD Are Elucidated

PracticeUpdate, 2017

---

Powered by **TREND MD**

**WE USE COOKIES ON THIS SITE TO ENHANCE YOUR USER EXPERIENCE**

By clicking any link on this page you are giving your consent for us to set cookies.

[OK, I agree](#)

[No, give me more info](#)