

their close relatives experienced the investigation and the subsequent social medicine intervention, and to give recommendations based on the results. The project is based on qualitative interviews with 11 families.

Background: The Coordinated Investigation Model of Dementia implies that the primary investigation is carried out in the primary sector by a general practitioner in cooperation with a local dementia nurse in accordance to a specified procedure. Further investigation may hereafter be carried out by specialists in the secondary sector. Especially after The Coordinated Investigation Model was evaluated in 2005 and found useful from a professional point of view, we found it relevant to investigate the experiences by the patients and their close relatives.

Results: The project shows that the patients and their close relatives don't relate to the model – but rather to the results of the investigation, responsiveness and comfort. The investigation in the secondary sector seems very important to the patients and their relatives, who generally expressed that important decisions and information were handled by the secondary sector. A follow-up interview in the primary sector after investigation, which is mandatory according to the model, was practiced to a very limited extent. The role of the local dementia nurse as a coordinator was widely recommended by the patients and their close relatives, since many have experienced an accessible support and a relevant help.

Recommendations: The primary sector's role in investigation and follow-up should be strengthened with the aim of executing the model with lowest effective cost. A formal agreement regarding follow-up should be implemented. The relatives should be more involved during both investigation period and in the socio-medical follow-up.

P1-10

The Effect of Learning Therapy on Improving The Cognition Function and Psychological , Behavior Symptoms among the Elderly with Dementia in Institution

Author: Professor Shu-Yuan Chao
Hungkung University

Co-Authors: Hsiao-Mei Chen, Li-Jane Tsai

The study used a quasi-experimental design research method to investigate the effect of learning therapy on the improvement of cognition and psychological, behavior symptoms among the elderly with dementia in institution. There were two institutions, in total 44 participants were recruited in this study, including 23 and 21 participants were assigned to experimental group and control group, respectively. The elderly in experimental group were invited to read aloud and conduct digital computation, 15 minutes per time, five days a week, and continuously for 3 months. The MMSE and NPI were used for measuring the improvement of

elder's cognition function, and psychological, behavior symptoms, before and after intervention. We applied multiple linear regression with the generalized estimating equation (GEE) statistic method to evaluate the effect of learning therapy on the cognition function and symptoms improvement after intervention. The results indicated significant improvement on experimental group elders, both cognition function and neuropsychiatric symptoms, achieving statistic level. The program could be the reference for the elderly care in institution.

Key words: Dementia, Learning therapy, Cognitive function, Elderly.

P1-11

Risk of falling in elderly patients with Chronic Obstructive Pulmonary Disease

Author: Dr. Cristina Jácome

Escola Superior de Saúde da Universidade de Aveiro

Co-Authors: Prof. Alda Marques, Prof. Daniela Figueiredo, Dr. Raquel Gabriel

Falls are a major problem among elderly adults that negatively impact on their functional independency and social interaction. The presence of chronic disease and impaired balance are well known risk factors for falling. Chronic Obstructive Pulmonary Disease (COPD) is one of the highest prevalent chronic diseases in elderly adults, which severely impacts on patients' mobility and balance. However, limited information is available on risk of falling in this population. Therefore, the aim of this study was to determine the risk of falling in elderly patients with COPD. A cross-sectional study was carried out with 31 elderly outpatients with COPD in the central region of Portugal. Socio-demographic data were collected via a questionnaire based on the International Classification of Functioning, Disability and Health checklist. Spirometry was performed to assess participants' respiratory function. Balance and risk of falling were assessed with the Timed Up and Go Test (TUG). Descriptive statistics, the Mann-Whitney U test and Chi-square test were applied using PASW Statistics version 18.0. Participants were mostly male (n=21;67.7%), with a mean age of 76.71±6.45 years old. According to the Global Initiative for Chronic Obstructive Lung Disease criteria, participants were at advanced COPD (FEV1%predicted 35.45±8.15%). Considering the TUG cutoff point of 14 seconds for predicting falls, 38.70% of the participants were at high risk of falling. Participants with more than 74 years old (58.06%) were significantly slower (15.40±6.18 vs 10.62±4.45;p=0.020) and presented a significantly higher risk of falling (55.6% vs 15.4%;p=0.023) compared with younger participants. Elderly patients with COPD are at high risk of falling, especially those over 74 years old. Hence, pulmonary rehabilitation, a recommended standard of care for patients with COPD, should include a specific component of balance training and strategies to prevent falling, in order respond to elderly patient's needs.