

MINIREVIEW

The role of non-standard translation in *Candida albicans* pathogenesis

Ana Rita Bezerra^{*,†}, Carla Oliveira, Inês Correia, Ana Rita Guimarães, Gonçalo Sousa, Maria João Carvalho, Gabriela Moura and Manuel A. S. Santos

Department of Medical Sciences, Institute of Biomedicine – iBiMED, University of Aveiro, 3810-193 Aveiro, Portugal

*Corresponding author: Department of Medical Sciences, Institute of Biomedicine – iBiMED, University of Aveiro, 3810-193 Aveiro, Portugal.

Tel: +351 234 247 336; E-mail: armbezerra@ua.pt

One sentence summary: *Candida albicans* uses the fidelity of protein synthesis to spawn novel virulence traits.

Editor: Miguel Teixeira

[†]Ana Rita Bezerra, <http://orcid.org/0000-0003-4613-1711>

ABSTRACT

Candida albicans typically resides in the human gastrointestinal tract and mucosal membranes as a commensal organism. To adapt and cope with the host immune system, it has evolved a variety of mechanisms of adaptation such as stress-induced mutagenesis and epigenetic regulation. Niche-specific patterns of gene expression also allow the fungus to fine-tune its response to specific microenvironments in the host and switch from harmless commensal to invasive pathogen. Proteome plasticity produced by CUG ambiguity, on the other hand is emerging as a new layer of complexity in *C. albicans* adaptation, pathogenesis, and drug resistance. Such proteome plasticity is the result of a genetic code alteration where the leucine CUG codon is translated mainly as serine (97%), but maintains some level of leucine (3%) assignment. In this review, we dissect the link between *C. albicans* non-standard CUG translation, proteome plasticity, host adaptation and pathogenesis. We discuss published work showing how this pathogen uses the fidelity of protein synthesis to spawn novel virulence traits.

Keywords: *Candida albicans*; non-standard translation; pathogenesis; drug resistance; genetic diversity; evolution

INTRODUCTION

The genetic code establishes the rules that govern the transfer of genetic information from nucleic acids to proteins. By establishing a unambiguous correspondence between codons and amino acids, the genetic code allows for stable inheritance of phenotypic variation produced by proteins upon which natural selection acts (Crick 1968). Despite this, numerous deviations to the standard genetic code have been discovered in both prokaryotes and eukaryotes (Knight, Freeland and Landweber 2001a; Koonin and Novozhilov 2009; Keeling 2016). Most of these alterations occurred in mitochondria due to their small genomes and

independent translation machinery—ribosomes and tRNAs (Knight, Landweber and Yarus 2001b; Ling, O'Donoghue and Soll 2015). Although alterations in nuclear genomes are less common, they include sense and nonsense codon reassignments, as well as codon unassignments (Miranda, Silva and Santos 2006; Kollmar and Muhlhausen 2017), with sense-to-sense codon reassignments having the highest potential to boost proteome plasticity.

The only reported cases of sense-to-sense codon reassignment in nuclear genomes invariably involves the leucine CUG codon. The identity of this codon changed three times during

Received: 1 March 2021; Accepted: 20 May 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

the evolution of budding yeasts, with two clades translating the CUG codon as serine (CUG-Ser), one clade translating it as alanine (CUG-Ala) and two clades translating it canonically as leucine (CUG-Leu) (Krassowski et al. 2018). Among the CUG-Ala species are *Pachysolen* and *Nakazawaea*, which are mostly used in biotechnological applications (Muhlhausen et al. 2016; Riley et al. 2016). Most species with biomedical relevance belong to the CTG-clade that comprise the most pathogenic *Candida* species (Fitzpatrick et al. 2006; Butler et al. 2009). Within this clade, *C. cylindracea* translates CUGs as serine only, while *C. guilliermondii*, *C. dubliniensis*, *C. zeylanoides*, *C. tropicalis* and *C. albicans* ambiguously decode the CUG codon as serine and leucine (Tuite and Santos 1996; M. A. Santos et al. 1997; Suzuki, Ueda and Watanabe 1997; M. A. Santos et al. 2011). Another example of dual codon identity occurs in the Ascoidea-clade species, where Ascoidea asiatica translates CUG stochastically as serine or leucine. In this species, residues encoded by CUG codons in key structural sites are strictly avoided (Muhlhausen et al. 2018).

Unlike *A. asiatica*, CUG-encoded residues may exist at functionally relevant positions in *C. albicans* and other CTG-clade spp. Most residues are located at the protein surface, where both leucine and serine can be incorporated without major impact on protein structure or function, but a few are located in positions where a serine or other polar amino acid is conserved in homologous proteins (Rocha et al. 2011). This is particularly intriguing because inaccurate production of proteins is generally viewed as a nuisance to biological systems (Kapur and Ackerman 2018; M. Santos et al. 2018). Nonetheless, recent data challenged these assumptions and strengthen the idea that protein mistranslation is a double-edged sword with potentially adaptive functions (Ribas de Pouplana et al. 2014; Schwartz and Pan 2017). Several studies show that, under stress, increased dual-translation of a codon enhances microbial fitness and modulates host-microbe interactions (Carlson et al. 2009; Netzer et al. 2009; Evans et al. 2018).

The concept of adaptive translation is greatly exemplified in *C. albicans*, where natural dual-translation of CUGs (which are normally translated as ~3% Leu and ~97% Ser but can change) expands the proteome exponentially (6438 CUG-containing genes can potentially encode 283 billion proteins, under a 50%–50% scenario) (Gomes et al. 2007; Rocha et al. 2011; M. A. Santos et al. 2011). The biological implications of this are profound, as each cell may contain a unique combination of protein molecules, generating an enormous biological complexity in this important human fungal pathogen. In this review, we dissect how protein diversity caused by variable translation of the CUG codon is emerging as a new key player in *C. albicans* adaptation, pathogenesis and drug resistance.

NON-STANDARD TRANSLATION IN CANDIDA SPP

At least 31 *Candida* spp. are known to cause disease in humans. Although the incidence of *Candida* spp. varies geographically, globally it is estimated that ~92% of the infections are caused by only five species: *C. albicans*, *C. glabrata*, *C. tropicalis*, *C. parapsilosis* and *C. krusei* (*Pichia kudriavzevii*) (Pfaller et al. 2010; Guinea 2014; Gabaldon et al. 2016). *C. albicans* is the most common pathogen isolated, but the prevalence of non-*albicans* species has been increasing in the past years. Other increasingly important causes of candidiasis include: *C. lusitaniae*, *C. guilliermondii*, *C. dubliniensis*, *C. kefyr* (*Kluyveromyces marxianus*), *C. famata* (*Debaryomyces hansenii*), *C. inconspicua*, *C. rugosa*, *C.*




Figure 1. Schematic illustration of the *C. albicans* tRNA_{CAG^{Ser}. **(A) Dual recognition of the *C. albicans* tRNA_{CAG^{Ser}.}** The *C. albicans* tRNA_{CAG^{Ser} contains identity elements for both LeuRS and SerRS. The LeuRS recognizes A₃₅ and m¹G₃₇ in the anticodon loop (red circles). The SerRS recognizes the discriminator base (G₇₃) and three GC base pairs of the extra-arm (blue circles). Notably, the presence of G₃₃, instead of U₃₃, in the anticodon U-turn decreases leucylation efficiency of the tRNA_{CAG^{Ser} in vitro (Miranda, Silva and Santos 2006). This double recognition by LeuRS and SerRS leads to the synthesis of two aminoacyl-tRNAs (charged with Ser or Leu) that compete for CUG codons. **(B) CUG decoding during mRNA translation.** Under standard growth conditions, *C. albicans* translates approximately 97% of CUG codons as Ser and approximately 3% as Leu (Miranda, Silva and Santos 2006; Gomes et al. 2007). Figure created with Biorender.com.}}}

dubliniensis, *C. norvegensis* (*Pichia norvegensis*) and the multi-drug resistant *C. auris* (Pfaller et al. 2010; Papon et al. 2013; Gabaldon et al. 2016; CDC 2019). Apart from *C. glabrata*, *C. krusei*, *C. kefyr*, *C. norvegensis*, and *C. inconspicua*, all *Candida* spp. listed belong to the CTG clade, a large group of yeasts that reassigned the leucine CUG codon (Butler et al. 2009; M. A. Santos et al. 2011; Papon et al. 2013; Du et al. 2020). Within this clade, *C. albicans*, *C. zeylanoides*, *C. lusitaniae*, *C. tropicalis*, *C. parapsilosis*, *C. guilliermondii* and *C. rugosa* naturally translate the leucine CUG codons as both serine and leucine (Suzuki, Ueda and Watanabe 1997; Gomes et al. 2007; M. A. Santos et al. 2011). This group of *Candida* spp. with dual CUG codon translation comprises most of the pathogenic *Candida* species, whose change of identity of the CUG codon is mediated by a novel tRNA_{CAG^{Ser} that contains identity elements for both the seryl- and leucyl-tRNA synthetases (SerRS and LeuRS, respectively). SerRS recognizes the 3 GC base pairs in the extra arm and the discriminator base (G₇₃), while LeuRS recognizes the A₃₅ and m¹G₃₇ in the anticodon loop of the tRNA molecule. This leads to double recognition by SerRS and LeuRS and to synthesis of two different aminoacyl-tRNAs, specifically a Ser-tRNA_{CAG^{Ser} and a Leu-tRNA_{CAG^{Ser}, that compete for CUG codons at the ribosome A-site during mRNA translation (Fig. 1). The mischarged Leu-tRNA_{CAG^{Ser} is not edited by the LeuRS nor}}}}

discriminated by the translation elongation factor 1 (eEF1A) and both Leu and Ser are incorporated into the proteome at CUG positions (Santos, Perreau and Tuite 1996; M. A. Santos et al. 1997; Gomes et al. 2007).

The evolutionary pathway of the CUG codon reassignment has been thoroughly discussed (Schultz and Yarus 1996; M. A. Santos et al. 2004; Miranda, Silva and Santos 2006; Moura, Paredes and Santos 2010), and is not in the scope of this review. However, these studies were the first to shed light on the mechanisms underlying the tolerance to codon ambiguity. Early reports noted that CUG is a rare codon in CTG clade species (Brown et al. 1991; Sugiyama et al. 1995; Suzuki, Ueda and Watanabe 1997) and tends to appear in non-housekeeping genes (Suzuki, Ueda and Watanabe 1997). Indeed, codon usage analysis revealed that the CUG codon is repressed in *C. albicans* (unlike *C. cylindracea*, the CUG 100% Ser decoder, which uses CUG as the preferred Ser codon) and CUG codons appear at serine codon sites in *S. cerevisiae* proteins rather than in Leu CUG codon positions (Massey et al. 2003; Gomes et al. 2007; Butler et al. 2009). This indicates that CUG reassignment and ambiguous decoding erased most of the CUG codons in the CTG clade ancestor and that new CUG sites arose in Ser codon positions (Massey et al. 2003; M. A. Santos et al. 2011).

Gomes and colleagues demonstrated that CUG is ambiguously decoded *in vivo* in *C. albicans* by quantifying Leu and Ser levels at CUG sites (Gomes et al. 2007), using a direct protein mass spectrometry (MS) analysis. The levels of leucine incorporation at CUG sites in *C. albicans* grown in standard growth conditions were 2.96% (Gomes et al. 2007). In the same study, recombinant *C. albicans* cells tolerated incorporation of 28% of Leu at CUG sites without significant impact on fitness. It was calculated that 13 074 CUG codons were distributed over more than a half (66%) of *C. albicans* genes. Considering that each gene has on average three CUGs, an ambiguity level of 2.96% could generate 6.7×10^6 novel proteins per cell in standard growth conditions. For this reason, it was considered that *C. albicans* proteome has a statistical nature (Gomes et al. 2007).

To evaluate the impact of CUG codon ambiguity on the proteome, Rocha and colleagues conducted a deep structural analysis of proteins containing CUG-encoded residues (Rocha et al. 2011). The alignment of 680 protein sequences with CUG-encoded residues in *C. albicans* with orthologs of six other fungal species revealed that 90% of CUG codons are located at non-conserved positions. The same pattern was observed for the other CTG clade species analysed, whereas CUGs in *S. cerevisiae* are evenly distributed in the protein sequences (Rocha et al. 2011). The distribution pattern of CUGs in the CTG clade species suggested that reintroduction of CUGs in the genomes, after the CUG reassignment, was selected to avoid protein misfolding caused by serine substitution for leucine. Proteins with CUG-encoded residues were uniformly distributed across diverse functional categories. However, the 10% of conserved CUG-encoded residues were mainly located in proteins involved in biological processes associated with virulence and pathogenesis, such as biofilm formation, mating, morphogenesis and adhesion. They also correlated with signal transduction, suggesting a pivotal role for CUG decoding ambiguity in pathogen-host interaction (Rocha et al. 2011).

To validate the functional impact of Ser-for-Leu substitutions, Rocha and colleagues determined the crystal structures of the two isoforms of SerRS (SerRS.Ser197 and SerRS.Leu197) (Rocha et al. 2011). SerRS has a CUG-encoded serine residue at position 197 located at the dimer interface. The Ser-for-Leu substitution induces a local rearrangement, which increases

slightly (27%) the activity of the SerRS.Leu197 isoform (Rocha et al. 2011; Robbins, Caplan and Cowen 2017). Similarly, Zhou and co-workers showed that the unique CUG-encoded residue present at position 919 in LeuRS is not conserved among eukaryotes, but is involved in tRNA binding and aminoacylation. Ser-for-Leu substitution at the unique CUG site increased ~30% the LeuRS activity without causing major structural alterations (Zhou et al. 2013). These studies suggest that leucine incorporation at CUG sites could be regulated by a fine balance of the SerRS and LeuRS isoforms activity. However, further investigation is needed to clarify this issue as we are still far from understanding the molecular mechanisms that regulate CUG codon ambiguity.

Another example of the impact of CUG codon ambiguity on protein function is the *C. albicans* eukaryotic translation initiation factor (eIF)4E. This cap-binding protein mediates mRNA binding to the ribosome and to translation initiation factors (Hinnebusch and Lorsch 2012). Feketová and colleagues showed that eIF4E has two variants due to CUG ambiguous decoding: eIF4E.Leu116 and eIF4E.Ser116 (Feketova et al. 2010). This residue is located in the surface of the protein, adjacent to the eIF4G binding region. Ser-for-Leu did not impair activity, but the eIF4E.Leu116 was temperature sensitive (Feketova et al. 2010), suggesting that *C. albicans* may reprogramme translation from CAP-dependent to CAP-independent mRNAs at high temperatures.

In another study, Leu-CUG translation decreased the stability and activity of the protein kinase Cek1 without major structural alterations. Cek1 contains a single CUG-encoded residue at a conserved position and is a key kinase of the MAPK cascade directly linked to morphogenesis in *C. albicans*. Incorporation of Ser at this CUG site induced the autophosphorylation of the conserved tyrosine residue of the Cek1 ²³¹TEY²³³ motif and increased its intrinsic kinase activity *in vitro*. Unlike the Ser-Cek1 variant, the Leu-Cek1 variant is not autophosphorylated at the ²³¹TEY²³³ motif within the kinase activation loop (Fraga et al. 2019). Therefore, these *in vitro* studies demonstrate that Leu/Ser-CUG isoforms of *C. albicans* proteins can be functional despite some differences in activity and/or specificity. It is also possible that CUG ambiguous decoding could affect protein-protein interactions, disrupting existing networks or producing new ones.

As the CTG clade shows some variability in reassignments and *C. albicans* tolerates high level of ambiguity at CUG sites (Gomes et al. 2007), the question if *C. albicans* tolerates ambiguity at other codons arose. To answer this, Simões and colleagues tested if *C. albicans* tolerates codon ambiguity at other codons (Simoes et al. 2016). For this, *C. albicans* strains were engineered to express tRNA_{UGA}^{Ser} genes with mutated anticodons. Several mutant tRNAs that were able to read codons encoding amino acids with different chemical properties, namely Leu, Ala, Gly, Lys, Thr and Tyr were engineered. This approach relied on the fact that eukaryotic seryl-tRNA synthetase (SerRS) does not recognize the tRNA Ser anticodon loop (Lenhard et al. 1999), so alterations in the anticodon do not affect tRNA serylation. In other words, each targeted codon was decoded by the mutant tRNA_{UGA}^{Ser} and by its cognate tRNA. It is important to note that the codons targeted in this study follow a similar codon usage as CUG. Interestingly, expression of the recombinant tRNAs had little impact on fitness (Simoes et al. 2016).

The high tolerance of *C. albicans* to codon ambiguity, alongside the functional consequences of Ser/Leu decoding mentioned above, raise the question of the overall implications of such ambiguity to *Candida* spp. biology. Recent studies provide

evidence that CUG ambiguity may indeed be relevant for adaptation. The double identity of the CUG codon creates statistical proteins whose implications in the many facets of *C. albicans* virulence will be dissected in the following sections.

PHENOTYPIC AND GENOMIC IMPACT OF CUG AMBIGUITY

Quantification of Leu incorporation at CUG sites using an MS-based reporter system revealed that wild-type *C. albicans* cells can increase CUG codon ambiguity up to ~5%, when grown in low pH conditions (Gomes et al. 2007). Although small, an increase of incorporation of Leu at CUG sites from the basal 3% to 5% could generate an additional 4.2×10^6 statistical polypeptides (Gomes et al. 2007). Remarkably, recombinant *C. albicans* strains encoding a yeast Leu tRNA_{CAG}^{Leu}, and incorporating 28% Leu at CUG sites displayed very high morphologic diversity, including ovoid opaque cells, pseudo-hyphal and hyphal morphologies. The phenotypes observed in highly ambiguous cells were similar to the phenotypes that are induced by environmental signals, such as high temperature, low pH and serum. Expression of a mutant *S. cerevisiae* Leu-tRNA_{CAG} in *C. albicans* strains increased Leu incorporation at CUG sites and exposed hidden phenotypic variability, including the same morphological changes reported by Gomes and colleagues, but also enhanced activity of extracellular hydrolases (aspartic proteinases and phospholipases). High-level CUG codon ambiguity also induced mating by upregulating the key regulator of white-opaque switching, which was reflected on the increase of mating competent opaque cells. Furthermore, it revealed up-regulated expression of genes involved in cell adhesion and hyphal development (Miranda et al. 2007). This is in line with other data showing that CUG-encoded residues are enriched at conserved positions in proteins associated with biofilm formation, morphological switching, and adhesion (Rocha et al. 2011).

Miranda and co-workers also showed that *C. albicans* cells that incorporated 28% Leu at CUG sites had higher adherence to fibronectin and gelatin (Miranda et al. 2013). This is relevant because *C. albicans* is a successful commensal of mammalian hosts due to its capacity to adhere to host constituents and to avoid immune clearance. The authors proposed two mechanisms to explain the high adherence phenotype. First, knowing that most CUG-encoded residues are located on the surface of proteins and that membrane and cell wall genes are particularly enriched in CUG codons (Rocha et al. 2011), CUG mistranslation may increase cell surface hydrophobicity due to increased substitution of polar Ser for the non-polar Leu. Second, presence of Leu at the two CUG-encoded residues (positions 379 and 433) in the adhesin Als3 altered its function and increased cell flocculation and substrate adhesion. Moreover, high level of Leu-CUG translation produced variability in the cell wall proteome that altered the cell surface exposure of 1,3- β -glucan and reduced phagocytosis by macrophages, compared to wild-type cells (Miranda et al. 2013). Together, these results indicate that proteome expansion generated by codon ambiguity can be a mechanism to generate cell surface variability and possibly to contribute to antigenic variation to avoid recognition by the immune system, which is particularly relevant in the fungal host-interaction.

To test the limits of CUG-Leu decoding and its role in virulence, a set of recombinant *C. albicans* strains incorporating between ~20% and ~98% of Leu at CUG sites were constructed (Bezerra et al. 2013). Although high-level Leu-CUG translating

strains displayed slower growth in rich medium, when grown with oxidative agents or antifungal drugs, particularly azoles, recombinant strains grew better than the wild-type strain (Bezerra et al. 2013). Similarly, Simões and coworkers reported that *C. albicans* strains that misincorporate Ser at other codons generally had increased growth rate when exposed to oxidative and osmotic stress (Simoes et al. 2016). Interestingly, the strain that incorporates ~98% of Leu at CUG sites was resistant to SDS and CuSO₄. Considering that SDS limits cell growth in yeasts by disrupting cell membranes and cell walls (Heilmann et al. 2013; Reyna-Beltrán et al. 2019; Schroeder and Ikui 2019), and copper toxicity is used by the innate immune system to kill fungal cells by causing membrane damage (Douglas and Konopka 2019), these findings indicate once again that high levels of Leu incorporation at CUG sites may induce extensive cell wall remodeling. Furthermore, experiments where high CUG mistranslating strains were exposed to human monocyte-derived dendritic cells showed increased in vitro release of the inflammatory ILs IL-1 β , IL-10, and IL-12p70 cytokines, supporting that CUG ambiguity also modulates the immune responses to the fungus. To assess the pathogenic potential of high Leu-CUG translating strains in vivo, C57BL/6 mice were inoculated intragastrically. Although fungal load was higher in mice infected with the wild-type strain, colonization by Leu misincorporating strains was associated with a clear inflammatory pathology in the stomach of infected mice, as revealed by the increased mucin content of goblet cells and the number of infiltrating cells. Moreover, levels of the proinflammatory cytokines TNF α and IL-17A were higher, and the anti-inflammatory IL-10 levels were lower relative to the wild-type strain. These results suggested that a diversified inflammatory response is elicited by dynamic Leu misincorporation at CUGs (Bezerra et al. 2013).

Genome analysis of high-level Leu-CUG translating and wild-type strains showed that increasing levels of CUG ambiguity have a profound impact on the genome (Bezerra et al. 2013). Indeed, high levels of CUG-Leu translation induced loss of heterozygosity (LOH) in a 300-kb region of the chromosome V in strains that incorporate ~80% and ~98% of Leu and for the entire chromosome R in the strain incorporating ~80% of Leu at CUG sites. The observed LOH events affected mainly genes with unknown function, but it also affected genes involved in regulation of biological processes, organelle organization and stress response. Sequence analysis also indicated that the number of SNPs increased proportionally to the level of Leu-CUG translation. The strain with ~98% of Leu incorporation at CUG sites, for example, accumulated higher number of unique SNPs and a higher number of SNPs was found in genes enriched with CUG codons. SNPs accumulated mainly in genes with unknown function, followed by genes associated with filamentous growth, cell adhesion, metabolic processes, and transcription regulation. Thus, the genomic changes were found to be correlated with some of the phenotypes observed in these recombinant cells (Bezerra et al. 2013). In another genomic study, experimental evolution of strains misincorporating Ser at other codons showed a similar great impact on the genome. Genome analysis showed stronger accumulation of LOH events and SNPs in cells with higher codon ambiguity than in control cells (Simoes et al. 2016). Interestingly, a SNP in the *JAB1* gene was found in all highly ambiguous strains; a gene involved in protein degradation by neddylation (Sela, Atir-Lande and Kornitzer 2012; Simoes et al. 2016). Taken together, it shows that codon ambiguity can lead to genome and phenotypic diversification with high adaptation potential, namely in the evasion of immune system and in azole tolerance.

CUG AMBIGUITY ACCELERATES ACQUISITION OF DRUG RESISTANCE

The increasing impact of *Candida* species on human health is mostly due to the existence of a limited number of approved antifungal drugs and the emergence of drug resistance. Currently, there are only five classes of antifungal agents: polyenes (amphotericin B), azoles (fluconazole, itraconazole, posaconazole, voriconazole and isavuconazole), echinocandins (caspofungin, micafungin and anidulafungin), allylamines (terbinafine) and antimetabolites (flucytosine). The scarcity of antifungals is partially explained by the conservation of eukaryotic gene functions between fungal pathogens and the human host (Perlin, Rautemaa-Richardson and Alastruey-Izquierdo 2017; Robbins, Caplan and Cowen 2017; Fisher et al. 2018), leading to a reduced number of viable safe targets. Of all classes, the most used against *Candida* infections belong to the azoles, mainly fluconazole, due to its great efficacy, low toxicity, bioavailability and low cost. However, the extensive use of fluconazole has resulted in increased resistance to azoles, turning therapies increasingly ineffective (Castanheira et al. 2017; Robbins, Caplan and Cowen 2017). Therefore, understanding the molecular mechanisms of drug resistance is of crucial importance for the medical community.

Several studies have identified two main molecular mechanisms of acquisition of azole resistance in *C. albicans*. First, overexpression of the drug target gene *ERG11* (14 α -lanosterol demethylase) reduces the direct impact of the drug (Oliver et al. 2007). Second, increased efflux of the drug from cells by ABC transporters (encoded by *CDR1* and *CDR2*) (Coste et al. 2006) or by the major facilitator superfamily efflux pump (encoded by *MDR1*) (Dunkel et al. 2008) reduces the effective intracellular drug concentration. In both cases, such alterations can result from point mutations in genes encoding those proteins, in transcription factors regulating mRNA expression levels (Coste et al. 2006; Dunkel et al. 2008), or from increased copy number of the relevant genes, via genome rearrangements (Selmecki, Forche and Berman 2006).

The importance of epigenetic pathways in mediating drug resistance in fungi is also well established and involve tuning of a pre-programmed response without permanent genetic changes (Chang et al. 2019). Regulation of translational fidelity may act similarly, as it affords the cell the opportunity of producing an entirely novel set of proteins from the same genetic background. Although most of these proteins will likely be neutral or deleterious in function, a small subset may acquire novel functions entirely. This is the case in bacteria, where Javid et al. showed that sequence variation of antibiotic targets results in resistant phenotypes (Javid et al. 2014). In this study, wild-type mycobacteria and strains with high and low translational fidelity were compared to investigate a direct role of mistranslation in antibiotic tolerance. Low translational fidelity strains had increased antibiotic tolerance both *in vivo* and *in vitro*. When the antibiotic target was purified from low and high mistranslating strains, the mistranslated protein from the high mistranslating strains had increased resistance. This corroborated that protein variation was responsible for the observed phenotype (Javid et al. 2014).

Although mistranslation is associated with antibiotic resistance in bacteria (Balashov and Humayun 2002; Amar, Al Mamun and Humayun 2009; Javid et al. 2014), this link is far less explored in fungal pathogens. As discussed in the previous sections of this review, in yeast and fungi, codon ambiguity leads to genome




Figure 2. Diagram representing the impact of CUG codon ambiguity in *C. albicans* biology. To date, it has been demonstrated three main consequences of Ser/Leu dual translation. (1) **Proteome plasticity:** it is estimated that the 6438 *C. albicans* genes containing CUGs generate 283 billion different proteins (Gomes et al. 2007). The impact of the proteome expansion was shown to generate (2) **Phenotypic diversity**, with high Leu-CUG translating cells show increased filamentation and adhesion to host substrates (Miranda et al. 2007; Miranda et al. 2013). CUG codon ambiguity also induces (3) **Genetic diversity**. *C. albicans* cells with high Leu-CUG levels displayed a higher number of LOH events and accumulation of SNPs into their genomes. Although, *C. albicans* signalling pathways directly affected by CUG codon ambiguity are still largely unexplored, the phenotypes uncovered support the hypothesis that CUG ambiguity modulates immune responses and accelerates ecological adaptation and acquisition of drug resistance (Bezerra et al. 2013; Weil et al. 2017). Figure created with Biorender.com.

and phenotypic diversification and increases adaptation potential, including azole tolerance (Gomes et al. 2007; Miranda et al. 2007; Paredes et al. 2012; Bezerra et al. 2013; Miranda et al. 2013; Kalapis et al. 2015). In *C. albicans*, Weil and colleagues investigated how CUG non-standard translation contributes to the acquisition of antifungal resistance by evolving high-level Leu-CUG translating and wild-type strains in the absence and presence of fluconazole and comparing their resistance trajectories during evolution (Weil et al. 2017). Results showed that increased levels of mistranslation lead to increased tolerance and accelerate the emergence of fluconazole resistance. Genome sequencing, array-based comparative genome analysis, and transcriptional profiling revealed that during evolution in fluconazole, the range of mutational and gene deregulation variations were distinctively different and broader in strains that incorporate Leu at high level. While the expression of genes of the ergosterol biosynthesis pathway, encoding the molecular target of fluconazole, was upregulated in both wild-type and high mistranslating resistant strains, the drug efflux pathway was affected differently (Weil et al. 2017). The wild-type strain acquired a known gain-of-function mutation in the transcription factor *MRR1*, a positive transcriptional regulator of the multidrug efflux pump gene *MDR1* (Morschhauser et al. 2007; Dunkel et al. 2008). On the other hand, the high-level Leu-CUG translating strain acquired a previously described A736V gain-of-function mutation in *TAC1*,

the transcriptional activator of the *CDR1* and *CDR2* genes, which encode the ABC transporters (Coste et al. 2006; Lohberger, Coste and Sanglard 2014). Taken together, and although the stochastic nature of mutations should be stressed here, these results indicated that both strains activated the traditional mechanisms of azole resistance, nonetheless through different drug efflux pathways. Also, the evolution of the hypermistranslating strain, in both the absence and presence of fluconazole, resulted in the fast accumulation of LOH events. A large LOH region in chromosome 5 (Chr5) appeared exclusively in the presence of fluconazole, which is consistent with previous studies of fluconazole-resistant *C. albicans* isolates (Selmecki, Forche and Berman 2006; Ford et al. 2015). Interestingly, analysis of the relative synonymous codon usage in genes affected by LOH events during evolution revealed three key aspects: (i) the number of CUG codons differed between alleles (a and b) of genes that lost heterozygosity; (ii) genes with a higher number of CUG codons in allele a than allele b ($a > b$) underwent a reduction in CUGs by changing them to other codons; (iii) genes with a lower number of CUGs in allele a than allele b ($a < b$) had an increase in the number of CUG codons by changing other codons to CUG. The same pattern was observed in copy number variation analysis of the evolved high-level Leu-CUG translating strain. This resistant strain showed loss of chromosomal repeat regions in Chr2, Chr4 and Chr6, and increased copies of Chr1, Chr4, Chr5 and ChrR. Within these amplified chromosomal regions were a large proportion of genes involved in both translation and drug resistance. Therefore allele-specific gene functions may contribute to rapid adaptation to fluconazole and CUG codon usage and translation seems to have a balancing effect on such genomic events (Bezerra et al. 2013; Kalapis et al. 2015; Weil et al. 2017).

CONCLUSION

The effect of codon ambiguity on phenotypic and genetic diversity of *Candida* spp. adds a new dimension to the study of genome evolution, phenotypic variation, ecological adaptation, drug resistance and virulence in humans. The direct impact of CUG ambiguity in *C. albicans* signalling pathways is still largely unexplored, but the accumulating evidence on phenotypes uncovered support the hypothesis that CUG ambiguity modulates host-pathogen interaction and immune responses and accelerates the evolution of drug resistance (Fig. 2). The implications are wide and should be further explored at the molecular level.

FUNDING

This work was supported by FEDER (Fundo Europeu de Desenvolvimento Regional) funds through the COMPETE 2020, Operational Programme for Competitiveness and Internationalization (POCI), and by Portuguese national funds via Fundação para a Ciência e a Tecnologia, I.P. (FCT) under the projects PBE-POCI-01-0145-FEDER-031 238, GenomePT POCI-01-0145-FEDER-022 184 and PTDC/BIA-MIC/31 849/2017. The iBiMED is supported by FCT funds under UIDB/04 501/2020. CO is supported directly by a FCT grant (SFRH/BD/131637/2017). ARB is supported by national funds (OE), through FCT, I.P., in the scope of the framework contract foreseen in the numbers 4, 5 and 6 of the article 23, of the Decree-Law 57/2016, of August 29, changed by Law 57/2017, of July 19.

Conflicts of interest. None declared.

REFERENCES

- Amar A, Al Mamun M, Humayun MZ. Spontaneous mutagenesis is elevated in protease-defective cells. *Mol Microbiol* 2009;**71**:629–39.
- Balashov S, Humayun MZ. Mistranslation induced by streptomycin provokes a RecABC/RuvABC-dependent mutator phenotype in *Escherichia coli* cells. *J Mol Biol* 2002;**315**:513–27.
- Bezerra AR et al. Reversion of a fungal genetic code alteration links proteome instability with genomic and phenotypic diversification. *Proc Natl Acad Sci* 2013;**110**:11079–84.
- Brown AJ et al. Codon utilisation in the pathogenic yeast, *Candida albicans*. *Nucleic Acids Res* 1991;**19**:4298.
- Butler G et al. Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. *Nature* 2009;**459**:657–62.
- Carlson BA et al. Selenoproteins regulate macrophage invasiveness and extracellular matrix-related gene expression. *BMC Immunol* 2009;**10**:57.
- Castanheira M et al. Monitoring Antifungal Resistance in a Global Collection of Invasive Yeasts and Molds: application of CLSI Epidemiological Cutoff Values and Whole-Genome Sequencing Analysis for Detection of Azole Resistance in *Candida albicans*. *Antimicrob Agents Chemother* 2017;**61**.
- CDC. Antibiotic Resistance Threats in the United States. *Department of Health and Human Services*. Atlanta, GA, United States: U.S. Department of Health and Human Sources, Centers for Disease Control and Prevention. 2019.
- Chang Z et al. Epigenetic mechanisms of drug resistance in fungi. *Fungal Genet Biol* 2019;**132**:103253.
- Coste A et al. A mutation in *Tac1p*, a transcription factor regulating *CDR1* and *CDR2*, is coupled with loss of heterozygosity at chromosome 5 to mediate antifungal resistance in *Candida albicans*. *Genetics* 2006;**172**:2139–56.
- Crick FH. The origin of the genetic code. *J Mol Biol* 1968;**38**:367–79.
- Douglas LM, Konopka JB. Plasma membrane architecture protects *Candida albicans* from killing by copper. *PLoS Genet* 2019;**15**:e1007911.
- Du H et al. *Candida auris*: epidemiology, biology, antifungal resistance, and virulence. *PLoS Pathog* 2020;**16**:e1008921.
- Dunkel N et al. Mutations in the multi-drug resistance regulator *MRR1*, followed by loss of heterozygosity, are the main cause of *MDR1* overexpression in fluconazole-resistant *Candida albicans* strains. *Mol Microbiol* 2008;**69**:827–40.
- Evans CR et al. Errors during Gene Expression: single-Cell Heterogeneity, Stress Resistance, and Microbe-Host Interactions. *mBio* 2018;**9**.
- Feketova Z et al. Ambiguous decoding of the CUG codon alters the functionality of the *Candida albicans* translation initiation factor 4E. *FEMS Yeast Res* 2010;**10**:558–69.
- Fisher MC et al. Worldwide emergence of resistance to antifungal drugs challenges human health and food security. *Science* 2018;**360**:739–42.
- Fitzpatrick David A et al. A fungal phylogeny based on 42 complete genomes derived from supertree and combined gene analysis. *BMC Evol Biol* 2006;**6**.
- Ford CB et al. The evolution of drug resistance in clinical isolates of *Candida albicans*. *Elife* 2015;**4**:e00662.
- Fraga JS et al. Genetic code ambiguity modulates the activity of a *C. albicans* MAP kinase linked to cell wall remodeling. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* 2019;**1867**:654–61.
- Gabalton T, Naranjo-Ortiz MA, Marcet-Houben M. Evolutionary genomics of yeast pathogens in the Saccharomycotina. *FEMS Yeast Res* 2016;**16**.

- Gomes AC et al. A genetic code alteration generates a proteome of high diversity in the human pathogen *Candida albicans*. *Genome Biol* 2007;**8**:R206.
- Guinea J. Global trends in the distribution of *Candida* species causing candidemia. *Clin Microbiol Infect* 2014;**20**:5–10.
- Heilmann CJ et al. Surface stress induces a conserved cell wall stress response in the pathogenic fungus *Candida albicans*. *Eukaryotic Cell* 2013;**12**:254–64.
- Hinnebusch AG, Lorsch JR. The mechanism of eukaryotic translation initiation: new insights and challenges. *Cold Spring Harb Perspect Biol* 2012;**4**.
- Javid B et al. Mycobacterial mistranslation is necessary and sufficient for rifampicin phenotypic resistance. *Proc Natl Acad Sci* 2014;**111**:1132–7.
- Kalapis D et al. Evolution of Robustness to Protein Mistranslation by Accelerated Protein Turnover. *PLoS Biol* 2015;**13**:e1002291.
- Kapur M, Ackerman SL. mRNA Translation Gone Awry: translation Fidelity and Neurological Disease. *Trends Genet* 2018;**34**:218–31.
- Keeling PJ. Genomics: evolution of the Genetic Code. *Curr Biol* 2016;**26**:R851–R3.
- Knight RD, Freeland SJ, Landweber LF. Rewiring the keyboard: evolvability of the genetic code. *Nat Rev Genet* 2001a;**2**:49–58.
- Knight RD, Landweber LF, Yarus M. How mitochondria redefine the code. *J Mol Evol* 2001b;**53**:299–313.
- Kollmar M, Muhlhausen S. Nuclear codon reassignments in the genomics era and mechanisms behind their evolution. *Bioessays* 2017;**39**.
- Koonin EV, Novozhilov AS. Origin and evolution of the genetic code: the universal enigma. *IUBMB Life* 2009;**61**:99–111.
- Krassowski T et al. Evolutionary instability of CUG-Leu in the genetic code of budding yeasts. *Nat Commun* 2018;**9**:1887.
- Lenhard B et al. tRNA recognition and evolution of determinants in seryl-tRNA synthesis. *Nucleic Acids Res* 1999;**27**:721–9.
- Ling J, O'Donoghue P, Soll D. Genetic code flexibility in microorganisms: novel mechanisms and impact on physiology. *Nat Rev Microbiol* 2015;**13**:707–21.
- Lohberger A, Coste AT, Sanglard D. Distinct roles of *Candida albicans* drug resistance transcription factors TAC1, MRR1, and UPC2 in virulence. *Eukaryotic Cell* 2014;**13**:127–42.
- Massey SE et al. Comparative evolutionary genomics unveils the molecular mechanism of reassignment of the CTG codon in *Candida* spp. *Genome Res* 2003;**13**:544–57.
- Miranda I, Silva R, Santos MA. Evolution of the genetic code in yeasts. *Yeast* 2006;**23**:203–13.
- Miranda I et al. A genetic code alteration is a phenotype diversity generator in the human pathogen *Candida albicans*. *PLoS One* 2007;**2**:e996.
- Miranda I et al. *Candida albicans* CUG mistranslation is a mechanism to create cell surface variation. *mBio* 2013;**4**.
- Morschhauser J et al. The transcription factor Mrr1p controls expression of the MDR1 efflux pump and mediates multidrug resistance in *Candida albicans*. *PLoS Pathog* 2007;**3**:e164.
- Moura GR, Paredes JA, Santos MA. Development of the genetic code: insights from a fungal codon reassignment. *FEBS Lett* 2010;**584**:334–41.
- Muhlhausen S et al. A novel nuclear genetic code alteration in yeasts and the evolution of codon reassignment in eukaryotes. *Genome Res* 2016;**26**:945–55.
- Muhlhausen S et al. Endogenous Stochastic Decoding of the CUG Codon by Competing Ser- and Leu-tRNAs in *Ascoidea asiatica*. *Curr Biol* 2018;**28**:2046–57 e5.
- Netzer N et al. Innate immune and chemically triggered oxidative stress modifies translational fidelity. *Nature* 2009;**462**:522–6.
- Oliver BG et al. cis-acting elements within the *Candida albicans* ERG11 promoter mediate the azole response through transcription factor Upc2p. *Eukaryotic Cell* 2007;**6**:2231–39.
- Papon N et al. Emerging and emerged pathogenic *Candida* species: beyond the *Candida albicans* paradigm. *PLoS Pathog* 2013;**9**:e1003550.
- Paredes JA et al. Low level genome mistranslations deregulate the transcriptome and translome and generate proteotoxic stress in yeast. *BMC Biol* 2012;**10**:55.
- Perlin DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. The global problem of antifungal resistance: prevalence, mechanisms, and management. *Lancet Infect Dis* 2017;**17**:e383–e92.
- Pfaller MA et al. Results from the ARTEMIS DISK Global Antifungal Surveillance Study, 1997 to 2007: a 10.5-year analysis of susceptibilities of *Candida* Species to fluconazole and voriconazole as determined by CLSI standardized disk diffusion. *J Clin Microbiol* 2010;**48**:1366–77.
- Reyna-Beltrán E et al. The Cell Wall of *Candida albicans*: a Proteomics View. *Candida Albicans* 2019.
- Ribas de Pouplana L et al. Protein mistranslation: friend or foe? *Trends Biochem Sci* 2014;**39**:355–62.
- Riley R et al. Comparative genomics of biotechnologically important yeasts. *Proc Natl Acad Sci* 2016;**113**:9882–7.
- Robbins N, Caplan T, Cowen LE. Molecular Evolution of Antifungal Drug Resistance. *Annu Rev Microbiol* 2017;**71**:753–75.
- Rocha R et al. Unveiling the structural basis for translational ambiguity tolerance in a human fungal pathogen. *Proc Natl Acad Sci* 2011;**108**:14091–6.
- Santos MA, Perreau VM, Tuite MF. Transfer RNA structural change is a key element in the reassignment of the CUG codon in *Candida albicans*. *EMBO J* 1996;**15**:5060–8.
- Santos MA et al. Driving change: the evolution of alternative genetic codes. *Trends Genet* 2004;**20**:95–102.
- Santos MA et al. The genetic code of the fungal CTG clade. *C R Biol* 2011;**334**:607–11.
- Santos MA et al. The non-standard genetic code of *Candida* spp.: an evolving genetic code or a novel mechanism for adaptation? *Mol Microbiol* 1997;**26**:423–31.
- Santos M et al. Codon misreading tRNAs promote tumor growth in mice. *RNA Biol* 2018;**15**:773–86.
- Schroeder L, Ikui AE. Tryptophan confers resistance to SDS-associated cell membrane stress in *Saccharomyces cerevisiae*. *PLoS One* 2019;**14**:e0199484.
- Schultz DW, Yarus M. On malleability in the genetic code. *J Mol Evol* 1996;**42**:597–601.
- Schwartz MH, Pan T. Function and origin of mistranslation in distinct cellular contexts. *Crit Rev Biochem Mol Biol* 2017;**52**:205–19.
- Sela N, Atir-Lande A, Kornitzer D. Neddylation and CAND1 independently stimulate SCF ubiquitin ligase activity in *Candida albicans*. *Eukaryotic Cell* 2012;**11**:42–52.
- Selmecki A, Forche A, Berman J. Aneuploidy and isochromosome formation in drug-resistant *Candida albicans*. *Science* 2006;**313**:367–70.
- Simoes J et al. The Fungus *Candida albicans* Tolerates Ambiguity at Multiple Codons. *Front Microbiol* 2016;**7**:401.
- Sugiyama H et al. In vivo evidence for non-universal usage of the codon CUG in *Candida maltosa*. *Yeast* 1995;**11**:43–52.
- Suzuki T, Ueda T, Watanabe K. The 'polysemous' codon—a codon with multiple amino acid assignment caused

- by dual specificity of tRNA identity. *EMBO J* 1997;**16**:1122–34.
- Tuite MF, Santos MA. Codon reassignment in *Candida* species: an evolutionary conundrum. *Biochimie* 1996;**78**:993–9.
- Weil T et al. Adaptive Mistranslation Accelerates the Evolution of Fluconazole Resistance and Induces Major Genomic and Gene Expression Alterations in *Candida albicans*. *mSphere* 2017;**2**.
- Zhou XL et al. Aminoacylation and translational quality control strategy employed by leucyl-tRNA synthetase from a human pathogen with genetic code ambiguity. *Nucleic Acids Res* 2013;**41**:9825–38.