

# Journal Pre-proof

Test-retest reliability, agreement and construct validity of the International Physical Activity Questionnaire short-form (IPAQ-sf) in people with COPD

Sofia Flora, Alda Marques, Nádia Hipólito, Nuno Morais, Cândida G. Silva, Filipa Januário, Fátima Rodrigues, Bruno P. Carreira, J. Cruz



PII: S0954-6111(22)00352-3

DOI: <https://doi.org/10.1016/j.rmed.2022.107087>

Reference: YRMED 107087

To appear in: *Respiratory Medicine*

Received Date: 23 August 2022

Revised Date: 7 November 2022

Accepted Date: 5 December 2022

Please cite this article as: Flora S, Marques A, Hipólito Ná, Morais N, Silva CâG, Januário F, Rodrigues Fá, Carreira BP, Cruz J, Test-retest reliability, agreement and construct validity of the International Physical Activity Questionnaire short-form (IPAQ-sf) in people with COPD, *Respiratory Medicine* (2023), doi: <https://doi.org/10.1016/j.rmed.2022.107087>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 The Author(s). Published by Elsevier Ltd.

**Author contributions**

**Sofia Flora:** Methodology, Formal analysis, Investigation, Writing - Original Draft, Visualization. **Alda Marques:** Conceptualization, Writing - Review & Editing, Supervision. **Nádia Hipólito:** Investigation, Writing - Review & Editing. **Nuno Morais:** Validation, Formal analysis, Resources, Writing - Review & Editing, Supervision. **Cândida G. Silva:** Validation, Formal analysis, Resources, Writing - Review & Editing, Supervision. **Filipa Januário:** Writing - Review & Editing, Supervision. **Fátima Rodrigues:** Writing - Review & Editing, Supervision. **Bruno P. Carreira:** Writing - Review & Editing, Supervision. **Joana Cruz:** Conceptualization, Methodology, Validation, Formal analysis, Resources, Writing - Review & Editing, Visualization, Supervision, Project administration

1 **Test-retest reliability, agreement and construct validity of the International Physical**  
2 **Activity Questionnaire short-form (IPAQ-sf) in people with COPD**

3 Sofia Flora<sup>1</sup>, Alda Marques<sup>2</sup>, Nádia Hipólito<sup>3</sup>, Nuno Morais<sup>4</sup>, Cândida G. Silva<sup>5</sup>, Filipa Januário<sup>6</sup>,  
4 Fátima Rodrigues<sup>7</sup>, Bruno P. Carreira<sup>8</sup>, J. Cruz<sup>9</sup>

5 <sup>1</sup>Center for Innovative Care and Health Technology (ciTechCare), Polytechnic of Leiria - Leiria (Portugal)

6 <sup>2</sup>Lab 3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences (ESSUA) and Institute of  
7 Biomedicine (iBiMED), University of Aveiro - Aveiro (Portugal)

8 <sup>3</sup>Center for Innovative Care and Health Technology (ciTechCare), Polytechnic of Leiria - Leiria (Portugal)

9 <sup>4</sup>Center for Innovative Care and Health Technology (ciTechCare), Polytechnic Institute of Leiria; School of Health Sciences,  
10 Polytechnic Institute of Leiria; Centre for Rapid and Sustainable Product Development (CDRSP), Polytechnic Institute of  
11 Leiria, Leiria, Portugal

12 <sup>5</sup>Center for Innovative Care and Health Technology (ciTechCare), Polytechnic Institute of Leiria; School of Health Sciences,  
13 Polytechnic Institute of Leiria; Coimbra Chemistry Centre, Department of Chemistry, University of Coimbra, Coimbra,  
14 Portugal

15 <sup>6</sup>Physical Medicine and Rehabilitation Department, Leiria Hospital Center, Leiria, Portugal

16 <sup>7</sup>Institute of Health Environmental, Faculty of Medicine, University of Lisbon; Pulmonary Rehabilitation Unit, Hospital Pulido  
17 Valente, University Hospital Center North Lisbon – Lisboa (Portugal)

18 <sup>8</sup>Center for Innovative Care and Health Technology (ciTechCare), Polytechnic of Leiria; School of Health Sciences,  
19 Polytechnic of Leiria - Leiria; Unidade de Saúde Familiar Pedro e Inês, ACES Oeste Norte, Alcobaca, Portugal

20 <sup>9</sup>Center for Innovative Care and Health Technology (ciTechCare), Polytechnic of Leiria; School of Health Sciences,  
21 Polytechnic of Leiria - Leiria (Portugal)

22

23 **Corresponding author:**

24 Joana Cruz, Center for Innovative Care and Health Technology (ciTechCare), School of Health Sciences  
25 (ESSLei), Polytechnic of Leiria, Campus 2, Morro do Lena – Alto do Vieiro, Apartado 4163, 2411-901 Leiria –  
26 PORTUGAL, [Joana.cruz@ipleiria.pt](mailto:Joana.cruz@ipleiria.pt)

27

28 **Declarations of interest:** none.

29

30 **Funding:** This work was funded by FEDER - Fundo Europeu de Desenvolvimento Regional by  
31 COMPETE 2020 – Programa Operacional Competitividade e Internacionalização (POCI) and national  
32 funds by Fundação para a Ciência e a Tecnologia (FCT), under the “OnTRACK project - Time to  
33 Rethink Activity Knowledge: a personalized mHealth coaching platform to tackle physical inactivity in  
34 COPD” (POCI-01-0145-FEDER-028446, PTDC/SAU-SER/28446/2017); national funds by FCT

35 (UIDB/05704/2020, UIDP/05704/2020, UIDB/04501/2020). SF and NH were financially supported by  
36 PhD fellowships DFA/BD/6954/2020 and 2021.05188.BD, respectively, funded by FCT/MCTES, FSE,  
37 Por\_Centro and UE.

### 38 **ABSTRACT (250 words)**

39 **Introduction:** This study assessed the test-retest reliability/agreement and construct validity of the  
40 International Physical Activity Questionnaire short-form (IPAQ-sf) in patients with chronic obstructive  
41 pulmonary disease (COPD). It also explored differences in its validity according to age, sex and GOLD  
42 airflow obstruction levels.

43 **Methods:** 62 participants (68±8 years, 53 males, FEV<sub>1</sub> 51±23%pred) completed the Portuguese  
44 IPAQ-sf, wore an accelerometer for 7 days and completed a second IPAQ-sf. Test-retest  
45 reliability/agreement was assessed with Intraclass Correlation Coefficient (ICC<sub>2,1</sub>), 95% Limits of  
46 Agreement (LoA), standard error of measurement (SEM) and minimal detectable change (MDC<sub>95</sub>) for  
47 continuous variables, and percentage of agreement (%agreement) for categories (“active”/“inactive”).  
48 Validity was assessed with 95% LoA and Spearman’s correlations ( $\rho$ ) between IPAQ-sf 2 (METs-  
49 min/week, time in vigorous [VPA], moderate PA [MPA] and walking) and accelerometry (time in  
50 MVPA, VPA, MPA and step counts) for continuous variables; %agreement, Cohen’s kappa, and  
51 sensitivity specificity and +/- predictive values for categories. Correlations were also performed for  
52 age, sex and GOLD airflow obstruction grades.

53 **Results:** Reliability was good (ICC<sub>2,1</sub>=0.707) with wide LoA (-6446—6409 METs-min/week). SEM and  
54 MDC<sub>95</sub> were 1840 and 4971 METs-min/week, respectively. %agreement between the two IPAQ-sf  
55 was 84% (kappa=0.660). Positive, moderate and significant correlations were found between IPAQ-  
56 sf and accelerometry ( $0.396 \leq \rho \leq 0.527$ ,  $p < 0.001$ ), except for VPA ( $p > 0.05$ ). The strongest correlations  
57 were found in age (<65 years) and male ( $0.466 \leq \rho \leq 0.653$ ,  $p < 0.05$ ). %agreement between tools was  
58 65% (kappa=0.313), with high sensitivity (0.830) but low specificity (0.500).

59 **Conclusions:** The IPAQ-sf seems valid to be used in COPD but caution on its widespread use is  
60 recommended as its accuracy may be limited.

61 **Keywords:** Accelerometer, chronic obstructive pulmonary disease, physical activity, psychometric  
62 properties, validation study.

### 63 **Abbreviations' list**

64 ACSM - American College of Sports Medicine

65 BMI - Body mass index

66 CCI - Charlson Comorbidity Index

67 CI - Confidence Intervals

68 ciTechCare - Centre for Innovative Care and Health Technology

69 COPD – Chronic Obstructive Pulmonary Disease

70 COSMIN - Consensus-based Standards for the selection of health Measurement INstruments

71 FEV<sub>1</sub> - Forced Expiratory Volume in first second

72 GOLD - Global Initiative for Chronic Obstructive Lung Disease

73 ICC - Intraclass Correlation Coefficient

74 IPAQ-sf - International Physical Activity Questionnaire short-form

75 LoA - Limits of agreement

76 MDC<sub>95</sub> - Minimal detectable change

77 METs - Metabolic equivalents

78 mMRC - modified Medical Research Council dyspnoea scale

79 MPA – moderate physical activity

- 80 MVPA - Moderate and vigorous physical activity
- 81 NPV - Negative predictive value
- 82 PA – Physical Activity
- 83 PAR - Stanford Seven-Day Physical Activity Recall
- 84 PPV – Positive predictive value
- 85 SD - Standard deviation
- 86 SEM - Standard error of measurement
- 87 SPSS - Statistical Package for the Social Sciences
- 88 VPA – vigorous physical activity
- 89 WHO - World Health Organisation

Journal Pre-proof

## 90 Introduction

91 People with chronic obstructive pulmonary disease (COPD) are markedly inactive in daily life [1] which  
92 contributes to a worsening of lung function, health status [2], increased risk of acute exacerbations,  
93 hospitalizations and mortality in this population [3]. Physical activity (PA) is a modifiable factor with  
94 potential to improve COPD prognosis, therefore the latest Global Initiative for Chronic Obstructive  
95 Lung Disease (GOLD) guidelines [4] have underlined the importance of assessing and promoting  
96 regular PA as part of COPD management.

97 The International Physical Activity Questionnaire short-form (IPAQ-sf) is one of the most widely used  
98 self-reported questionnaires to assess PA. Although good measurement properties were reported in  
99 the healthy population of the original study [5], measurement properties are population-specific. In  
100 fact, a systematic review has shown that studies assessing the validity of this instrument presented  
101 conflicting results, suggesting that evidence to support its use as an indicator of PA is weak [6].  
102 Moreover, poor validity results have also been found in populations with chronic conditions, such as  
103 in rheumatoid arthritis [7], fibromyalgia [8] or systemic lupus erythematosus [9]. In COPD, the IPAQ-  
104 sf has been used in several studies to estimate patients' PA levels [10-12]. This study showed strong,  
105 positive and significant correlations between the IPAQ-sf METs-min/week and moderate and vigorous  
106 physical activity (MVPA) measured with an accelerometer ( $r=0.729$ ,  $p=0.017$ ), but low percentage of  
107 agreement (% agreement) in identifying "physically active" and "physically inactive" patients (%  
108 agreement=20%,  $\kappa=-0.538$ ), and poor to moderate test-retest reliability (Intraclass Correlation  
109 Coefficient [ICC]=0.439, 95% Confidence Intervals [95%CI] -0.267 — 0.838). The small sample size  
110 of this study hinders the generalisability of the findings. Further research is therefore needed to assess  
111 the measurement properties of the IPAQ-sf in COPD. Furthermore, previous studies have shown  
112 differences in PA levels among GOLD airflow obstruction levels [13], and an influence of age and sex  
113 in patients' PA behaviour [14], hence it may be important to explore the performance of the IPAQ-sf  
114 in these specific subgroups.

115 This study aimed to assess the test-retest reliability/agreement and construct validity of the IPAQ-sf  
116 in people with COPD. A secondary aim of this study was to explore potential differences in the validity  
117 of the tool among groups of age, sex, and COPD levels of airflow obstruction.

## 118 **Methods**

### 119 **Study design**

120 This was a cross-sectional study which was part of a larger study (ref. POCI-01-0145-FEDER-  
121 028446; PTDC/SAU-SER/28446/2017). Construct validity of the IPAQ-sf was assessed using  
122 accelerometer-based data. Test-retest reliability/agreement was calculated using the IPAQ-sf results  
123 obtained in two different occasions separated by 7 days, corresponding to the time participants used  
124 the accelerometer.

### 125 **Ethical considerations**

126 Ethical approval was obtained prior to study commencement from the Health Units participating in  
127 this study. Participants received verbal and written information about the study and provided written  
128 informed consent before data collection.

### 129 **Sample size**

130 Sample size was defined according to the COnsensus-based Standards for the selection of health  
131 Measurement INstruments (COSMIN) guidelines [15, 16], which recommend that a minimum of 50  
132 individuals should be recruited to ensure the quality of studies assessing the measurement properties  
133 of instruments.

### 134 **Participants**

135 Patients with COPD were identified by physicians of the Leiria Hospital Centre, Baixo Vouga Hospital  
136 Centre, University Hospital Centre North Lisbon and a primary care centre (USF Santiago Marrazes),  
137 who ensured the fulfilment of the eligibility criteria. Patients included in the study had to be: 18 years  
138 old or more; diagnosed with COPD according to the GOLD criteria [4]; clinically stable in the previous



139 month (i.e., no hospital admissions or acute exacerbations); able to understand Portuguese and to  
140 provide informed consent. Exclusion criteria consisted of the presence of severe neurologic (e.g.,  
141 Parkinson, stroke), musculoskeletal (e.g., severe osteoarthritis) or psychiatric disorders (e.g.,  
142 schizophrenia), unstable cardiovascular disease, or other health condition/impairment (e.g., severe  
143 visual or hearing impairment) that could preclude patients from understanding the study and/or  
144 participating in data collection. Data were collected at the Centre for Innovative Care and Health  
145 Technology (ciTechCare) of the Polytechnic of Leiria, at the Respiratory Research and Rehabilitation  
146 Laboratory – School of Health Sciences, University of Aveiro (Lab3R-ESSUA), or at the health units,  
147 depending on patients' and services' availability.

#### 148 **Data collection**

149 Participants completed a structured questionnaire with sociodemographic (age, sex, education level  
150 and work status) and general clinical information such as smoking status (never, current or former  
151 smokers), dyspnoea perception (modified Medical Research Council dyspnoea scale [mMRC] [17])  
152 and presence of comorbidities (Charlson Comorbidity Index [CCI] [18]) to characterise the sample.  
153 Comorbidities were classified as mild (CCI scores of 1–2), moderate (CCI scores of 3–4) or severe  
154 (CCI scores  $\geq 5$ ) [18]. Height and weight were collected to calculate the body mass index (BMI). Lung  
155 function was assessed according to standardised guidelines [19] with a portable spirometer  
156 (MicroLoop, CareFusion, Kent, UK) to characterise airflow obstruction limitation [4]: GOLD grades  
157 1–4 (considering patients' Forced Expiratory Volume in first second percentage predicted [ $FEV_1$  %  
158 predicted]: GOLD 1 –  $FEV_1 \geq 80\%$ ; GOLD 2 –  $50 \leq FEV_1 \leq 79\%$ ; GOLD 3 –  $30 \leq FEV_1 \leq 49\%$  and  
159 GOLD 4 –  $FEV_1 < 30\%$ ). All patients were advised to take their usual medication before data collection.  
160 Then, participants completed the IPAQ-sf (IPAQ-sf 1) and received an accelerometer (ActiGraph  
161 GT3X+, Pensacola, FL) to use for 7 days [20]. Patients were instructed to wear the accelerometer at  
162 the waist, on the dominant side, during waking hours, except for bathing or swimming. A second  
163 appointment was scheduled 8 days after the first appointment to retrieve the accelerometers and

164 complete the IPAQ-sf once more (IPAQ-sf 2), for further assessment of test-retest reliability and  
165 agreement of the tool.

## 166 **Measures**

### 167 International Physical Activity Questionnaire short-form (IPAQ-sf)

168 The IPAQ-sf is composed of 7 questions, simple to administer in clinical practice, and provides  
169 information on the number of days/week and average time/day spent walking, in moderate- and  
170 vigorous-intensity activities and sitting, based on the previous 7 days, to further calculate energy  
171 expenditure in metabolic equivalents (METs) [5]. The continuous score of the IPAQ-sf can be  
172 calculated as “MET level × minutes of activity per day × days per week” and is expressed in METs-  
173 min/week. It can be calculated for walking (3.3 METs), MPA (4 METs) and VPA (8 METs). The  
174 categorical score of the IPAQ-sf classifies a patients’ PA level as “low”, “moderate” or “high” [21].  
175 These classifications can be then translated to “physically active” (corresponding to “moderate” or  
176 “high” PA levels) and “physically inactive” (which corresponds to “low” PA level) (Table 1). The  
177 Portuguese version of IPAQ-sf was used in this study [5] and it takes about 10 minutes to complete.  
178 The questionnaire is free of charge and can be found in the IPAQ website  
179 (<https://sites.google.com/site/theipaq/home>), along with a detailed scoring information.

### 180 Accelerometry

181 Accelerometry was used as a criterion measure to validate the IPAQ-sf, similarly to other validation  
182 studies [9, 22-24]. In this study, the triaxial accelerometer ActiGraph GT3X+ was used, which has  
183 been validated in the COPD population [20, 25]. The device collects and stores PA data which can  
184 be downloaded and converted into time-stamped PA counts and step counts using specific software  
185 (ActiLife 6, version 6.13.3, Pensacola, FL). A valid day was defined as a minimum of 8 hours of  
186 wearing time [26]. Patients who had less than 5 days of valid data from the 7-day wear interval were  
187 excluded, since 4 days are the minimum number of days needed for an accurate assessment of  
188 patients’ PA using accelerometers [26], and at least 5 days are required to assess whether patients

189 are physically active or not (considering the moderate-intensity PA – Table 1) [27]. Accelerometer-  
 190 based data were then downloaded and analysed using the algorithms of Freedson (1998) [28] with  
 191 60-s epoch, incorporated in the Actilife software: daily time (in min) spent in light-intensity PA (100—  
 192 1951 counts-per-minute [CPM]), MPA (1952—5724 CPM), VPA ( $\geq 5725$  CPM), and a combination of  
 193 both (MVPA) [28]. Data were retrieved in min/week to facilitate the comparison with the results from  
 194 IPAQ-sf. The number of steps per day and per week was also collected. Participants were classified  
 195 as “physically active” or “physically inactive” using two approaches, an intensity-based approach and  
 196 a step-based approach, according to the American College of Sports Medicine (ACSM) guidelines  
 197 [27] and World Health Organisation (WHO) [29] (Table 1).

198 **Table 1 - Categories of “physically active” and “physically inactive” obtained with the IPAQ-sf**  
 199 **and accelerometer-based data.**

Category	Physically active	Physically inactive
IPAQ-sf	Correspond to “high” and “moderate” scores of the IPAQ-sf:  <p style="text-align: center;"><b>“High PA level”</b></p> a) vigorous-intensity PA on $\geq 3$ days achieving $\geq 1500$ MET-min/week  <p style="text-align: center;"><u>OR</u></p> b) 7 days of any combination of walking, moderate- or vigorous-intensity PA achieving $\geq 3000$ MET-min/week  <p style="text-align: center;"><b>“Moderate PA level”</b></p> a) $\geq 3$ days of vigorous-intensity PA of $\geq 20$ min/day  <p style="text-align: center;"><u>OR</u></p> b) $\geq 5$ days of moderate-intensity PA and/or walking of $\geq 30$ min/day  <p style="text-align: center;"><u>OR</u></p>	Correspond to “low” score of the IPAQ-sf:  <p style="text-align: center;"><b>“Low PA level”</b></p> a) No PA is reported  <p style="text-align: center;"><u>OR</u></p> b) Some PA is reported but not enough to meet categories “high” or “moderate”

	c) $\geq 5$ days of any combination of walking, moderate- or vigorous-intensity PA achieving $\geq 600$ MET-min/week	
<b>Accelerometer (intensity-based approach) [27]</b>	<p>a) <math>\geq 20</math> min/day of vigorous-intensity PA on <math>\geq 3</math> days, to reach a total of at least 75 min/week</p> <p style="text-align: center;"><u>OR</u></p> <p>b) <math>\geq 30</math> min/day of moderate-intensity PA on <math>\geq 5</math> days, to reach a total of at least 150 min/week</p> <p style="text-align: center;"><u>OR</u></p> <p>c) a combination of both</p>	<p>a) No PA is reported</p> <p style="text-align: center;"><u>OR</u></p> <p>b) Some PA is reported but not enough to meet the guidelines</p>
<b>Accelerometer (step-based approach) [27]</b>	a) $\geq 7000$ steps/day	a) Not achieving the minimum of 7000 steps/day

200 **Legend:** IPAQ-sf, International Physical Activity Questionnaire - short form; METs, metabolic  
 201 equivalent; PA, physical activity.

## 202 **Data analysis**

203 Descriptive statistics were used to characterise the sample regarding age, sex, FEV<sub>1</sub> % predicted,  
 204 BMI, education level, work status, smoking status, GOLD airflow obstruction limitation (1–4),  
 205 dyspnoea (mMRC), comorbidities (CCI) and PA results (IPAQ-sf and accelerometer-based data).

## 206 Reliability and Agreement

207 Test-retest reliability of the IPAQ-sf was assessed using: 1) continuous values of IPAQ-sf 1 and IPAQ-  
 208 sf 2 (METs-min/week); and 2) categories of IPAQ-sf 1 and 2 (i.e., “low PA”, “moderate PA” and “high  
 209 PA”; and “physically active” vs. “physically inactive”). According to the guidelines [16, 30, 31], the  
 210 following analyses were conducted:

211 1) For continuous variables:

- 212 a. Reliability was assessed using  $ICC_{2,1}$  and its 95% CI [32]. An ICC of at least 0.70 was  
 213 considered as a minimum standard for good reliability [33].
- 214 b. Agreement was calculated using the standard error of measurement ( $SEM =$   
 215  $\frac{SD_{differences}}{\sqrt{2}}$ ), minimal detectable change at the 95% confidence level ( $MDC_{95} =$   
 216  $SEM \times \sqrt{2} \times 1.96$ ), and the Bland and Altman 95% limits of agreement (LoA) [34].
- 217 2) For categorical variables:
- 218 a. Percentage of agreement was defined as the total number of participants assigned to  
 219 the same category (either “physically active” or “physically inactive”) by both measures,  
 220 divided by the total number of participants.
- 221 b. Cohen’s weighted kappa coefficient and its 95% CI were used for ordinal variables  
 222 (“low PA”, “moderate PA” and “high PA”) and Cohen’s kappa for nominal variables  
 223 (“physically inactive” and “physically active”). Results were interpreted as follows [35]:  
 224 slight ( $\leq 0.20$ ), fair (0.21—0.40), moderate (0.41—0.60), substantial (0.61—0.80) and  
 225 almost perfect (0.81—1.00). An acceptable value of kappa was considered as  $\geq 0.70$   
 226 [33].

### 227 Construct Validity

228 The IPAQ-sf 2 and accelerometer-based data were used to assess the construct validity of the PA  
 229 assessment tool, since they referred to the same period. Criterion validity was not possible to assess  
 230 as there is still no gold standard for the assessment of daily PA [16, 36].

231 The variables used from IPAQ-sf were the following (all in min/week): METs-min/week, time spent in  
 232 VPA (i.e., product of IPAQ-sf questions 1 and 2), time spent in MPA (i.e., product of IPAQ-sf questions  
 233 3 and 4) and in time spent in walking (i.e., product of IPAQ-sf questions 5 and 6). From accelerometry,  
 234 the following variables were used: time spent in VPA, MPA and MVPA (combination between VPA  
 235 and MPA) (in min/week), and step counts per week. The question regarding the time spent sitting  
 236 (Q7) is not included as part of the continuous score and was not addressed in the present study.

237 Normality of data distribution was assessed using the Kolmogorov-Smirnov test for each variable.

238 The following analyses were conducted, according to the guidelines [30]:

239 1) For continuous variables:

240 a. Spearman's rank-order correlations ( $\rho$ ) or Pearson's correlation coefficient ( $r$ )  
241 (according to the [non-]normality of data distribution) were used in the total sample and  
242 in the following subgroups: 1) age ( $< 65$  and  $\geq 65$  years old); 2) sex (male and female);  
243 and 3) GOLD airflow obstruction levels (GOLD 1, 2, 3 and 4). Construct validity is often  
244 considered good if correlations are positive, significant and  $\geq 0.50$  [33]. Strength of the  
245 correlations were based on criteria from Evans [37]: very weak (0.00—0.19), weak  
246 (0.20—0.39), moderate (0.40—0.59), strong (0.60—0.79) and very strong (0.80—1.0).  
247 b. Bland and Altman's 95% LoA were used to compare the two measurement methods  
248 on variables that have used same units: weekly time spent on vigorous activity (VPA),  
249 moderate activity (MPA) and walking.

250 2) For categorical variables:

251 a. The ability of the IPAQ-sf for classifying "physically active" and "physically inactive"  
252 patients was evaluated against the accelerometer-based data, using the cut-off points  
253 previously described (Table 1). Percentage of agreement and Cohen's kappa  
254 coefficient were used.

255 b. Sensitivity (i.e., those who were correctly classified as "physically active" by the IPAQ-  
256 sf using the accelerometer-based data) and specificity (i.e., those who were correctly  
257 classified as "physically inactive" by the IPAQ-sf using the same criteria) were also  
258 calculated, including the 95% CI. The 95% CI were calculated for sensitivity and  
259 specificity using the following formula  $= p \pm 1.96 \sqrt{\frac{p(1-p)}{n}}$ , where "p" is the relevant  
260 proportion (i.e., sensitivity or specificity) and "n" is the total sample [32].

261 c. Positive and negative predictive values (PPV and NPV, respectively) were calculated  
262 and refer to the proportion of "physically active" (PPV) and "physically inactive" (NPV)

263 participants classified by the IPAQ-sf who were “truly physically active” and “truly  
264 physically inactive”, respectively, having the accelerometer as the reference standard.

265 All data were analysed using SPSS version 24 (IBM Corp., Armonk, USA) and GraphPad Prism  
266 Version 8.0.1. (263). Statistical significance was set at  $p < 0.05$ .

## 267 Results

### 268 Participants

269 A total of 103 patients with COPD were identified. From these, 18 refused to participate, 2 withdrew  
270 from participating and 1 died. Additionally, 2 reported having had an exacerbation in the previous  
271 days and 7 were not available to participate at the moment of data collection. When considering the  
272 IPAQ scoring guidelines [38] eleven participants were excluded from the analysis due to: presenting  
273 a very high score, i.e., > 16 hours at walking, moderate and vigorous PA (n=3); being significant  
274 outliers, i.e.,  $\geq 16$ h of different intensities PA (n=3) and missing data (n=5; 2 in the IPAQ-sf and 3 in  
275 accelerometry). The final sample was composed of 62 participants.

276 Participants had a mean ( $\pm$  standard deviation) age of  $68 \pm 8$  years old and 53 (86%) participants were  
277 male. They were slightly overweight (BMI= $27 \pm 5$  kg/m<sup>2</sup>) and presented a FEV<sub>1</sub> of  $51 \pm 23\%$  predicted.  
278 Their detailed sociodemographic and clinical characteristics are presented in Table 2. Most  
279 participants were in GOLD 2 (n=25, 40%) and GOLD 3 (n=20, 32%) of airflow obstruction. All  
280 participants reported comorbidities, the most common being arterial hypertension (n=26, 43%),  
281 dyslipidemia (n=18, 30%) and mental health problems, such as anxiety and depression (n=23, 43%).

### 282 Table 2 - Participants' sociodemographic and clinical characteristics (n=62).

Participants' characteristics (n=62)	
Age (years), mean (SD)	68 (8)
Sex (male), n (%)	53 (86%)
FEV <sub>1</sub> % predicted, mean (SD)	51 (23)
BMI (kg/m <sup>2</sup> ), mean (SD)	27 (5)

Education Level, n (%)	
No qualifications	2 (3%)
1 <sup>st</sup> cycle (years 1-4)	26 (42%)
2 <sup>nd</sup> cycle (years 5-6)	7 (11%)
3 <sup>rd</sup> cycle (years 7-9)	7 (11%)
High school (years 10-12)	14 (23%)
University	6 (10%)
Work status, n (%)	
Retired	50 (82%)
Full/part-time employment	5 (8%)
Unemployed (health-related reason)	5 (8%)
Smoking status, n (%) <sup>1</sup>	
Never	8 (14%)
Current smokers	11 (19%)
Former smokers	39 (68%)
GOLD airflow obstruction levels, n (%)	
GOLD 1	5 (8%)
GOLD 2	25 (40%)
GOLD 3	20 (32%)
GOLD 4	12 (19%)
mMRC, median [Q1; Q3]	2 [1; 2]
CCI, n (%)	
Mild	8 (13%)
Moderate	43 (71%)
Severe	10 (16%)

283 **Legend:** BMI, body mass index; CCI, Charlson comorbidity index; FEV<sub>1</sub>, forced expiratory volume in  
 284 first second; FVC, forced vital capacity; mMRC, Modified Medical Research Council; SD, standard  
 285 deviation. Q, quartile. <sup>1</sup>Missing cases: 4.



286 **Physical activity levels**

287 Physical activity data are presented in Table 3. None of the variables from the IPAQ-sf or the  
 288 accelerometer followed a normal distribution, hence data are presented as median (quartile  
 289 [Q1; Q3]). More than 50% of the sample did not meet the international PA recommendations  
 290 (median of MPA=85 min/week), which is lower than the 150 min/week recommended [27].  
 291 From 62 participants, 56 used the accelerometer for 7 days (4 used for 6 days and 2 used for  
 292 5 days).

293 **Table 3 – Data from the IPAQ-sf, IPAQ-sf 2 (retest) and accelerometer-based data (n=62).**


<b>IPAQ-sf 1 (min/week)</b>	
Total energy expenditure (METs-min/week)	1193 [220; 2996]
Time in moderate PA	60 [0; 285]
Time in vigorous PA	0 [0; 0]
Time in walking	130 [0; 300]
<b>IPAQ-sf 2, median (min/week)</b>	
Total energy expenditure (METs-min/week)	1550 [309; 3254]
Time in moderate PA	73 [0; 304]
Time in vigorous PA	0 [0; 180]
Time in walking	140 [28; 360]
<b>Accelerometry (min/week)</b>	
Time in moderate PA	85 [46; 248]
Time in vigorous PA	1 [1; 2]
Total time in MVPA	87 [47; 248]
Steps (per day)	3504 [2313; 5766]

294 **Legend:** IPAQ-sf, International Physical Activity Questionnaire-short form; METs, metabolic  
 295 equivalent; Min, minutes; MVPA, moderate and vigorous physical activity; PA, physical activity; SD,  
 296 standard deviation. The results are presented in median (the percentile 25 [Q1]; percentile 75 [Q3]).

297 **Test-retest reliability and agreement of IPAQ-sf**298 IPAQ-sf continuous scores

299 Test-retest reliability and agreement of the IPAQ-sf were first analysed using the continuous scores  
 300 from IPAQ-sf 1 and 2 (in METs-min/week). The ICC was 0.707 (95% CI 0.515—0.823), and the values  
 301 of the SEM and MDC<sub>95</sub> were 1840 METs-min/week and 4971 METs-min/week, respectively.

302 Figure 1 presents a Bland and Altman plot with the 95% LoA between the IPAQ-sf 1 and 2 (METs-  
 303 min/week). A bias (i.e., mean differences between IPAQ-sf 1 and 2) of -18.6 METs-min/week  
 304 (standard deviation of bias= 3279 METs-min/week) was observed, with wide 95% LoA ranging from  
 305 -6446 to 6409 METs-min/week, and no evidence of consistent bias was found.



306

307 **Figure 1 - Bland and Altman plots between IPAQ-sf 1 e 2 (total METs-min/week) in patients with**  
 308 **chronic obstructive pulmonary disease (n=62).**

309 IPAQ-sf categories

310 The percentage of agreement among IPAQ-sf categories (“low PA”, “moderate PA” and “high PA”)  
 311 obtained from IPAQ-sf 1 and 2 was 66% and the weighted Cohen’s kappa was 0.496 (95% CI 0.329—  
 312 0.663), as shown in the appendix A. When considering the categories “physically inactive” (i.e., low  
 313 PA) and “physically active” (i.e., moderate to high PA), the agreement was 84% and the Cohen’s  
 314 kappa was 0.660 (95% CI 0.493—0.827), as shown in Table 4.

315 **Table 4 - Percentage of agreement and weighted Cohen's kappa among IPAQ-sf categories**  
 316 **(“physically inactive” and “physically active”) (n=62).**

		IPAQ-sf 2		% Agreement	Kappa (95% CI)
		Physically Inactive	Physically Active		
IPAQ-sf 1	Physically Inactive	19	6	84%	0.660 (0.493 to 0.827)
	Physically Active	4	33		
	Physically Inactive				
	Physically Active				

317 **Legend:** CI, confidence intervals; IPAQ-sf, International Physical Activity Questionnaire-short form.

### 318 **Validity of the IPAQ-sf**

#### 319 IPAQ-sf and accelerometry - continuous variables

320 Correlations between measurement methods were positive, moderate and significant in all PA  
 321 variables ( $0.396 \leq \rho \leq 0.527$ ,  $p < 0.001$ ), except for VPA ( $\rho = 0.006$ ,  $p > 0.05$ ) (appendix B). Overall, the  
 322 IPAQ-sf overestimated the weekly time spent in activity (mean differences between methods [95%  
 323 LoA] for VPA = 45 min/week [135 – 224], MPA = 18 min/week [-480 – 515] and Walking = 35  
 324 min/week, [-491 – 561] and this was more evident the longer the patients report being active,  
 325 particularly in VPA (Figures 2, 3 and 4).

#### 326 *Subgroup analyses*

327 Significant, positive and moderate correlations were found between the IPAQ-sf and accelerometry  
 328 in patients independently of the age group and in male patients (except for VPA in both groups,  
 329  $p > 0.05$ ). The highest values were obtained in patients with <65 years ( $0.467 \leq \rho \leq 0.651$ ,  $p < 0.05$ ) and  
 330 in male patients ( $0.466 \leq \rho \leq 0.653$ ,  $p < 0.001$ ). Correlations were negative and non-significant for female  
 331 patients ( $-0.594 \leq \rho \leq -0.159$ ,  $p > 0.05$ ). In GOLD grades, significant correlations were only found for:  
 332 IPAQ-sf total score and total duration in MVPA (accelerometry) (GOLD 2 and 4), time in MPA (GOLD

333 2), time in walking and in MPA (GOLD 4), and time in walking and steps per week (GOLD 1 and 4)  
 334 ( $p < 0.05$ ). All correlations can be found in the appendix C.

335 IPAQ-sf and accelerometry - categorical variables

336 The agreement between instruments to identify “physically active” or “physically inactive” participants  
 337 was 65% and Cohen’s kappa was 0.313 (95% CI 0.146—0.480) (Table 5Table 5). The sensitivity and  
 338 specificity of IPAQ-sf 2 were 0.830 (95% CI 0.739—0.921) and 0.500 (95% CI 0.380—0.621),  
 339 respectively. PPV and NPV were 0.564 (95% CI 0.503—0.625) and 0.783 (95% CI 0.731—0.833),  
 340 respectively (Table 5).

341 **Table 5 – Comparison of the activity categories (“physically active” and “physically inactive”)**  
 342 **obtained from the IPAQ-sf 2 and accelerometer-based data (n=62).**

		Accelerometer		% agreement	Kappa (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)
		Physically Inactive	Physically Active						
IPAQ-sf 2	Physically Inactive	18	5	65%	0.313 (0.146	0.830 (0.739–	0.500 (0.380–	0.564 (0.503–	0.783 (0.731–
	Physically Active	17	22		– 0.480)				

343 **Legend:** CI, confidence intervals; IPAQ-sf, International Physical Activity Questionnaire-short form;  
 344 NPV, Negative predictive value; PPV, Positive predictive value.

345 **Discussion**

346 The present study suggests that the IPAQ-sf is valid to be used in patients with COPD and has good  
 347 test-retest reliability but with wide limits of agreement which may limit the accuracy of this instrument.  
 348 When stratifying patients by age, sex and GOLD airflow obstruction levels, the highest correlations  
 349 were found in patients with <65 years and in male patients.

350 These findings show that the IPAQ-sf may not be a reliable measure, nevertheless, patients may have  
351 also increased awareness of their PA levels by wearing the accelerometer [39, 40]. Similar results  
352 have been reported in other studies assessing the test-retest reliability of IPAQ-sf in several  
353 populations [5, 41, 42]. Results from the present study were, in general, more positive than the results  
354 from a previous exploratory study conducted in COPD [41], which revealed a lower ICC in test-retest  
355 reliability (ICC=0.439, 95% CI -0.267—0.838) and even wider limits of agreement (-10361—4548  
356 METs-min/week).

357 This study showed that, when considering the test-retest agreement using the LoA, the standard  
358 deviation of the bias of the IPAQ-sf (3279 MET-min/week) was higher than the IPAQ-sf cut-off scores  
359 for categorising individuals as “physically active” (i.e., at least 600 MET-min/week) [21]. A similar  
360 finding was observed in the MDC<sub>95</sub> (4971 METs-min/week). When analysing the IPAQ-sf categories  
361 “physically active” and “physically inactive”, the percentage of agreement was higher than when the  
362 categories “high PA level”, “moderate PA level” and “low PA level” were considered (84% vs. 66%,  
363 respectively), and above the recommended standard for reliability coefficients [16]. This can be  
364 justified by the fact that the category “physically active” includes both “high PA level” and “moderate  
365 PA level”. LoA (or the MDC<sub>95</sub>) can be considered “true” changes after an intervention [16], and the  
366 LoA were wide (i.e., higher than the IPAQ-sf cut-off scores for categorising individuals as “physically  
367 active” - at least 600 MET-min/week), the IPAQ-sf may not be appropriate to assess patients’ PA  
368 levels throughout time. This was somewhat expected since the IPAQ-sf questionnaire was originally  
369 designed for PA surveillance studies [5] and not for assessing PA changes or the impact of  
370 interventions on individuals’ PA levels. Thus, caution is needed when using the IPAQ-sf to register  
371 patients’ PA evolution/progression in PA levels in clinical practice to avoid imprecise assessment  
372 which may interfere with the tailored intervention.

373 The Stanford Seven-Day Physical Activity Recall (PAR), which was previously tested for construct  
374 validity in patients with COPD using accelerometry, showed similar results to the ones provided in the  
375 present study ( $r=0.54$ ,  $p<0.001$ ) [43]. When comparing the IPAQ-sf to the Clinical Visit PROactive tool

376 [44], this instrument presented slightly higher correlations with related constructs ( $r > 0.6$ ) and higher  
377 test-retest reliability ( $ICC \geq 0.9$ ). However, the PROactive tool is a hybrid tool (i.e., combines a short  
378 patient-reported outcome questionnaire and an activity monitor), which makes it less feasible to be  
379 used in clinical settings with low resources. Therefore IPAQ-sf seems to be an applicable  
380 questionnaire to assess PA in patients with COPD since the correlations were higher than the  
381 threshold recommended [33] in most variables and presented similar results compared to more  
382 complex instruments [45, 46]. Nevertheless, no significant correlations were found in VPA measured  
383 with the two instruments. This is somewhat expected as few patients engage in vigorous-intensity PA  
384 and its duration is normally limited [47]; and IPAQ-sf may overestimate time spent in VPA in this  
385 population. The exploratory study carried out in patients with COPD [41] revealed a higher correlation  
386 between IPAQ-sf and accelerometry than in the present study ( $r = 0.729$ ,  $p = 0.017$ ). The bigger sample  
387 size of the present study may justify the differences found between studies and suggests that larger  
388 studies should be carried out in this population to ensure more robust results.

389 In clinical practice, an accurate tool for assessing PA levels and identifying physically inactive patients  
390 is crucial to enable healthcare professionals to provide adequate advice. The IPAQ-sf may be useful  
391 for this purpose in COPD but caution is required, since it has high sensitivity but a low specificity  
392 (0.830 and 0.500, respectively), which means that the IPAQ-sf may wrongly classify individuals as  
393 “active” when they are actually “truly inactive” (low specificity, i.e., a high number of false positives  
394 having the accelerometer as the reference standard). These results are in line with a previous study  
395 [48], which has adapted and validated the IPAQ-sf to the elderly population (IPAQ-E). The authors  
396 found sensitivity results similar to the present study (81%) but higher specificity (85%), since it was  
397 an adapted version of IPAQ-sf. Future research should explore whether the IPAQ-E is more suitable  
398 for the COPD population, as most patients are older [49]. To the best of the authors’ knowledge,  
399 previous studies on measurement properties of instruments for PA assessment in patients with COPD  
400 have not provided information of sensitivity and specificity. To overcome the uncertainty of classifying  
401 a “truly inactive patient” as “physically active” with the IPAQ-sf in clinical practice, the authors suggest

402 healthcare professionals to confirm this categorisation through other methods, such as asking  
403 patients about PA routines or, if possible, perform an objective assessment using PA monitors.

404 When stratifying patients by subgroups, correlations in the subgroup of  $\geq 65$  years were below the  
405 recommended threshold ( $\rho > 0.50$ ), although significant; additionally, the strongest correlations were  
406 found in total METs-min/week in males ( $\rho = 0.653$ ,  $p < 0.001$ ) and  $< 65$  years ( $\rho = 0.651$ ,  $p < 0.001$ ), which  
407 is in line with the fact that IPAQ-sf was initially developed to people with  $< 65$  years [5] and, thus, it  
408 may not be adjusted to older people. The study of Hurting-Wennlöf's et al. [48] presented a positive  
409 correlation between self-reported activity domains with the objectively assessed PA by an  
410 accelerometer ( $\rho = 0.277$ — $0.471$ ), but with a systematic error observed.

411 Although the IPAQ-sf is widely used in several populations [6-9], this study highlights that caution  
412 should be taken when using it as an isolated indicator of PA in COPD [6].

#### 413 **4.1 Limitations and future work**

414 This study has some limitations that need to be acknowledged. The IPAQ-sf was designed to be used  
415 by adults aged 18—65 years [5] and, in this study, participants had a mean ( $\pm$ SD) age higher than  
416 that range ( $68 \pm 8$  years) which may have had influenced the results. Additionally, the original authors  
417 of the IPAQ-sf [5] recommended the “last 7 days recall” version of IPAQ-sf for studies assessing PA.  
418 However, the last 7 days may not represent the usual pattern of patients' weekly PA, which is  
419 dependent of several factors, such as weather conditions [50]. Further studies should explore the  
420 “usual week” IPAQ-sf to understand if the correlations remain consistent. Nevertheless, there was  
421 only a small percentage (10%) of patients who had less than 7 valid days of PA monitoring. Another  
422 limitation concerns to the use of accelerometers as the comparator (*gold standard*). Although they  
423 are valid to assess PA of patients with COPD [20, 25], some activities such as water-based activities  
424 and movement of the upper limbs cannot be assessed [51]. This study was conducted with stable  
425 patients with COPD hence, generalisability of results to other states of COPD and/or to other diseases  
426 is not possible. In addition, most participants in this sample were male. In female patients, no  
427 significant correlations were found between the IPAQ-sf 2 and any of the PA variables obtained

428 through accelerometry. This could be justified by the lower sample size in the female subgroup (n=9).  
429 In addition, there was some variability in the correlation results in the different GOLD grades. This  
430 finding may be partially explained by the unbalanced sample sizes in the groups, but it may also  
431 indicate that the IPAQ-sf is not an adequate tool for assessing PA levels in different airflow obstruction  
432 levels of the disease. Further research with a larger, more balanced sample of female patients and  
433 patients in the different GOLD groups and different countries, as well as longitudinal studies, are  
434 needed to reinforce these findings and to ensure external validity of findings.

### 435 **Conclusions**

436 Findings from this study showed that the IPAQ-sf presents positive and significant correlations with  
437 accelerometry, as well as high test-retest reliability but with large 95% limits of agreement, suggesting  
438 that the IPAQ-sf may not be appropriate to assess patients' PA levels throughout time. This was  
439 somehow expected since the IPAQ-sf questionnaire was originally designed for PA surveillance  
440 studies and not for assessing PA changes or the impact of interventions on individuals' PA levels.



441 **References**

- 442 1. Vorrink, S.N.W., et al., *Level of daily physical activity in individuals with COPD compared with*  
443 *healthy controls*. *Respiratory Research*, 2011. **12**: p. 1-8.
- 444 2. Waschki, B., et al., *Disease Progression and Changes in Physical Activity in Patients with*  
445 *Chronic Obstructive Pulmonary Disease*. *American journal of respiratory and critical care*  
446 *medicine*, 2015. **192**(3): p. 295-306.
- 447 3. Watz, H., et al., *An official European Respiratory Society statement on physical activity in*  
448 *COPD*. *Eur Respir J*, 2014. **44**(6): p. 1521-37.
- 449 4. Global Strategy for the Diagnosis Management and Prevention of COPD, *Global Initiative for*  
450 *Chronic Obstructive Lung Disease (GOLD)*. Available from: <https://goldcopd.org/gold-reports/>.  
451 2022.
- 452 5. Craig, C.L., et al., *International physical activity questionnaire: 12-country reliability and*  
453 *validity*. *Med Sci Sports Exerc*, 2003. **35**(8): p. 1381-95.
- 454 6. Lee, P.H., et al., *Validity of the International Physical Activity Questionnaire Short Form (IPAQ-*  
455 *SF): a systematic review*. *The international journal of behavioral nutrition and physical activity*,  
456 2011. **8**: p. 115-115.
- 457 7. Tierney, M., A. Fraser, and N. Kennedy, *Criterion validity of the International Physical Activity*  
458 *Questionnaire Short Form (IPAQ-SF) for use in patients with rheumatoid arthritis: comparison*  
459 *with the SenseWear Armband*. *Physiotherapy*, 2015. **101**(2): p. 193-197.
- 460 8. Kaleth, A.S., et al., *Validity and reliability of community health activities model program for*  
461 *seniors and short-form international physical activity questionnaire as physical activity*  
462 *assessment tools in patients with fibromyalgia*. *Disability and Rehabilitation*, 2010. **32**(5): p.  
463 353-359.
- 464 9. Ahn, G.E., et al., *Self-reported and objectively measured physical activity in adults with*  
465 *systemic lupus erythematosus*. *Arthritis care & research*, 2015. **67**(5): p. 701-707.
- 466 10. Nyssen, S.M., et al., *Levels of physical activity and predictors of mortality in COPD*. *Jornal*  
467 *Brasileiro de Pneumologia*, 2013. **39**: p. 659-666.
- 468 11. Saglam, M., et al., *Functional capacity, physical activity, and quality of life in hypoxemic*  
469 *patients with chronic obstructive pulmonary disease*. *Int J Chron Obstruct Pulmon Dis*, 2015.  
470 **10**: p. 423-8.
- 471 12. Voncken-Brewster, V., et al., *A randomized controlled trial evaluating the effectiveness of a*  
472 *web-based, computer-tailored self-management intervention for people with or at risk for*  
473 *COPD*. *Int J Chron Obstruct Pulmon Dis*, 2015. **10**: p. 1061-73.
- 474 13. Zogg, S., et al., *Differences in classification of COPD patients into risk groups A-D: a cross-*  
475 *sectional study*. *BMC Research Notes*, 2014. **7**(1): p. 562.
- 476 14. Gimeno-Santos, E., et al., *Determinants and outcomes of physical activity in patients with*  
477 *COPD: a systematic review*. *Thorax*, 2014. **69**(8): p. 731-739.
- 478 15. Lidwine B Mokkink, et al., *COSMIN methodology for systematic reviews of Patient-Reported*  
479 *Outcome Measures (PROMs)*. 2018.
- 480 16. Terwee, C.B., et al., *Qualitative attributes and measurement properties of physical activity*  
481 *questionnaires: a checklist*. *Sports Med*, 2010. **40**(7): p. 525-37.
- 482 17. Doherty, D.E., et al., *Chronic Obstructive Pulmonary Disease: Consensus Recommendations*  
483 *for Early Diagnosis and Treatment*. *J Fam Pract*, 2006. **Supplement**: p. S1-S8.
- 484 18. Charlson, M.E., et al., *A new method of classifying prognostic comorbidity in longitudinal*  
485 *studies: development and validation*. *J Chronic Dis*, 1987. **40**(5): p. 373-83.

- 486 19. Miller, M.R., et al., *Standardisation of spirometry*. European Respiratory Journal, 2005. **26**(2):  
487 p. 319-338.
- 488 20. Rabinovich, R.A., et al., *Validity of physical activity monitors during daily life in patients with*  
489 *COPD*. Eur Respir J, 2013. **42**(5): p. 1205-1215.
- 490 21. IPAQ, *Guidelines for the data processing and analysis of the International Physical Activity*  
491 *Questionnaire*. 2005.
- 492 22. Marshall, A.L., et al., *Reliability and validity of a brief physical activity assessment for use by*  
493 *family doctors*. Br J Sports Med, 2005. **39**(5): p. 294-7; discussion 294-7.
- 494 23. Puig-Ribera, A., et al., *Screening Physical Activity in Family Practice: Validity of the Spanish*  
495 *Version of a Brief Physical Activity Questionnaire*. PLOS ONE, 2015. **10**(9): p. e0136870.
- 496 24. Wanner, M., et al., *Validation of the long international physical activity questionnaire: Influence*  
497 *of age and language region*. Preventive Medicine Reports, 2016. **3**: p. 250-256.
- 498 25. Van Remoortel, H., et al., *Validity of activity monitors in health and chronic disease: a*  
499 *systematic review*. International Journal of Behavioral Nutrition and Physical Activity, 2012.  
500 **9**(1): p. 84.
- 501 26. Demeyer, H., et al., *Standardizing the analysis of physical activity in patients with copd*  
502 *following a pulmonary rehabilitation program*. Chest, 2014. **146**(2): p. 318-327.
- 503 27. Garber, C.E., et al., *American College of Sports Medicine position stand. Quantity and quality*  
504 *of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and*  
505 *neuromotor fitness in apparently healthy adults: guidance for prescribing exercise*. Medicine  
506 and Science in Sports and Exercise, 2011. **43**(7): p. 1334-1359.
- 507 28. Freedson, P.S., E. Melanson, and J. Sirard, *Calibration of the Computer Science and*  
508 *Applications, Inc. accelerometer*. Medicine and Science in Sports and Exercise, 1998. **30**(5):  
509 p. 777-781.
- 510 29. WHO. *WHO Guidelines on Physical Activity and Sedentary Behaviour 2020*; Available from:  
511 [https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-](https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi)  
512 [lifestyle/body-mass-index-bmi](https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi).
- 513 30. Kottner, J., et al., *Guidelines for Reporting Reliability and Agreement Studies (GRRAS) were*  
514 *proposed*. International Journal of Nursing Studies, 2011. **48**(6): p. 661-671.
- 515 31. Prinsen, C.A.C., et al., *COSMIN guideline for systematic reviews of patient-reported outcome*  
516 *measures*. Quality of life research : an international journal of quality of life aspects of  
517 treatment, care and rehabilitation, 2018. **27**(5): p. 1147-1157.
- 518 32. Watson, P.F. and A. Petrie, *Method agreement analysis: a review of correct methodology*.  
519 Theriogenology, 2010. **73**(9): p. 1167-79.
- 520 33. Terwee, C.B., *Qualitative Attributes and Measurement Properties of Physical Activity*  
521 *Questionnaires*. Sports Medicine, 2010. **40**(7): p. 525-537.
- 522 34. Bland, M.J. and D.G. Altman, *Statistical Methods for Assessing Agreement between Two*  
523 *Methods of Clinical Measurement*. The Lancet, 1986. **327**(8476): p. 307-310.
- 524 35. Landis, G.K.J., *The measurement of observer agreement for categorical data*. Biometrics,  
525 1977. **33**(1): p. 159-174.
- 526 36. Sattler, M.C., et al., *Current Evidence of Measurement Properties of Physical Activity*  
527 *Questionnaires for Older Adults: An Updated Systematic Review*. Sports Medicine, 2020.  
528 **50**(7): p. 1271-1315.
- 529 37. Evans, J.D., *Straightforward statistics for the behavioral sciences*. 1996, Pacific Groove:  
530 Brooks/Cole Publishing.

- 531 38. IPAQ Research Committee, *Guidelines for Data Processing and Analysis of the International*  
532 *Physical Activity Questionnaire (IPAQ) – Short and Long Form*. 2005:  
533 <https://sites.google.com/site/theipaq/scoring-protocol>.
- 534 39. Blikman, T., et al., *Reliability and validity of the Dutch version of the International Physical*  
535 *Activity Questionnaire in patients after total hip arthroplasty or total knee arthroplasty*. *J Orthop*  
536 *Sports Phys Ther*, 2013. **43**(9): p. 650-9.
- 537 40. Wendel-Vos, G.C., et al., *Reproducibility and relative validity of the short questionnaire to*  
538 *assess health-enhancing physical activity*. *J Clin Epidemiol*, 2003. **56**(12): p. 1163-9.
- 539 41. Cruz, J., C. Jácome, and A. Marques, *Is the international physical activity questionnaire*  
540 *(IPAQ-sf) valid to assess physical activity in patients with COPD? Comparison with*  
541 *accelerometer data*. *BMC Health Services Research*, 2018. **18**(2): p. 684.
- 542 42. Faulkner, G., T. Cohn, and G. Remington, *Validation of a physical activity assessment tool for*  
543 *individuals with schizophrenia*. *Schizophr Res*, 2006. **82**(2-3): p. 225-31.
- 544 43. Garfield, B.E., et al., *Stanford Seven-Day Physical Activity Recall questionnaire in COPD*.  
545 *European Respiratory Journal*, 2012. **40**(2): p. 356.
- 546 44. Gimeno-Santos, E., et al., *The PROactive instruments to measure physical activity in patients*  
547 *with chronic obstructive pulmonary disease*. *Eur Respir J*, 2015. **46**(4): p. 988-1000.
- 548 45. Copeland, J.L. and D.W. Esliger, *Accelerometer assessment of physical activity in active,*  
549 *healthy older adults*. *J Aging Phys Act*, 2009. **17**(1): p. 17-30.
- 550 46. Swartz, A.M., et al., *Estimation of energy expenditure using CSA accelerometers at hip and*  
551 *wrist sites*. *Med Sci Sports Exerc*, 2000. **32**(9 Suppl): p. S450-6.
- 552 47. Hartman, J.E., et al., *Physical Activity Recommendations in Patients with Chronic Obstructive*  
553 *Pulmonary Disease*. *Respiration Physiology*, 2014.
- 554 48. Hurtig-Wennlöf, A., M. Hagströmer, and L.A. Olsson, *The International Physical Activity*  
555 *Questionnaire modified for the elderly: aspects of validity and feasibility*. *Public Health Nutr*,  
556 2010. **13**(11): p. 1847-54.
- 557 49. Raheison, C. and P.O. Girodet, *Epidemiology of COPD*. *European Respiratory Review*, 2009.  
558 **18**(114): p. 213.
- 559 50. Chan, C.B. and D.A. Ryan, *Assessing the effects of weather conditions on physical activity*  
560 *participation using objective measures*. *Int J Environ Res Public Health*, 2009. **6**(10): p. 2639-  
561 54.
- 562 51. Lee, I.M. and E.J. Shiroma, *Using accelerometers to measure physical activity in large-scale*  
563 *epidemiological studies: issues and challenges*. *Br J Sports Med*, 2014. **48**(3): p. 197-201.

564

565

566 **Appendix A- Percentage of agreement and weighted Cohen's kappa among IPAQ-sf categories**  
 567 **(“low PA”, “moderate PA” and “high PA”) (n=62).**

	IPAQ-sf 2			% agreement	Kappa (95% CI)
	Low PA	Moderate PA	High PA		
IPAQ-sf 1	Low PA	19	2	66%	0.496 (0.329 to 0.663)
	Moderate PA	4	13		
	High PA	0	2		

568 **Legend:** CI, confidence intervals; IPAQ-sf, International Physical Activity Questionnaire-short form;  
 569 PA, physical activity.

570 **Appendix B - Correlations ( $\rho$ ) between IPAQ-sf 2 and accelerometry (n=62).**

Source	Correlations (min/week)	$\rho$
IPAQ-sf	Total METs-min/week	0.527**
Accelerometry	Time in MVPA	
IPAQ-sf	Time in VPA	0.006
Accelerometry	Time in VPA	
IPAQ-sf	Time in MPA	0.444**
Accelerometry	Time in MPA	
IPAQ-sf	Time in Walking	0.396**
Accelerometry	Time in MPA	
IPAQ-sf	Time in walking	0.434**
Accelerometry	Number of steps/week	

571 **Legend:** IPAQ-sf, International Physical Activity Questionnaire-short form; MPA, moderate physical  
 572 activity; MVPA, moderate to vigorous physical activity; PA, physical activity; VPA, vigorous physical  
 573 activity. \*  $p < 0.05$  \*\* $p < 0.001$ .

574 **Appendix C - Correlations ( $\rho$ ) between IPAQ-sf 2 and accelerometer-based data stratified by**  
 575 **age, sex and GOLD grades.**


Source	(min/week)	Age		Sex		GOLD airflow obstruction levels			
		< 65 years (n=20)	$\geq$ 65 years (n=42)	Male (n=53)	Female (n=9)	GOLD 1 (n=5)	GOLD 2 (n=25)	GOLD 3 (n=20)	GOLD 4 (n=12)
IPAQ-sf 2	Total METs- min	0.651**	0.443**	0.653**	-0.450	0.300	0.491**	0.437	0.635*
AC	Total MVPA								
IPAQ-sf 2	Time in VPA	0.240	-0.152	0.092	-0.359	0.057	0.242	-0.317	0.305
AC	Time in VPA								
IPAQ-sf 2	Time in MPA	0.517*	0.393*	0.524**	-0.294	0.051	0.431*	0.352	0.541
AC	Time in MPA								
IPAQ-sf 2	Time in Walking	0.467*	0.395**	0.466**	-0.159	0.564	0.159	0.312	0.640*
AC	Time in MPA								
IPAQ-sf 2	Time in MPA and walking	0.377	0.444**	0.507**	-0.594	0.975**	0.232	0.282	0.707*
AC	Number of steps/week								

576 **Legend:** AC, accelerometry; COPD, Chronic Obstructive Pulmonary Disease; IPAQ-sf, International  
 577 Physical Activity Questionnaire-short form; MPA, moderate physical activity; MVPA, moderate to  
 578 vigorous physical activity; PA, physical activity; VPA, vigorous physical activity. \*  $p < 0.05$  \*\* $p < 0.001$ .

579

580 **Appendix D – Bland and Altman plots (n=62).**


581 Figures 2, 3 and 4 present a Bland and Altman plot with the 95% LoA between the IPAQ-sf 2 and  
 582 accelerometry regarding VPA (mean differences (bias) =45 min/week, SD of bias= 91 min/week,  
 583 95% LoA= -135 – 224 min/week), MPA (bias= 18 min/week, SD of bias= 254 min/week, 95% LoA= -  
 584 480 – 515 min/week) and Walking (bias= 35 min/week, SD of bias= 268 min/week, 95% LoA= -491 –  
 585 561 min/week), respectively.



586


587 **Figure 2 - Bland and Altman plots for vigorous physical activity (VPA) in patients with chronic**  
 588 **obstructive pulmonary disease (n=62). Comparison between IPAQ-sf 2 and accelerometry**  
 589 **measurements (min/week).**

590



591

592 **Figure 3 - Bland and Altman plots for moderate physical activity (MPA) in patients with chronic**  
 593 **obstructive pulmonary disease (n=62). Comparison between IPAQ-sf 2 and accelerometry**  
 594 **measurements (min/week).**



595

596 **Figure 4 - Bland and Altman plots for walking in patients with chronic obstructive pulmonary disease**597 **(n=62). Comparison between IPAQ-sf 2 and accelerometry measurements (min/week).**

Journal Pre-proof

### Highlights

1. The IPAQ-sf presented good test-retest reliability results but wide limits of agreement.
2. The wide LoA suggest that it may not be appropriate to assess patients' PA levels throughout time.
3. Overall, the IPAQ-sf showed positive and moderate correlations with accelerometry.
4. The IPAQ-sf seems to be valid in COPD but caution on its widespread use is recommended as its specificity is low.



**Declarations of interest:**

All authors declare that they have no conflicts of interest.

Journal Pre-proof