

REGULARIZATION ALGORITHMS FOR LINEAR COPOSITIVE PROBLEMS

OLGA I. KOSTYUKOVA^{1,2}  AND TATIANA V. TCHEMISOVA^{3,*} 

Abstract. The paper is devoted to the regularization of linear Copositive Programming problems which consists of transforming a problem to an equivalent form, where the Slater condition is satisfied and therefore the strong duality holds. We describe regularization algorithms based on a concept of immobile indices and on the understanding of the important role that these indices play in the feasible sets' characterization. These algorithms are compared to some regularization procedures developed for a more general case of convex problems and based on a facial reduction approach. We show that the immobile-index-based approach combined with the specifics of copositive problems allows us to construct more explicit and detailed regularization algorithms for linear Copositive Programming problems than those already available.

Mathematics Subject Classification. 90C25, 90C46, 90C30, 90C34.

Received December 15, 2020. Accepted May 4, 2022.

1. INTRODUCTION

Conic optimization is a subfield of convex optimization that studies the problems of minimizing a convex function over the intersection of an affine subspace and a convex cone. For a gentle introduction to conic optimization and a survey of its applications in Operations Research and related areas, we refer interested readers to [15] and the references therein.

Copositive Programming (CoP) problems form a special class of conic problems and can be considered as an optimization over the convex cone of so-called *copositive matrices* (*i.e.* matrices which are positive semi-defined on the non-negative orthant). Copositive models arise in many important applications, including \mathcal{NP} -hard problems. For the references on motivation and application of CoP see, *e.g.* [3, 7, 9].

In *linear* CoP, the objective function is linear and the constraints are formulated with the help of linear matrix functions. Linear copositive problems are closely related to that of linear *Semi-Infinite Programming* (SIP) and *Semidefinite Programming* (SDP). Copositive and semidefinite problems are particular cases of SIP problems, but CoP deals with more challenging and less studied problems than SDP. The literature on the theory and

Keywords. Linear copositive programming, strong duality, normalized immobile index set, regularization, minimal cone, facial reduction, constraint qualifications.

¹ Institute of Mathematics, National Academy of Sciences of Belarus, Surganov str. 11, 220072 Minsk, Belarus.

² Department of Applied Mathematics and Computer Science, Belarusian State University, Nezavisimosti av. 4, 220030 Minsk, Belarus.

³ Mathematical Department, University of Aveiro, Campus Universitário Santiago, 3810-193 Aveiro, Portugal.

*Corresponding author: tatiana@ua.pt

methods of SIP, CoP, and SDP is quite extensive. We refer the interested readers to [1–3, 9, 25, 26], and the references in these works.

In convex and conic optimization, optimality conditions, and duality results are usually formulated under certain regularity conditions, so-called constraint qualifications (CQ) (see, *e.g.* [2, 10, 22, 26]). Such conditions should guarantee the fulfillment of the Karush–Kuhn–Tucker (KKT)-type optimality conditions and the *strong duality* property consisting in the fact that the optimal values of the primal problem and the corresponding Lagrangian dual one, are equal and the dual problem attains its maximum. Strong duality is the cornerstone of convex optimization, playing a particularly important role in the stability of numerical methods.

Unfortunately, even in convex optimization, many problems cannot be classified as regular (*i.e.* satisfying some regularity conditions such as, for example, strict feasibility). In [8], we read: “. . . *new optimization modeling techniques and convex relaxations for hard nonconvex problems have shown that the loss of strict feasibility is a more pronounced phenomenon than has previously been realized*”. This phenomenon can occur because of either the poor choice of functions that describe feasible sets or the degeneration of the feasible sets themselves. According to [23], sometimes the loss of a certain CQ “. . . *is a modeling issue rather than inherent to the problem instance. . .*” which “. . . *justifies the pleasing paradigm: efficient modeling provides for a stable program*”.

Thus, the idea of a *regularization* appears quite naturally which is aimed at obtaining an equivalent and more convenient reformulation of the problem with some required properties, one of which is that the regularized problem must satisfy the generalized Slater condition.

The first works on the regularization of abstract convex problems (regularization procedures are called *pre-processing* there) appeared in the 1980-s, followed by various publications on special classes of conic problems (see, *e.g.* [5, 6]). Nevertheless, as Drusvyatskiy and Wolkowicz wrote in their paper [8] published in 2017, for conic optimization in general, the research in the field of regularization algorithms is still in its infancy. At the same time, the authors of [8] confirm that in order to make a regularization algorithm viable, it is necessary to actively explore the structure of the problem, since for some specific applications of conic optimization, a rich basic structure makes regularization quite possible and leads to significantly simplified models and enhanced algorithms.

Several approaches to the regularization of conic optimization problems are proposed in the literature. In [5, 6], the concept of *minimal cone* of constraints was used by Borwein and Wolkowicz to regularize abstract convex and conic convex problems for which any CQ fails. The algorithm proposed there to describe the minimal cone is based on the sequential reduction of the cone’s faces and was named by the authors *Facial Reduction Algorithm (FRA)*.

Another approach called the *dual regularization* or *conic expansion* was proposed by Luo, Sturm, and Zhang (see [17] and the references therein). This approach tries to close the *duality gap* (the difference between the primal and dual optimal values) of the regularized problems by expanding the dual constraints’ cone.

In [24], Waki and Muramatsu applied the facial reduction approach to a conic optimization problem in such a way that each primal reduced cone is dual to the cone generated by the conic expansion approach.

The facial reduction approach has been successfully applied to SDP and second-order cone programming problems, as well as to certain classes of optimization problems over symmetric (*i.e.* self-dual and homogeneous) and nice cones (see, *e.g.* [18–21]). At the same time, the question of effective constructive application of this approach to other classes of problems remains open. This is because the known FRAs are more *conceptual* than practical.

In this paper, based on the results from [11, 12, 14], we develop a different approach to regularization of linear CoP problems. This approach is based on the concept of *immobile indices*, *i.e.* indices of the constraints that are active for all feasible solutions.

The purpose of the paper is to

- (a) describe in details a finite algorithm for regularization of linear CoP problems that is based on the concept of immobile indices but does not require any additional information about them;

- (b) analogize two approaches to the regularization of linear CoP problems, one based on facial reduction and the other on the concept of immobile indices, and to compare the corresponding regularized problems constructed using these approaches.

To the best of our knowledge, in CoP there has never been an attempt to develop detailed and easy-to-use algorithms, based on the minimal cone representation (see, *e.g.* the FRA in [5, 6] and the modified FRA in [24]). Nor do we have any information about any other attempts to describe constructive regularization procedures for linear copositive problems. The regularization algorithms presented in the paper are new, original, and timely due to the growing number of eminent applications of CoP.

The paper is organized as follows. Section 2 contains equivalent formulations of the linear CoP problem and the basic definitions. In Section 3, we consider two different approaches to regularization of copositive problems. In Section 3.1, we show how the minimal face regularization from [5, 6] can be applied to linear CoP problems; in Section 3.2, we briefly describe the one-step regularization proposed in [14] and based on the concept of immobile indices, and compare the regularized problems obtained in this subsection with the problem in Section 3.1. Section 4 is devoted to iterative algorithms for regularization of linear copositive problems. The Waki and Muramatsu’s facial reduction algorithm is described in Section 4.1, a new regularization algorithm **REG-LCoP** based on the immobile index set together with its compressed modification is introduced, justified, and compared with the Waki and Muramatsu FRA in Section 4.2. A small clarifying example is proposed. We conclude Section 4 with a brief discussion on the algorithms considered there. Section 5 contains some conclusions.

2. LINEAR COPOSITIVE PROGRAMMING PROBLEM: EQUIVALENT FORMULATIONS AND BASIC DEFINITIONS

Given an integer $p > 1$, denote by \mathbb{R}_+^p the set of all p vectors with non-negative components, by $\mathcal{S}(p)$ and $\mathcal{S}_+(p)$ the space of real symmetric $p \times p$ matrices and the cone of symmetric positive semidefinite $p \times p$ matrices, respectively, and let \mathcal{COP}^p stay for the cone of symmetric copositive $p \times p$ matrices:

$$\mathcal{COP}^p := \{D \in \mathcal{S}(p) : t^\top D t \geq 0 \ \forall t \in \mathbb{R}_+^p\}.$$

The space $\mathcal{S}(p)$ is considered here as a vector space with the trace inner product

$$A \bullet B := \text{trace}(AB).$$

Consider a linear copositive programming problem in the form

$$\min_{x \in \mathbb{R}^n} c^\top x, \quad \text{s.t. } \mathcal{A}(x) \in \mathcal{COP}^p, \tag{2.1}$$

where $x = (x_1, \dots, x_n)^\top$ is the vector of decision variables. The data of the problem are presented by vector $c \in \mathbb{R}^n$ and the constraints matrix function $\mathcal{A}(x)$ defined in the form

$$\mathcal{A}(x) := \sum_{i=1}^n A_i x_i + A_0, \tag{2.2}$$

with given matrices $A_i \in \mathcal{S}(p), i = 0, 1, \dots, n$. It is well known (see *e.g.* [1]) that the copositive problem (2.1) is equivalent to the following convex SIP problem:

$$\min_{x \in \mathbb{R}^n} c^\top x, \quad \text{s.t. } t^\top \mathcal{A}(x) t \geq 0 \ \forall t \in T, \tag{2.3}$$

with a p -dimensional compact index set in the form of a simplex

$$T := \{t \in \mathbb{R}_+^p : \mathbf{e}^\top t = 1\}, \tag{2.4}$$

where $\mathbf{e} = (1, 1, \dots, 1)^\top \in \mathbb{R}^p$.

Denote by X the feasible set of the equivalent problems (2.1) and (2.3):

$$X := \{x \in \mathbb{R}^n : \mathcal{A}(x) \in \mathcal{COP}^p\} = \{x \in \mathbb{R}^n : t^\top \mathcal{A}(x)t \geq 0 \ \forall t \in T\}. \tag{2.5}$$

In what follows, we will suppose that $X \neq \emptyset$. Evidently, the set X is convex.

Remark 2.1. Since $X \neq \emptyset$, then without loss of generality, we can consider that $A_0 \in \mathcal{COP}^p$. Indeed, by fixing a feasible solution $y \in X$ and substituting the variable x by a new variable $z := x - y$, we can replace the original problem (2.1) by the following one in terms of z :

$$\min_{z \in \mathbb{R}^n} c^\top z, \quad \text{s.t. } \bar{\mathcal{A}}(z) \in \mathcal{COP}^p,$$

with $\bar{\mathcal{A}}(z) := \sum_{i=1}^n A_i z_i + \bar{A}_0$, $\bar{A}_0 := \mathcal{A}(y) \in \mathcal{COP}^p$.

According to the commonly used definition, the constraints of the copositive problem (2.1) satisfy the Slater condition if

$$\exists \bar{x} \in \mathbb{R}^n \text{ such that } \mathcal{A}(\bar{x}) \in \text{int } \mathcal{COP}^p = \{D \in \mathcal{S}(p) : t^\top D t > 0 \ \forall t \in \mathbb{R}_+^p, t \neq \mathbf{0}\}. \tag{2.6}$$

Here $\text{int } \mathcal{B}$ stands for the interior of a set \mathcal{B} .

Following [11, 14], let's define the set of normalized immobile indices T_{im} in problem (2.1):

$$T_{im} := \{t \in T : t^\top \mathcal{A}(x)t = 0 \ \forall x \in X\}. \tag{2.7}$$

In what follows, the elements of the set T_{im} are called immobile indices.

The following lemma follows from Proposition 1 and Lemma 1 in [14].

Lemma 2.2. Given the linear copositive problem (2.1),

- (i) the Slater condition (2.6) is equivalent to the emptiness of the set T_{im} ,
- (ii) the normalized immobile index set T_{im} is either empty or can be represented as a union of a finite number of convex closed bounded polyhedra.

For a vector $t = (t_k, k \in P)^\top \in \mathbb{R}_+^p$ with $P := \{1, 2, \dots, p\}$, define the sets

$$P_+(t) := \{k \in P : t_k > 0\}, \quad P_0(t) := P \setminus P_+(t).$$

Given a set \mathcal{B} and a point $l = (l_k, k \in P)^\top$ in \mathbb{R}^p , denote by $\rho(l, \mathcal{B})$ the distance between these set and point, $\rho(l, \mathcal{B}) := \min_{\tau \in \mathcal{B}} \sum_{k \in P} |l_k - \tau_k|$, and by $\text{conv } \mathcal{B}$ the convex hull of the set \mathcal{B} .

Suppose that in problem (2.1), the normalized immobile index set T_{im} is non-empty. Consider a finite non-empty subset of T_{im} :

$$V = \{\tau(i), i \in I\} \subset T_{im}, \quad 0 < |I| < \infty. \tag{2.8}$$

For this set, define the following number and sets:

$$\sigma(V) := \min\{\tau_k(i), k \in P_+(\tau(i)), i \in I\} > 0, \tag{2.9}$$

$$\Omega(V) := \{t \in T : \rho(t, \text{conv } V) \geq \sigma(V)\}, \tag{2.10}$$

$$\mathcal{X}(V) := \{x \in \mathbb{R}^n : \mathcal{A}(x)\tau(i) \geq 0 \ \forall i \in I; t^\top \mathcal{A}(x)t \geq 0 \ \forall t \in \Omega(V)\}. \tag{2.11}$$

In [12], the following theorem is proved (see Thm. 7.1 in [12]).

Theorem 2.3. Consider problem (2.1) with the feasible set X . For any subset (2.8) of the set of normalized immobile indices of this problem, the equality $X = \mathcal{X}(V)$, holds true, where the set $\mathcal{X}(V)$ is defined in (2.11).

Let $\{e_k, k \in P\}$ be the standard basis of \mathbb{R}^P .

In what follows we will need the following proposition.

Proposition 2.4. *For any $\tau \in T_{im}$, the following relations hold true:*

$$e_k^\top \mathcal{A}(x)\tau = 0 \quad \forall k \in P_+(\tau), \quad e_k^\top \mathcal{A}(x)\tau \geq 0 \quad \forall k \in P_0(\tau), \quad \forall x \in X. \tag{2.12}$$

Proof. Remind that

$$t^\top \mathcal{A}(x)t \geq 0 \quad \forall t \in T, \quad \forall x \in X, \tag{2.13}$$

and by definition, for any vector $\tau \in T_{im}$, we have $\tau^\top \mathcal{A}(x)\tau = 0 \quad \forall x \in X$. Hence, for any $x \in X$, this vector τ is an optimal solution to the following quadratic programming problem:

$$\text{QP} : \min t^\top \mathcal{A}(x)t \quad \text{s.t. } t \in T.$$

Notice that relations (2.12) are nothing else than the first order necessary optimality conditions for the vector τ in problem (QP). The proposition is proved. \square

3. REGULARIZATION OF COPOSITIVE PROBLEMS

In this section, first, we remind a known regularization approach developed in [5, 6] for conic optimization problems and based on the concept of the minimal face. We briefly describe how this approach can be applied to linear CoP problems. After, for the copositive problem (2.1), we present another regularization approach based on the concept of immobile indices and compare the regularized problems obtained using two considered approaches.

3.1. Minimal face regularization

Let us, first, recall the necessary terms and notions.

For a given cone $\mathbf{F} \subset \mathcal{S}(p)$, its dual cone is defined as follows:

$$\mathbf{F}^* := \{D \in \mathcal{S}(p) : D \bullet B \geq 0 \quad \forall B \in \mathbf{F}\}.$$

By definition, a convex subset \mathbf{F} of the cone \mathcal{COP}^p is its *face* if for any $x \in \mathcal{COP}^p, y \in \mathcal{COP}^p$, the inclusion $x + y \in \mathbf{F}$ implies $x \in \mathbf{F}, y \in \mathbf{F}$. It is evident that any face of the cone \mathcal{COP}^p is also a cone.

Given the copositive problem (2.1) with the feasible set X presented in (2.5), let \mathbf{F}_{\min} be the smallest (by inclusion) face of \mathcal{COP}^p containing a set \mathcal{D} defined in terms of the constraints of this problem as follows:

$$\mathcal{D} := \{\mathcal{A}(x), x \in X\}. \tag{3.1}$$

In what follows, the face \mathbf{F}_{\min} will be called *the minimal face of the optimization problem (2.1)*.

Generally speaking, for the copositive problem (2.1), the approach suggested in [5, 6], is to replace the constraint $\mathcal{A}(x) \in \mathcal{COP}^p$ with an equivalent constraint $\mathcal{A}(x) \in \mathbf{F}_{\min}$. The resulting regularized problem takes the form

$$\min_{x \in \mathbb{R}^n} c^\top x, \quad \text{s.t. } \mathcal{A}(x) \in \mathbf{F}_{\min}. \tag{3.2}$$

The dual problem to (3.2) can be written in the form

$$\max_{U \in \mathcal{S}(p)} -A_0 \bullet U, \quad \text{s.t. } A_j \bullet U = c_j \quad \forall j = 1, \dots, n; \quad U \in \mathbf{F}_{\min}^*, \tag{3.3}$$

where \mathbf{F}_{\min}^* is the dual cone to the cone \mathbf{F}_{\min} .

It is proved in [5, 6], that the constraints of problem (3.2) satisfy the *generalized Slater condition*: there exists $\bar{x} \in X$ such that $\mathcal{A}(\bar{x}) \in \text{reint } \mathbf{F}_{\min}$ and hence the duality gap between the dual problems (3.2) and (3.3) vanishes. Here $\text{reint } \mathcal{B}$ stays for the relative interior of a set \mathcal{B} .

Unfortunately, there is no information available about how to explicitly construct the cones \mathbf{F}_{\min} and \mathbf{F}_{\min}^* in a general case and, in particular, in the case of copositive problems.

3.2. One-step regularization based on the concept of immobile indices

In our paper [14], for the copositive problem (2.1), we obtained a regularized dual problem that is different from (3.3). The construction of this dual is based on the concept of immobile indices and can be thought of as *one-step regularization* because it contains a unique step.

Consider the copositive problem (2.1). Let T_{im} be the normalized set of immobile indices of this problem defined in (2.7).

If $T_{im} = \emptyset$, then problem (2.1) satisfies the Slater condition, which means that it is already regular and no regularization is required. Now, suppose that $T_{im} \neq \emptyset$. In this case, the Slater condition is not satisfied and the problem is not regular. Let us describe how one can convert problem (2.1) into a regularized one.

Consider the set $\text{conv } T_{im}$ and the set W of all vertices of $\text{conv } T_{im}$:

$$W := \{t(j), j \in J\}, \quad 0 < |J| < \infty. \tag{3.4}$$

Suppose that the elements $t(j)$, $j \in J$, of the set W are known. Then we can regularize problem (2.1) in just **one step**.

In fact, it follows from Theorem 2.3 above and Theorem 3.2 and Corollary 3.3 in [12], that the set X of feasible solutions of the original problem (2.1) coincides with the set of feasible solutions of the following system:

$$t^\top \mathcal{A}(x)t \geq 0 \quad \forall t \in \Omega(W); \quad \mathcal{A}(x)t(i) \geq 0 \quad \forall i \in J,$$

and the next condition is satisfied:

$$\exists \bar{x} \in X \quad \text{such that} \quad t^\top \mathcal{A}(\bar{x})t > 0 \quad \forall t \in \Omega(W). \tag{3.5}$$

Here the set $\Omega(W)$ is defined by the rules (2.10) with $V = W$.

Consequently, the original copositive problem (2.1) is equivalent to the following SIP problem:

$$\min_{x \in \mathbb{R}^n} c^\top x, \tag{3.6}$$

$$\text{s.t. } t^\top \mathcal{A}(x)t \geq 0 \quad \forall t \in \Omega(W), \tag{3.7}$$

$$\mathcal{A}(x)t(i) \geq 0 \quad \forall i \in J. \tag{3.8}$$

Problem (3.6)–(3.8) can be considered as a *regularized* primal problem since

- it possesses a finite number of linear inequality constraints (3.8),
- the first group of constraints (3.7), satisfies the Slater type condition (3.5),
- the set $\Omega(W)$ is compact.

Let us stress that in problem (3.6)–(3.8), the infinite index set $\Omega(W)$ is obtained by removing the set T_{im} together with the $\sigma(W)$ -neighborhood of its convex hull, from the original index set T . Note here that the set $\Omega(W)$

- (a) is explicitly constructed by the rules (2.9), (2.10), using the finite set $W = \{t(j), j \in J\}$ of vertices of $\text{conv } T_{im}$,
- (b) does not contain the set $\text{conv } T_{im}$,
- (c) may be sufficiently small.

All these properties may be useful for numerical solving the problem (3.6)–(3.8).

It is evident that problem (3.6)–(3.8) can be written in the equivalent conic form

$$\min_{x \in \mathbb{R}^n} c^\top x, \quad \text{s.t. } \mathcal{A}(x) \in \mathcal{K}_0, \tag{3.9}$$

where

$$\mathcal{K}_0 := \{D \in \mathcal{S}(p) : t^\top Dt \geq 0 \quad \forall t \in \Omega(W); \quad Dt(j) \geq 0 \quad \forall j \in J\}.$$

It can be shown that $\mathcal{K}_0 \subset \mathcal{COP}^p$.

The dual problem to (3.9) is as follows:

$$\max_{U \in \mathcal{S}(p)} -A_0 \bullet U, \quad \text{s.t. } A_j \bullet U = c_j \quad \forall j = 1, \dots, n; \quad U \in \mathcal{K}_0^*. \tag{3.10}$$

In the problem above, \mathcal{K}_0^* is the dual cone to \mathcal{K}_0 and has the form

$$\mathcal{K}_0^* = \text{cl}\{D \in \mathcal{S}(p) : D \in \mathcal{CP}(W) \oplus \mathcal{P}^*\}, \tag{3.11}$$

where

$$\begin{aligned} \mathcal{CP}(W) &:= \text{conv}\{tt^\top : t \in \Omega(W)\}, \\ \mathcal{P}^* &:= \left\{ D \in \mathcal{S}(p) : D = \sum_{j \in J} (\lambda(j)(t(j))^\top + t(j)(\lambda(j))^\top), \lambda(j) \geq 0 \quad \forall j \in J \right\}. \end{aligned}$$

Here and in what follows, for given sets \mathcal{B} and \mathcal{G} , $\text{cl}\mathcal{B}$ denotes the closure of the set \mathcal{B} and $\mathcal{B} \oplus \mathcal{G}$ stays for the Minkowski sum of the corresponding two sets.

Notice that for the pair of dual conic problems (3.9) and (3.10), the duality gap is zero.

As it was shown in [12], the cone (3.11) in problem (3.10) can be replaced by the following one (which has a more explicit form since it does not contain the closure operator):

$$\mathbf{K}_0^* := \{D \in \mathcal{S}(p) : D \in \mathcal{CP}^p \oplus \mathcal{P}^*\},$$

where \mathcal{CP}^p denotes the set of completely positive matrices:

$$\mathcal{CP}^p := \text{conv}\{tt^\top : t \in \mathbb{R}_+^p\}, \tag{3.12}$$

and there is no duality gap for problem (3.9) and its dual problem in the form (3.10) with \mathcal{K}_0^* replaced by \mathbf{K}_0^* .

Note that the cones \mathcal{K}_0 and \mathbf{K}_0^* are explicitly described in terms of indices (3.4) and this is an advantage of the approach presented here over the one from 3.1.

The only drawback of the regularization procedure described here is the following: *to apply the one-step regularization, one needs to know the finite number of indices (3.4) which are the vertices of the set $\text{conv } T_{im}$.*

Let us show that the regularized primal problem (3.9) can be modified as follows:

$$\min_{x \in \mathbb{R}^n} c^\top x, \quad \text{s.t. } \mathcal{A}(x) \in \bar{\mathcal{K}}_0, \tag{3.13}$$

where

$$\begin{aligned} \bar{\mathcal{K}}_0 = \left\{ D \in \mathcal{S}(p) : t^\top Dt \geq 0 \quad \forall t \in \Omega(W), \quad e_k^\top Dt(j) = 0 \quad \forall k \in P_+(t(j)), \right. \\ \left. e_k^\top Dt(j) \geq 0 \quad \forall k \in P \setminus P_+(t(j)), \quad \forall j \in J \right\}. \end{aligned}$$

In fact, due to Theorem 2.3 we have $X = \{x \in \mathbb{R}^n : \mathcal{A}(x) \in \mathcal{K}_0\}$. It is evident that $\bar{\mathcal{K}}_0 \subset \mathcal{K}_0$ and hence $\{x \in \mathbb{R}^n : \mathcal{A}(x) \in \bar{\mathcal{K}}_0\} \subset X$. On the other hand, taking into account the inclusion $\{t(j), j \in J\} \subset T_{im}$ and Proposition 2.4, we conclude that $X \subset \{x \in \mathbb{R}^n : \mathcal{A}(x) \in \bar{\mathcal{K}}_0\}$. Hence we have shown that $X = \{x \in \mathbb{R}^n : \mathcal{A}(x) \in \bar{\mathcal{K}}_0\}$ and, consequently, in problem (3.9) the cone \mathcal{K}_0 can be replaced by the cone $\bar{\mathcal{K}}_0$.

Note that the inclusions $\bar{\mathcal{K}}_0 \subset \mathcal{K}_0$ and $\mathcal{K}_0 \subset \mathcal{COP}^p$ imply $\bar{\mathcal{K}}_0 \subset \mathcal{COP}^p$.

To show that the regularizations presented above are themselves deeply connected, let us give an explicit description of the minimal face \mathbf{F}_{\min} in terms of the vertices of the set $\text{conv } T_{im}$ and the index sets $M(j)$, $j \in J$, defined as:

$$M(j) := \{k \in P : e_k^\top \mathcal{A}(x)t(j) = 0 \quad \forall x \in X\}, \quad j \in J. \tag{3.14}$$

The following theorem can be proved (see the results obtained in [12]).

Theorem 3.1. *Given the copositive problem (2.1), let $\{t(j), j \in J\}$ be the (finite) set of all vertices of the set $\text{conv } T_{im}$. Then the minimal face \mathbf{F}_{\min} of this problem can be described in two equivalent forms*

$$\begin{aligned} \mathbf{F}_{\min} &= K_{\min}(1) := \{D \in \mathcal{COP}^p : e_k^\top Dt(j) = 0 \quad \forall k \in M(j), \quad \forall j \in J\}, \text{ and} \\ \mathbf{F}_{\min} &= K_{\min}(2) := \{D \in \mathcal{COP}^p : e_k^\top Dt(j) = 0 \quad \forall k \in M(j), e_k^\top Dt(j) \geq 0 \quad \forall k \in P \setminus M(j), \quad \forall j \in J\}. \end{aligned}$$

Now, having described the minimal face \mathbf{F}_{\min} via immobile indices, we can compare the regularized problems (3.2), (3.9), and (3.13) in more detail.

The regularized problem (3.2) is formulated using the facial reduction approach to the copositive problem (2.1) and the regularized problems (3.9) and (3.13) are obtained using the immobile indices of this problem. The difference between these three problems is that in problem (3.2), the constraint set is determined by the minimal face \mathbf{F}_{\min} , while the constraints of problem (3.9) are formulated with the help of the cone \mathcal{K}_0 , and the constraints of problem (3.13) use the cone $\bar{\mathcal{K}}_0$.

It should be noticed that the minimal face \mathbf{F}_{\min} and the cones \mathcal{K}_0 and $\bar{\mathcal{K}}_0$ satisfy the inclusions

$$\mathbf{F}_{\min} \subset \bar{\mathcal{K}}_0 \subset \mathcal{K}_0.$$

At the same time, the cones \mathbf{F}_{\min} and $\bar{\mathcal{K}}_0$ are faces of the cone of copositive matrices \mathcal{COP}^p , while the cone \mathcal{K}_0 is generally not. In addition, one can show that $\bar{\mathcal{K}}_0$ is an exposed face while the face \mathbf{F}_{\min} as a whole is not.

For each of the conic problems mentioned above, we face certain challenges caused by the troubles associated with the *concrete construction* of the respective cones. For example, for the copositive problem (2.1), the following difficulties should be mentioned:

- to define the cones \mathcal{K}_0 and $\bar{\mathcal{K}}_0$, the elements $t(j)$, $j \in J$, of the finite set of indices (3.4) should be known;
- as far as we know, there are no explicit procedures for constructing the minimal face \mathbf{F}_{\min} and its dual cone \mathbf{F}_{\min}^* .

Theorem 3.1 shows how the minimal face \mathbf{F}_{\min} can be represented in the form of the cones $K_{\min}(1)$ and $K_{\min}(2)$ via immobile indices. Notice that to construct these cones, one has to find not only the set of indices (3.4), but also the corresponding sets $M(j)$, $j \in J$, defined in (3.14).

As mentioned above, regularity is an important property of optimization problems. As a rule, the regularity of copositive problems is characterized by the Slater condition. In this regard, it is important to note that the regularized problem (3.2) satisfies the *generalized* Slater condition while the regularized problems (3.9) and (3.13) obtained here satisfy the Slater type condition (3.5). This difference can be important for further study of linear CoP problems, as well as for the development of stable numerical methods for them.

4. ITERATIVE ALGORITHMS FOR REGULARIZATION OF LINEAR COPOSITVE PROBLEMS

In Section 3, we considered general schemes of two theoretical methods that made it possible to obtain regularizations of the linear copositive problem (2.1). In each of these schemes, we meet some difficulties associated with explicit representations of the respective “regularized” feasible cones and their dual ones. In this section, we consider and compare two different approaches to regularization aimed at overcoming these difficulties by using algorithmic procedures.

4.1. Waki and Muramatsu’s facial reduction algorithm

In [24] for linear conic problems, a regularization algorithm was proposed by Waki and Muramatsu. This algorithm can be thought of as the Facial Reduction Algorithm (FRA) from [5, 6], applied to linear conic problems in finite-dimensional spaces.

Let us describe the algorithm from [24] for the linear copositive problem (2.1) with the matrix constraint function $\mathcal{A}(x)$ defined in (2.2). Recall that here we suppose that problem (2.1) is feasible. Then, according to Remark 2.1, we can assume that $A_0 \in \mathcal{COP}^p$.

Denote

$$\text{Ker } \mathcal{A} := \{D \in \mathcal{S}(p) : A_j \bullet D = 0 \quad \forall j = 0, 1, \dots, n\}.$$

As above, let \mathcal{F}^* denote the dual cone of a given cone $\mathcal{F} \subset \mathcal{S}(p)$.

For a given feasible copositive problem (2.1), starting with \mathcal{COP}^p , the Waki and Muramatsu’s algorithm repeatedly finds smaller faces of \mathcal{COP}^p until it stops with the minimal face \mathbf{F}_{\min} .

Waki and Muramatsu’s FRA for the copositive problem (2.1)

Step 1. Set $i := 0$ and $\mathcal{F}_0 := \mathcal{COP}^p$.

Step 2. If $\text{Ker } \mathcal{A} \cap \mathcal{F}_i^* \subset \text{span}\{Y_1, \dots, Y_i\}$, then STOP: $\mathbf{F}_{\min} = \mathcal{F}_i$.

Step 3. Find $Y_{i+1} \in \text{Ker } \mathcal{A} \cap \mathcal{F}_i^* \setminus \text{span}\{Y_1, \dots, Y_i\}$.

Step 4. Set $\mathcal{F}_{i+1} := \mathcal{F}_i \cap \{Y_{i+1}\}^\perp$ and $i := i + 1$, and go to step 2.

The description of the algorithm is very simple but, in practice, its implementation presents serious difficulties which arise on step 2 and especially on step 3. As the matter of fact, in the case of the copositive problem (2.1), the fulfillment of step 3 is hard already on the first two iterations.

Let us consider the initial iteration when $i = 0$. On step 3, one has to find a matrix $Y_1 \in \text{Ker } \mathcal{A} \cap \mathcal{F}_0^*$. Since $\mathcal{F}_0 = \mathcal{COP}^p$, then at the current iteration ($i = 0$) we know the explicit description of the dual cone for \mathcal{F}_0 : $\mathcal{F}_0^* = \mathcal{CP}^p$, where the cone \mathcal{CP}^p is defined in (3.12). Therefore, the matrix Y_1 should have the form

$$Y_1 = \sum_{i \in I_1} t(i)(t(i))^\top, \quad t(i) \geq 0, \quad t(i) \neq 0 \quad \forall i \in I_1, \quad 0 < |I_1| \leq p(p+1)/2,$$

and the condition $\sum_{i \in I_1} (t(i))^\top A_j t(i) = 0 \quad \forall j = 0, 1, \dots, n$, has to be satisfied.

At the next iteration ($i = 1$), one is looking for a matrix Y_2 such that

C1: $Y_2 \in \mathcal{F}_1^* = \text{cl}\{D \in \mathcal{S}(p) : D \in \mathcal{CP}^p \oplus \alpha Y_1, \alpha \in \mathbb{R}\}$,

C2: $Y_2 \notin \text{span}\{Y_1\}$,

C3: $A_j \bullet Y_2 = 0 \quad \forall j = 0, 1, \dots, n$.

The first difficulty arises when trying to satisfy the condition **C1**, as there is no explicit description of the set \mathcal{F}_1^* . Notice that this set is defined using the closure operator, this operator being essential for the definition of \mathcal{F}_1^* . Therefore, in general, for a matrix Y_2 satisfying the condition **C1**, it may happen that $Y_2 \notin \{D \in \mathcal{S}(p) : D \in \mathcal{CP}^p \oplus \alpha Y_1, \alpha \in \mathbb{R}\}$.

In [24], there is also no any indication of how to find a matrix Y_2 satisfying the conditions **C2** and **C3**. Notice that the fulfillment of these conditions is a non-trivial task as well.

Thus, we can state that although the reported in [24] FRA is an easy-to-describe method, its practical implementation is not constructively described, which makes it difficult to apply. There is no information concerning which form should have the matrix Y_i at the i -th iteration ($i \geq 1$) of the algorithm and how to meet the conditions **C1–C3** for it.

4.2. A regularization based on the immobile indices

Here we will describe and justify a distinct algorithm for regularization of the copositive problem (2.1). This algorithm has a similar structure to the Waki and Muramatsu’s FRA considered in Section 4.1 but is based on

the concept of immobile indices and described in more detail, being, therefore, more constructive. Note from the outset that although our algorithm exploits the properties of the set of immobile indices, it does not require the initial knowledge of either this set or the vertices of its convex hull.

4.2.1. Algorithm **REG-LCoP** (*REGularization of Linear Copositive Problems*)

Consider the copositive problem in the form (2.1).

Iteration # 0. Using data of the original problem (2.1), let us form the following *regular* SIP problem:

$$\text{SIP}_0 : \min_{(x,\mu) \in \mathbb{R}^{n+1}} \mu, \text{ s.t. } t^\top \mathcal{A}(x)t + \mu \geq 0 \quad \forall t \in T,$$

with the index set T defined in (2.4).

If there exists a feasible solution $(\bar{x}, \bar{\mu})$ of this problem with $\bar{\mu} < 0$, then set $m_* := 0$ and go to the *Final step*.

Otherwise the vector $(x = \mathbf{0}, \mu = 0)$ is an optimal solution of the problem (SIP_0) .

It should be noticed that in the problem (SIP_0) , the index set T is compact and the constraints satisfy the Slater condition. Hence (see e.g. [4]), it follows from the optimality conditions for the vector $(x = \mathbf{0}, \mu = 0)$ that there exist indices and numbers

$$\tau(i) \in T, \quad \gamma(i) > 0 \quad \forall i \in I_1, \quad |I_1| \leq n + 1, \tag{4.1}$$

such that

$$\sum_{i \in I_1} \gamma(i) (\tau(i))^\top A_j \tau(i) = 0 \quad \forall j = 0, 1, \dots, n; \quad \sum_{i \in I_1} \gamma(i) = 1. \tag{4.2}$$

It follows from these relations (see Prop. 4.2 below) that $I_1 \neq \emptyset$, $\tau(i) \in T_{im} \subset T$, and $e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in P_+(\tau(i))$, $\forall i \in I_1, \quad \forall x \in X$. Set $L_1(i) := P_+(\tau(i))$, $i \in I_1$, and go to the next iteration.

Iteration # $m, m \geq 1$. By the beginning of the iteration, we have indices and sets $\tau(i), L_m(i), i \in I_m$, such that

$$\tau(i) \in T_{im}, \quad P_+(\tau(i)) \subset L_m(i) \subset P, \quad e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in L_m(i), \quad \forall i \in I_m, \quad \forall x \in X. \tag{4.3}$$

Consider a SIP problem

$$\begin{aligned} & \min_{(x,\mu) \in \mathbb{R}^{n+1}} \mu, \\ \text{SIP}_m : \text{ s.t. } & e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in L_m(i); \quad e_k^\top \mathcal{A}(x)\tau(i) \geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m, \\ & t^\top \mathcal{A}(x)t + \mu \geq 0 \quad \forall t \in \Omega(W_m), \end{aligned}$$

where $W_m := \{\tau(i), i \in I_m\}$ and the set $\Omega(W_m)$ is constructed by the rules (2.9), (2.10) with $V = W_m$.

In the problem (SIP_m) , the index set $\Omega(W_m)$ is compact and the constraints satisfy the following Slater type condition:

$$\begin{aligned} \exists (\hat{x}, \hat{\mu}) \text{ such that } & e_k^\top \mathcal{A}(\hat{x})\tau(i) = 0 \quad \forall k \in L_m(i); \quad e_k^\top \mathcal{A}(\hat{x})\tau(i) \geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m, \\ & t^\top \mathcal{A}(\hat{x})t + \hat{\mu} > 0 \quad \forall t \in \Omega(W_m). \end{aligned}$$

Hence, this problem is *regular*.

If problem (SIP_m) admits a feasible solution $(\bar{x}, \bar{\mu})$ with $\bar{\mu} < 0$, then STOP and go to the *Final step* with $m_* := m$.

Otherwise, the vector $(x = \mathbf{0}, \mu = 0)$ is an optimal solution of (SIP_m) . Since this problem is regular, the optimality of the vector $(x = \mathbf{0}, \mu = 0)$ provides (see [16]) that there exist indices, numbers, and vectors

$$\tau(i) \in \Omega(W_m), \quad \gamma(i), \quad i \in \Delta I_m, \quad 1 \leq |\Delta I_m| \leq n + 1; \quad \lambda^m(i) \in \mathbb{R}^p, \quad i \in I_m, \tag{4.4}$$

which satisfy the following conditions:

$$\sum_{i \in \Delta I_m} \gamma(i)(\tau(i))^\top A_j \tau(i) + 2 \sum_{i \in I_m} (\lambda^m(i))^\top A_j \tau(i) = 0 \quad \forall j = 0, 1, \dots, n; \tag{4.5}$$

$$\gamma(i) > 0 \quad \forall i \in \Delta I_m; \quad \lambda_k^m(i) \geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m. \tag{4.6}$$

Here and in what follows, without loss of generality, we suppose that $\Delta I_m \cap I_m = \emptyset$. Moreover, applying to the data (4.3), (4.4), the procedure **DAM** described in [13], it is possible to ensure that the following conditions are met:

$$P_0(\tau(i)) \cap P_+(\tau(j)) \neq \emptyset \quad \forall i \in \Delta I_m, \quad \forall j \in I_m. \tag{4.7}$$

Let us set

$$\begin{aligned} \Delta L(i) &:= \{k \in P \setminus L_m(i) : \lambda_k^m(i) > 0\}, & i \in I_m, \\ L_{m+1}(i) &:= L_m(i) \cup \Delta L(i), \quad i \in I_m; \quad L_{m+1}(i) := P_+(\tau(i)), \quad i \in \Delta I_m. \end{aligned} \tag{4.8}$$

It follows from (4.3) and (4.5) (see Prop. 4.3 below) that $e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in \Delta L(i), \quad \forall i \in I_m$, and $\tau(i) \in T_{im}, e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in P_+(\tau(i)), \quad \forall i \in \Delta I_m, \quad \forall x \in X$.

Go to the next iteration $\#(m + 1)$ with the new data

$$\tau(i), L_{m+1}(i), i \in I_{m+1} := I_m \cup \Delta I_m, \tag{4.9}$$

satisfying relations (4.3) with m replaced by $m + 1$.

Final step. It follows from Theorem 4.4 (see Sect. 4.2.2 below) that, in a finite number of iterations, the algorithm **REG-LCoP** comes to the final step. Therefore, for some $m_* \geq 0$, the problem (SIP $_{m_*}$) has a feasible solution $(\bar{x}, \bar{\mu})$ with $\bar{\mu} < 0$.

If $m_* = 0$, then the constraints of problem (2.1) satisfy the Slater condition with \bar{x} , and hence the problem is regular.

Suppose now that $m_* > 0$ and consider a problem

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & c^\top x, \\ \text{REG : s.t.} \quad & e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in L_{m_*}(i), \quad e_k^\top \mathcal{A}(x)\tau(i) \geq 0 \quad \forall k \in P \setminus L_{m_*}(i), \quad \forall i \in I_{m_*}, \\ & t^\top \mathcal{A}(x)t \geq 0 \quad \forall t \in \Omega(W_{m_*}), \end{aligned}$$

where the sets $W_{m_*} = \{\tau(i), i \in I_{m_*}\}$ and $\Omega(W_{m_*})$ are the same as in the problem (SIP $_{m_*}$). Problem (REG) has a finite number of linear equality/inequality constraints and according to Theorem 4.4 it possesses the following properties.

- (A) The set of feasible solutions X_{reg} of the problem (REG) coincides with the set of feasible solutions X of the original problem (2.1).
- (B) There exists $\bar{x} \in X$ such that $t^\top \mathcal{A}(\bar{x})t > 0 \quad \forall t \in \Omega(W_{m_*})$.

Hence the problem (REG) is equivalent to problem (2.1) and can be considered as its regularization. The algorithm is described.

Remark 4.1. In the described above algorithm **REG-LCoP**, it is assumed that $X \neq \emptyset$. It is easy to modify the algorithm so that this assumption can be removed.

4.2.2. Justification of Algorithm **REG-LCoP**

In this subsection, we prove some propositions and theorem that justify the algorithm **REG-LCoP**.

Proposition 4.2. *Let relations (4.1), (4.2) be satisfied. Then*

$$I_1 \neq \emptyset, \tau(i) \in T_{im}, e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in P_+(\tau(i)), \quad \forall i \in I_1, \quad \forall x \in X. \tag{4.10}$$

Proof. It is clear that the inequalities $\gamma(i) > 0, i \in I_1$, and the equality $\sum_{i \in I_1} \gamma(i) = 1$ imply that $I_1 \neq \emptyset$.

Consider the first group of equalities in (4.2). Add to the first equality of this group, corresponding to the $j = 0$, the remaining equalities, with $j = 1, \dots, n$, multiplied by x_j . As a result we get

$$\sum_{i \in I_1} \gamma(i) \left(\sum_{j=1}^n (\tau(i))^\top A_j \tau(i) x_j + (\tau(i))^\top A_0 \tau(i) \right) = 0 \quad \forall x \in \mathbb{R}^n$$

or equivalently

$$\sum_{i \in I_1} \gamma(i) (\tau(i))^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall x \in \mathbb{R}^n.$$

Taking into account the latest equalities, inequalities (2.13) and relations (4.1), we conclude that $(\tau(i))^\top \mathcal{A}(x) \tau(i) = 0$ for all $i \in I_1$ and $x \in X$. By definition, this means that $\tau(i) \in T_{im} \quad \forall i \in I_1$. Then it follows from Proposition 2.4 that for $i \in I_1$, the relations

$$e_k^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall k \in P_+(\tau(i)); \quad e_k^\top \mathcal{A}(x) \tau(i) \geq 0 \quad \forall k \in P_0(\tau(i)), \quad \forall x \in X, \tag{4.11}$$

hold true, wherefrom we conclude that relations (4.10) hold true too. The proposition is proved. □

Proposition 4.3. *Suppose that for $1 \leq m < m_*$, the indices and sets $\tau(i), L_m(i), i \in I_m$, satisfy (4.3) and the indices, numbers and vectors (4.4) satisfy (4.5), (4.6). Then*

$$\tau(i) \in T_{im}, e_k^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall k \in P_+(\tau(i)), \quad \forall i \in \Delta I_m, \quad \forall x \in X, \tag{4.12}$$

$$e_k^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall k \in \Delta L(i), \quad \forall i \in I_m, \quad \forall x \in X, \tag{4.13}$$

where the sets $\Delta L(i), i \in I_m$, are defined in (4.8).

Proof. Consider equalities (4.5). Add to the equality corresponding to $j = 0$ the remaining equalities corresponding to $j = 1, \dots, n$, multiplied by x_j . As a result, we get

$$\sum_{i \in \Delta I_m} \gamma(i) (\tau(i))^\top \left(\sum_{j=1}^n A_j x_j + A_0 \right) \tau(i) + 2 \sum_{i \in I_m} (\lambda^m(i))^\top \left(\sum_{j=1}^n A_j x_j + A_0 \right) \tau(i) = 0 \quad \forall x \in \mathbb{R}^n,$$

or equivalently

$$\sum_{i \in \Delta I_m} \gamma(i) (\tau(i))^\top \mathcal{A}(x) \tau(i) + 2 \sum_{i \in I_m} (\lambda^m(i))^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall x \in \mathbb{R}^n. \tag{4.14}$$

According to (4.3) we have that $\tau(i) \in T_{im}, i \in I_m$. Then it follows from Proposition 2.4 that for $i \in I_m$, relations (4.11) hold true. These relations and the equalities in (4.3) imply

$$e_k^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall k \in L_m(i); \quad e_k^\top \mathcal{A}(x) \tau(i) \geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m, \quad \forall x \in X. \tag{4.15}$$

Hence it follows from the relations above and (4.6), (2.13), (4.14) that for all $x \in X$

$$(\tau(i))^\top \mathcal{A}(x) \tau(i) = 0, \quad \forall i \in \Delta I_m; \quad \lambda_k^m(i) e_k^\top \mathcal{A}(x) \tau(i) = 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m,$$

wherefrom we obtain

$$\tau(i) \in T_{im} \quad \forall i \in \Delta I_m; \quad e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in \Delta L(i), \quad \forall i \in I_m, \quad \forall x \in X. \tag{4.16}$$

It follows from the inclusions $\tau(i) \in T_{im} \quad \forall i \in \Delta I_m$, and Proposition 2.4 that for all $i \in \Delta I_m$, relations (4.11) hold true. Taking into account these relations, and relations (4.16) we conclude that (4.12) and (4.13) hold true. The proposition is proved. \square

Theorem 4.4. *Given the CoP problem (2.1), in a finite number of iterations, the algorithm **REG-LCoP** constructs a problem (REG) possessing the properties (A) and (B).*

Proof. It follows from Propositions 4.2 and 4.3 that for any $m, 1 \leq m \leq m_* - 1$, at the beginning of the Iteration # m we have indices and sets $\tau(i), L_m(i), i \in I_m$, satisfying (4.3), and during this iteration the algorithm finds indices, numbers, and vectors (4.4) satisfying (4.5) and (4.6). Notice that by the definitions of the sets $\Omega(W_m), W_m := \{\tau(i), i \in I_m\}$ it holds:

$$\rho(t, \text{conv}W_m) \geq \sigma(W_m) > 0 \quad \forall t \in \Omega(W_m).$$

Hence the procedure **DAM** from [13] can be correctly applied to the data (4.3) and (4.5). This procedure consists of a finite number of operations and ensures the fulfilment of the conditions (4.7). It follows from these conditions that the algorithm **REG-LCoP** runs a finite number m_* of iterations and comes to final step with a vector $(\bar{x}, \bar{\mu}), \bar{\mu} < 0$, which is a feasible solution to the problem (SIP $_{m_*}$).

At the final step, the problem (REG) is formed on the basis of the problem (SIP $_{m_*}$). Let X_{reg} be the set of feasible solutions to the problem (REG):

$$X_{\text{reg}} := \{x \in \mathbb{R}^n : t^\top \mathcal{A}(x)t \geq 0 \quad \forall t \in \Omega(W_{m_*}); \\ e_k^\top \mathcal{A}(x)\tau(i) = 0 \quad \forall k \in L_{m_*}(i), \quad e_k^\top \mathcal{A}(x)\tau(i) \geq 0 \quad \forall k \in P \setminus L_{m_*}(i), \quad \forall i \in I_{m_*}\}$$

As before, let X be the set of feasible solutions of the original problem (2.1), and $\mathcal{X}(W_{m_*})$ the set defined by rules (2.9)–(2.11) with V replaced by W_{m_*} . For $m = m_*$, relations (4.15) imply that $X \subset X_{\text{reg}}$. On the other hand, it is clear that $X_{\text{reg}} \subset \mathcal{X}(W_{m_*})$. It follows from Proposition 4.3 that $W_{m_*} \subset T_{im}$, and consequently due to Theorem 2.3, we have $X = \mathcal{X}(W_{m_*})$. Hence we conclude that the problems (REG) and (2.1) have the same sets of feasible solutions: $X_{\text{reg}} = X$. The property (A) is proved.

By construction, the vector $(\bar{x}, \bar{\mu}), \bar{\mu} < 0$, is a feasible solution to the problem (SIP $_{m_*}$). Hence $\bar{x} \in X_{\text{reg}} = X$ and $t^\top \mathcal{A}(\bar{x})t \geq -\bar{\mu} > 0$ for all $t \in \Omega(W_{m_*})$. The property (B) is proved. \square

4.2.3. Example

Let us illustrate the iterations of the algorithm **REG-LCoP** with a small example.

Consider the CoP problem (2.1) with the following data:

$$n = 4, \quad p = 4, \quad c = (3, \quad 4, \quad -1, \quad -1)^\top, \\ A_0 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 4 & -1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \quad A_1 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 2 & 0 & 3 \\ 1 & 0 & -1 & 0 \\ 0 & 3 & 0 & 0 \end{pmatrix}, \\ A_2 = \begin{pmatrix} 0 & 1 & -2 & 0 \\ 1 & 4 & 1 & 0 \\ -2 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad A_4 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \tag{4.17}$$

Let us apply the algorithm **REG-LCoP** to this problem.

Iteration # 0. On this iteration we solve a semidefinite programming problem (SIP₀) with data (4.18). For this problem, relations (4.2) hold true with $I_1 = \{1\}$, $\tau(1) = e_1$, $\gamma_1 = 1$. These relations are the sufficient optimality conditions for the vector $(x = \mathbf{0}, \mu = 0)$ in this problem (SIP₀) and hence this vector is an optimal solution of the problem (SIP₀).

Set $L_1(1) := P_+(\tau(1)) = \{1\}$, $i \in I_1$, and go to the next iteration.

Iteration # 1. Using the indices and sets $\tau(i)$, $L_1(i)$, $i \in I_1$, found on the previous iteration, let us form the problem (SIP₁):

$$\begin{aligned} & \min \mu \\ & \text{s.t. } e_2^\top \mathcal{A}(x)\tau(1) = x_2 - x_3 + x_4 \geq 0, \quad e_3^\top \mathcal{A}(x)\tau(1) = x_1 - 2x_2 \geq 0, \\ & \quad t^\top \mathcal{A}(x)t + \mu \geq 0 \qquad \qquad \qquad \forall t \in \Omega(W_1), \end{aligned}$$

where $W_1 = \{\tau(1) = e_1\}$, $\Omega(W_1) := \{t \in T : \rho(t, \tau(1)) \geq \sigma(W_1) = 1\} = \{t \in \mathbb{R}_+^4 : e^\top t = 1, t_1 \leq 1/2\}$. Notice that in this example we have $e_1^\top \mathcal{A}(x)\tau(1) \equiv e_4^\top \mathcal{A}(x)\tau(1) \equiv 0$, and hence the constraints $e_1^\top \mathcal{A}(x)\tau(1) = 0$ and $e_4^\top \mathcal{A}(x)\tau(1) \geq 0$ are omitted in the problem (SIP₁).

The vector $(x = \mathbf{0}, \mu = 0)$ is an optimal solution to (SIP₁) since relations (4.5) hold true with $\Delta I_1 = \{2\}$, $\tau(2) = e_4 \in \Omega(W_1)$, $\gamma_2 = 1$, $\lambda^1(1) = \mathbf{0}$. Remind that these relations are the optimality conditions for $(x = \mathbf{0}, \mu = 0)$ in (SIP₁).

Hence we go to the next iteration with

$$\begin{aligned} I_2 &:= I_1 \cup \Delta I_1 = \{1, 2\}, \quad \tau(1) = e_1, \quad \tau(2) = e_4, \\ L_2(1) &:= L_1(1) \cup \Delta L_1(1) = L_1(1) = \{1\}, \quad L_2(2) := P_+(\tau(2)) = \{4\}. \end{aligned} \tag{4.18}$$

Iteration # 2. Using the known data (4.18), we form problem (SIP₂):

$$\begin{aligned} & \min \mu \\ & \text{s.t. } e_2^\top \mathcal{A}(x)\tau(1) = x_2 - x_3 + x_4 \geq 0, \quad e_3^\top \mathcal{A}(x)\tau(1) = x_1 - 2x_2 \geq 0, \\ & \quad e_2^\top \mathcal{A}(x)\tau(2) = 3x_1 \geq 0, \quad e_3^\top \mathcal{A}(x)\tau(2) = x_2 - 1 \geq 0, \\ & \quad t^\top \mathcal{A}(x)t + \mu \geq 0 \qquad \qquad \qquad \forall t \in \Omega(W_2), \end{aligned}$$

where $W_2 = \{\tau(1) = e_1, \tau(2) = e_4\}$, $\Omega(W_2) := \{t \in T : \rho(t, \text{conv}W_2) \geq \sigma(W_2) = 1\} = \{t \in \mathbb{R}_+^4 : e^\top t = 1, t_1 + t_4 \leq 1/2\}$. As before, since in this example, we have $e_1^\top \mathcal{A}(x)\tau(2) \equiv e_4^\top \mathcal{A}(x)\tau(2) \equiv 0$, the constraints $e_1^\top \mathcal{A}(x)\tau(2) \geq 0$ and $e_4^\top \mathcal{A}(x)\tau(2) = 0$ are not presented in our problem (SIP₂).

The problem (SIP₂) possesses a feasible solution $(\bar{x}, \bar{\mu})$ with $\bar{\mu} < 0$. For example, we can take $\bar{x} = (4, 1.5, 0.5, 1)^\top$, $\bar{\mu} = -0.4$. Hence, according to the algorithm, we should pass to the *Final step* with $m_* = 2$.

Final step. Consider a problem (REG) that is formed on the base of the problem (SIP₂), constructed at the last iteration with $m_* = 2$. In our example, the problem (REG) has the form

$$\begin{aligned} & \min c^\top x \\ & \text{s.t. } x_2 - x_3 + x_4 \geq 0, \quad x_1 - 2x_2 \geq 0, \quad 3x_1 \geq 0, \quad x_2 - 1 \geq 0, \\ & \quad t^\top \mathcal{A}(x)t \geq 0 \qquad \qquad \qquad \forall t \in \Omega(W_2), \end{aligned} \tag{4.19}$$

where, as before, $\Omega(W_2) = \{t \in \mathbb{R}_+^4 : e^\top t = 1, t_1 + t_4 \leq 1/2\}$.

As it was proved, this problem is equivalent to the original problem (2.1) with data (4.17) and possesses the properties (A) and (B). In particular, for this problem, there is a feasible solution $\bar{x} = (4, 1.5, 0.5, 1)^\top$, such that $t^\top \mathcal{A}(x)t \geq 0.4 \quad \forall t \in \Omega(W_2)$. Another useful property consists in the fact that in problem (4.19), the set of indices $\Omega(W_2)$ is smaller then the index set T in the original problem (2.1).

To illustrate the advantages of the regularized problem (REG), we solved this problem and the problem (2.1) with data (4.17) by a simple discretization on an uniform grid superimposed on the sets T and $\Omega(W_2)$, respectively. The auxiliary discretized linear programming (LP) problems were solved by a computer programme developed using the Matlab programming language, and all computations were performed with a personal computer. The accuracy of computations was 10^{-16} .

- By discretizing the regularized semi-infinite problem (4.19) using an uniform grid with a step $h = 0.1$ overlaid on the set $\Omega(W_2)$, we got an LP problem with **125** linear constraints. Having solved this problem (let us denote this problem by (LP1)), we obtained a solution $x^0 = (2.0000, 1.0000, 4.5000, 4.5000)$ and the optimal value of the problem $c^\top x^0 = 1$.

One can check that the found vector x^0 satisfies all the constraints of the problem (2.1) with the data (4.17), and hence is a feasible solution of this problem. Having verified the optimality conditions for linear copositive problems obtained in [14], one can conclude that x^0 is optimal in this problem.

- By discretization of the original problem (2.1) with the data (4.17) using the same uniform grid with the step $h = 0.1$ superimposed on the set T , we got an LP problem with **286** linear constraints. Denote this problem by (LP2). Having solved this problem, we obtained a solution $\tilde{x} = (0.7778, 0.5926, 2.0556, 2.4630)^\top$ and the corresponding value of the objective function $c^\top \tilde{x} = 0.1852$.

Recall that both discretized problems, (LP1) and (LP2), were obtained using the same grid's step $h = 0.1$. But in the case of the problem (LP2), due to the inclusion $\Omega(W_2) \subset T$, we have got more than twice as many constraints as in the LP problem (LP1).

Since we already know that the optimal value of the original problem (2.1) is equal to 1, but $c^\top \tilde{x} = 0.1852$, we can easily conclude that the found vector \tilde{x} does not belong to the feasible set of the original problem.

In order to get a more accurate solution of the original problem, we gradually reduced the step h of the grid.

For $h = 0.01$, we obtained an LP problem with **176 851** linear constraints whose optimal solution was $x^* = (1.8406, 0.9469, 4.1812, 4.2343)^\top$ and the optimal value $c^\top x^* = 0.8937$.

For $h = 0.0083$, we obtained an LP problem with **302 621** constraints, optimal solution $x^{**} = (1.8664, 0.9555, 4.2328, 4.2773)^\top$ and the optimal value $c^\top x^{**} = 0.9109$.

It is important to stress that all vectors \tilde{x} , x^* and x^{**} obtained by discretization of the original (not-regularized) problem are not feasible in this problem.

A further reducing the grid's step led to an increase in the number of constraints but not to an improvement in the quality of solution.

4.2.4. On the comparison of the algorithms

To give another interpretation of the algorithm **REG-LCoP** and to better trace the compliance of the algorithm **REG-LCoP** to the Waki and Muramatsu's FRA from [24] (presented here in Sect. 4.1), let us perform some additional constructions at the iterations of the algorithm **REG-LCoP**.

At the end of **Iteration # 0**, having data $\tau(i)$, $\gamma(i)$, $L_1(i)$, $i \in I_1$, let us set

$$\mathcal{F}_0 := \mathcal{COP}^p, \quad Y_1 := \sum_{i \in I_1} \gamma(i) \tau(i) (\tau(i))^\top, \quad \mathcal{F}_1 := \mathcal{F}_0 \cap \{Y_1\}^\perp.$$

Notice here that, by construction,

$$\begin{aligned} Y_1 &\neq \mathbb{O}_p, \quad Y_1 \in \text{Ker} A, \quad Y_1 \in \mathcal{F}_0^* = \mathcal{CP}^p, \\ \mathcal{F}_1 &:= \mathcal{F}_0 \cap \{Y_1\}^\perp = \{D \in \mathcal{COP}^p : D \bullet Y_1 = 0\} = \{D \in \mathcal{COP}^p : (\tau(i))^\top D \tau(i) = 0 \quad \forall i \in I_1\} \\ &= \{D \in \mathcal{COP}^p : e_k^\top D \tau(i) = 0 \quad \forall i \in L_1(i), \quad e_k^\top D \tau(i) \geq 0 \quad \forall i \in P \setminus L_1(i), \quad \forall i \in I_1\}, \end{aligned}$$

where \mathbb{O}_p is the $p \times p$ null matrix.

Consider **Iteration # m** , $1 \leq m \leq m_*$. By the beginning of the iteration, we have a cone $\mathcal{F}_m = \mathcal{F}_{m-1} \cap \{Y_m\}^\perp$ that can be described as follows:

$$\mathcal{F}_m = \{D \in \mathcal{CO}\mathcal{P}^p : e_k^\top D\tau(i) = 0 \quad \forall k \in L_m(i), e_k^\top D\tau(i) \geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m\}. \tag{4.20}$$

At the end of this iteration, we have new data (4.4). Let us set

$$Y_{m+1} := \sum_{i \in \Delta I_m} \gamma(i)\tau(i)(\tau(i))^\top + \sum_{i \in I_m} [\tau(i)(\lambda^m(i))^\top + \lambda^m(i)(\tau(i))^\top]. \tag{4.21}$$

From the equations in (4.5), we conclude: $Y_{m+1} \in \text{Ker}\mathcal{A}$. From (4.20), we get

$$\begin{aligned} \mathcal{F}_m^* &= \text{cl}\{D \in \mathcal{S}(p) : D \in \mathcal{CO}\mathcal{P}^p \oplus \mathcal{P}_m^*\}, \\ \mathcal{P}_m^* &:= \left\{ D \in \mathcal{S}(p) : D = \sum_{i \in I_m} [\tau(i)(\lambda(i))^\top + \lambda(i)(\tau(i))^\top], \quad \lambda_k(i) \geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m \right\}. \end{aligned}$$

Hence, by construction, $Y_{m+1} \in \mathcal{F}_m^*$.

Consider the cone $\mathcal{F}_{m+1} := \mathcal{F}_m \cap \{Y_{m+1}\}^\perp$ and show that it can be described as follows:

$$\mathcal{F}_{m+1} = \{D \in \mathcal{CO}\mathcal{P}^p : e_k^\top D\tau(i) = 0 \quad \forall k \in L_{m+1}(i), e_k^\top D\tau(i) \geq 0 \quad \forall k \in P \setminus L_{m+1}(i), \quad \forall i \in I_{m+1}\}. \tag{4.22}$$

In fact, it follows from (4.21) that the equality $D \bullet Y_{m+1} = 0$ can be rewritten in the form

$$\begin{aligned} 0 = D \bullet Y_{m+1} &= \sum_{i \in \Delta I_m} \gamma(i)(\tau(i))^\top D\tau(i) + 2 \sum_{i \in I_m} (\lambda^m(i))^\top D\tau(i), \\ \text{where } \gamma(i) > 0 \quad \forall i \in \Delta I_m; \quad \lambda_k^m(i) &\geq 0 \quad \forall k \in P \setminus L_m(i), \quad \forall i \in I_m. \end{aligned}$$

Taking into account (4.20) and the relations above, we conclude that for $D \in \mathcal{F}_m$, the equality $D \bullet Y_{m+1} = 0$ implies the equalities

$$(\tau(i))^\top D\tau(i) = 0 \quad \forall i \in \Delta I_m, e_k^\top D\tau(i) = 0 \quad \forall k \in \Delta L(i), \quad \forall i \in I_m.$$

Notice that the relations $D \in \mathcal{CO}\mathcal{P}^p$, $(\tau(i))^\top D\tau(i) = 0, \tau(i) \geq 0 \quad \forall i \in \Delta I_m$, imply

$$e_k^\top D\tau(i) = 0 \quad \forall k \in P_+(\tau(i)) \text{ and } e_k^\top D\tau(i) \geq 0 \quad \forall k \in P_0(\tau(i)), \quad \forall i \in \Delta I_m. \tag{4.23}$$

Representation (4.22) follows from (4.20) and (4.23).

The matrices and cones

$$Y_m, \mathcal{F}_m, m = 0, 1, \dots, m_*, \tag{4.24}$$

constructed by rules (4.21) and (4.22), satisfy the following relations:

$$Y_m \in \mathcal{F}_{m-1}^* \quad \forall m = 1, \dots, m_*, Y_0 = \mathbb{O}_p; \tag{I}$$

$$Y_m \in \text{Ker}\mathcal{A} \quad \forall m = 1, \dots, m_*; \tag{II}$$

$$\mathcal{F}_m = \mathcal{F}_{m-1} \cap \{Y_m\}^\perp \quad \forall m = 1, \dots, m_*, \mathcal{F}_0 = \mathcal{CO}\mathcal{P}^p. \tag{III}$$

Now we see that the algorithm **REG-LCoP** allows one to get a more clear description of the structure of the matrices $Y_m, m = 1, \dots, m_*$, satisfying conditions (I)–(III), and quite constructive rules of their formation:

- for a given m , the matrix Y_m has a form (4.21) and is built on the basis of the optimality conditions for the feasible solution ($x = \mathbf{0}, \mu = 0$) in the corresponding **regular SIP** problem (SIP_m).

As it was shown in Section 4.1, at each iteration, the Waki and Muramatsu’s FRA produces a set of matrices and cones (4.24) satisfying the conditions (I)–(III), and the condition

$$Y_m \notin \text{span}\{Y_0, Y_1, \dots, Y_{m-1}\} \quad \forall m = 1, \dots, m_* - 1. \tag{IV}$$

On the other hand, the algorithm **REG-LCoP** described in Section 4.2.1 at each iteration produces a set of matrices and cones (4.24) satisfying the conditions (I)–(III) but not necessarily the condition (IV).

Since in the algorithm **REG-LCoP**, the fulfillment of the condition (IV) is not guaranteed at each iteration, if compare this algorithm with the Waki and Muramatsu’s FRA, at the first glance it may seem that, in general, the number of iterations executed by the algorithm **REG-LCoP** is larger. Such an impression is caused by the fact that in Section 4.2.1, we described in more detail all the steps of the algorithm and explicitly indicated all the computations carried out at each iteration. As for the Waki and Muramatsu’s FRA, its iterations are described only in general terms.

In what follows, we set out a *modification* of the algorithm **REG-LCoP**, where the number of iterations is reduced and it is guaranteed that all conditions (I)–(IV) are satisfied on each *core* iteration. This modification is formal, being essentially another way of the iterations’ numbering. The real number of the calculations on the steps of this modified algorithm is the same as on the iterations of the original one.

4.2.5. *A compressed modification of the algorithm REG-LCoP*

Consider the algorithm **REG-LCoP** presented in Section 4.2.1. Evidently, one can reduce the number of iterations of the algorithm if squeeze into just one iteration that iterations of the algorithm which change the description of the dual cone \mathcal{F}_m^* , but do not change the cone \mathcal{F}_m itself. In other words, we will only move to the next *core* iteration when all conditions (I)–(IV) are satisfied. Formally, such a procedure can be described as follows.

Suppose that the algorithm **REG-LCoP** has constructed matrices and cones (4.24), satisfying the properties (I)–(III) and let $m_* > 0$. Denote by $m_s \in \{0, 1, \dots, m_* - 1\}$, $s = 0, 1, \dots, s_*$, the iterations’ numbers such that

$$\begin{aligned} m_0 &:= 0, \quad m_s < m_{s+1} && \forall s = 0, 1, \dots, s_* - 1; \quad m_{s_*} + 1 = m_*, \\ Y_{m_s+1} &\notin \text{span}\{Y_0, Y_1, \dots, Y_{m_s}\} && \forall s = 0, 1, \dots, s_* - 1; \\ Y_{m_s+1+i} &\in \text{span}\{Y_0, Y_1, \dots, Y_{m_s+i}\} && \forall i = 1, \dots, m_{s+1} - m_s - 1, \quad \forall s = 0, 1, \dots, s_* - 1. \end{aligned}$$

Here s_* denotes the number of iterations for which the conditions above are met. Notice that the set $\{l, l + 1, \dots, w\}$ is considered empty if $w < l$.

In other words, the condition (IV) is satisfied only for $m \in \{m_s + 1, s = 0, 1, \dots, s_* - 1\}$ and, possibly, for $m = m_*$. Set

$$\bar{Y}_0 := Y_0, \quad \bar{\mathcal{F}}_0 := \mathcal{F}_0, \quad \bar{Y}_{s+1} := Y_{m_s+1}, \quad \bar{\mathcal{F}}_{s+1} := \mathcal{F}_{m_s+1} \quad \forall s = 0, 1, \dots, s_*.$$

It is easy to check that the following conditions hold true:

$$\begin{aligned} \bar{Y}_s &\in \bar{\mathcal{F}}_{s-1}^*, \quad \bar{Y}_s \in \text{Ker } \mathcal{A}, \quad \bar{\mathcal{F}}_s = \bar{\mathcal{F}}_{s-1} \cap \{\bar{Y}_s\}^\perp && \forall s = 1, \dots, s_* + 1; \\ \bar{Y}_s &\notin \text{span}\{\bar{Y}_0, \bar{Y}_1, \dots, \bar{Y}_{s-1}\} && \forall s = 1, \dots, s_*. \end{aligned}$$

Thus, after the described above squeezing, we get s_* core iterations of the modified algorithm. It follows from the conditions above that $s_* \leq \dim(\text{Ker } \mathcal{A})$.

Notice that for any $s = 0, 1, \dots, s_* - 1$, the iterations of the algorithm **REG-LCoP** having the numbers $m_s + 1 + i$, where $i = 1, \dots, m_{s+1} - m_s - 1$ (the compressed iterations), are not useless. They can be considered as the steps of a regularization procedure for the cone \mathcal{F}_{m_s+1} at the current core iteration $\# s$. At each of these iterations, we reformulate the cone \mathcal{F}_{m_s+1} in a new equivalent form. This additional information allows us to improve (make more regular) the representation of the cone \mathcal{F}_{m_s+1} and get a more explicit and useful description of its dual cone $\mathcal{F}_{m_s+1}^*$.

4.2.6. A short discussion on the algorithms considered in this section

By analyzing and comparing the iterative algorithms presented above, we can draw the following conclusions.

- (1) The Waki and Muramatsu's facial reduction algorithm from [24], reformulated for copositive problems in Section 4.1, is very simple in the description and runs no more than $\dim(\text{Ker } \mathcal{A})$ iterations. But this algorithm is more conceptual than constructive since it does not provide any information about the structure of the matrix Y_m and the cone \mathcal{F}_m^* at its m -th iteration. Moreover, it is not explained in [24] how to fulfill steps 2 and 3 at each iteration.
- (2) The algorithm **REG-LCoP** proposed in Section 4.2.1 also runs a finite number of iterations. This algorithm is described in all details and justified. The quite constructive rules for calculating the matrix Y_m satisfying the condition $Y_m \in \mathcal{F}_m^*$, are presented using the information available at the Iteration $\#m$ of this algorithm. These rules are derived from the optimality conditions for the optimal solution ($x = \mathbf{0}, \mu = 0$) of the **regular** problem (SIP $_m$). Notice that it is possible to develop a modification of the algorithm **REG-LCoP** which runs no more than $2n$ iterations.
- (3) Finally, to show that the described in Section 4.2.1 algorithm **REG-LCoP** is not worse (by the number of iterations) than the FRA from the Section 4.1, we presented a compressed modification of the algorithm **REG-LCoP**. This modification consists of no more than $\dim(\text{Ker } \mathcal{A})$ iterations as well as the algorithm from Section 4.1.

5. CONCLUSIONS

The main contribution of the paper is that, based on the concept of immobile indices, previously introduced for semi-infinite optimization problems, we suggested new methods for regularization of copositive problems. The algorithmic procedure of regularization of copositive problems is described in the form of the algorithm **REG-LCoP** and is compared with the facial reduction approach based on the minimal cone representation. We show that, when applied to the linear CoP problem (2.1), the algorithm **REG-LCoP** possesses the same properties as the FRA suggested by Waki and Muramatsu in [24], but its iterations are explicit, described in more detail and hence more constructive.

The described in the paper algorithms are useful for the study of convex copositive problems. In particular, for the linear copositive problem, they allow to

- formulate an equivalent (regular) semi-infinite problem which satisfies the Slater type regularity condition and can be solved numerically;
- prove new optimality conditions without any CQs;
- develop strong duality theory based on an explicit representation of the “regularized” feasible cone and the corresponding dual (such as, *e.g.* the Extended Lagrange Dual Problem suggested for SDP by Ramana *et al.* [21]).

The described in the paper regularization approach is novel and rather constructive. It is important to stress that no constructive regularization procedures are known for linear copositive problems.

Acknowledgements. The authors thank the anonymous referees for their very helpful comments and suggestions which aided us in improving the presentation of this paper. This work was partially supported by the state research program “Convergence” (Republic Belarus), Task 1.3.01, by Portuguese funds through CIDMA – Center for Research and Development in Mathematics and Applications, and FCT – Portuguese Foundation for Science and Technology, within the project UIDB/04106/2020.

REFERENCES

- [1] F. Ahmed, M. Dür and G. Still, Copositive Programming via semi-infinite optimization. *J. Optim. Theory Appl.* **159** (2013) 322–340.
- [2] M.F. Anjos and J.B. Lasserre (editors), Handbook of Semidefinite, Conic and Polynomial Optimization. *International Series in Operational Research and Management Science*. Vol. 166. Springer US (2012) 138.
- [3] I.M. Bomze, Copositive optimization – recent developments and applications. *EJOR* **216** (2012) 509–520.

- [4] J.F. Bonnans and A. Shapiro, Perturbation analysis of optimization problems. Springer-Verlag, New-York, NY (2000) 601.
- [5] J.M. Borwein and H. Wolkowicz, Facial reduction for a cone-convex programming problem. *J. Austral. Math. Soc. Ser. A* **30** (1981) 369–380.
- [6] J.M. Borwein and H. Wolkowicz, Regularizing the abstract convex program. *J. Math. Anal. Appl.* **83** (1981) 495–530.
- [7] E. de Klerk and D.V. Pasechnik, Approximation of the stability number number of a graph via copositive programming. *SIAM J. Optim.* **12** (2002) 875–892.
- [8] D. Drusvyatskiy and H. Wolkowicz, The many faces of degeneracy in conic optimization. In: Foundations and Trends in Optimization. Vol. 3. Now Publishers Inc. (2017) 77–170.
- [9] M. Dür, Copositive programming – a survey. In: Recent Advances in Optimization and its Applications in Engineering, edited by M. Diehl, F. Glineur, E. Jarlebring and W. Michielis. Springer-Verlag, Berlin, Heidelberg (2010) 535.
- [10] K.O. Kortanek and Q. Zhang, Perfect duality in semi-infinite and semidefinite programming. *Math. Program. Ser. A* **91** (2001) 127–144.
- [11] O.I. Kostyukova and T.V. Tchemisova, Optimality conditions for convex semi-infinite programming problems with finitely representable compact index sets. *J. Optim. Theory Appl.* **175** (2017) 76–103.
- [12] O.I. Kostyukova and T.V. Tchemisova, On equivalent representations and properties of faces of the cone of copositive matrices. *Optimization* (2020). DOI: [10.1080/02331934.2022.2027939](https://doi.org/10.1080/02331934.2022.2027939).
- [13] O. Kostyukova and T. Tchemisova, On strong duality in linear copositive programming. *J. Global Optim.* (2021) 1–24. DOI: [10.1007/s10898-021-00995-3](https://doi.org/10.1007/s10898-021-00995-3).
- [14] O.I. Kostyukova, T.V. Tchemisova and O.S. Dudina, Immobile indices and CQ-free optimality criteria for linear copositive programming problems. *Set-Valued Var. Anal.* **28** (2020) 89–107.
- [15] A.N. Letchford and A.J. Parkes, A guide to conic optimisation and its applications. *RAIRO Oper. Res.* **52** (2018) 1087–1106.
- [16] V.L. Levin, Application of E. Helly’s theorem to convex programming, problems of best approximation and related questions. *Math. USSR Sbornik* **8** (1969) 235–247.
- [17] Z.-Q. Luo, F.J. Sturm and S. Zhang, Duality Results for Conic Convex Programming. Econometric institute report no. 9719/a. Erasmus University Rotterdam, Erasmus School of Economics (ESE), Econometric Institute (1997).
- [18] G. Pataki, A simple derivation of a facial reduction algorithm and extended dual systems. Preprint available at <http://www.unc.edu/pataki/papers/fr.pdf> (2000).
- [19] F. Permenter, H. Friberg and E. Andersen, Solving conic optimization problems via self-dual embedding and facial reduction: a unified approach. Tech. Report. (2015). DOI: [10.13140/RG.2.1.4340.7847](https://doi.org/10.13140/RG.2.1.4340.7847).
- [20] I. Pólik and T. Terlaky, Exact Duality for Optimization Over Symmetric Cones. AdvOL-Report No. 2007/10. McMaster University, Advanced Optimization Lab., Hamilton, Canada (2007).
- [21] M.V. Ramana, L. Tunçel and H. Wolkowicz, Strong duality for semidefinite programming. *SIAM J. Optim.* **7** (1997) 641–662.
- [22] M.V. Solodov, Constraint qualifications. In: Wiley Encyclopedia of Operations Research and Management Science, edited by J.J. Cochran, et al. John Wiley & Sons, Inc. (2010).
- [23] L. Tunçel and H. Wolkowicz, Strong duality and minimal representations for cone optimization. *Comput. Optim. Appl.* **53** (2013) 619–648.
- [24] H. Waki and M. Muramatsu, Facial reduction algorithms for conic optimization problems. *J. Optim. Theory Appl.* **158** (2013) 188–215.
- [25] G.-W. Weber, Generalized semi-infinite optimization and related topics. In: Research and Exposition in Mathematics, edited by K.H. Hofmann and R. Willem. Vol. 29. Heldermann Publishing House, Lemgo (2003).
- [26] H. Wolkowicz, R. Saigal and L. Vandenberghe, Handbook of Semidefinite Programming – Theory, Algorithms, and Applications. Kluwer Academic Publishers (2000).

Subscribe to Open (S2O)

A fair and sustainable open access model



This journal is currently published in open access under a Subscribe-to-Open model (S2O). S2O is a transformative model that aims to move subscription journals to open access. Open access is the free, immediate, online availability of research articles combined with the rights to use these articles fully in the digital environment. We are thankful to our subscribers and sponsors for making it possible to publish this journal in open access, free of charge for authors.

Please help to maintain this journal in open access!

Check that your library subscribes to the journal, or make a personal donation to the S2O programme, by contacting subscribers@edpsciences.org

More information, including a list of sponsors and a financial transparency report, available at: <https://www.edpsciences.org/en/maths-s2o-programme>