

# Accepted Manuscript

Bee pollen as a natural antioxidant source to prevent lipid oxidation in black pudding

Ofélia Anjos, Rodrigo Fernandes, Susana M. Cardoso, Teresa Delgado, Nelson Farinha, Vanessa Paula, Letícia M. Estevinho, Solange Teresinha Carpes



PII: S0023-6438(19)30536-5

DOI: <https://doi.org/10.1016/j.lwt.2019.05.105>

Reference: YFSTL 8207

To appear in: *LWT - Food Science and Technology*

Received Date: 23 February 2019

Revised Date: 20 May 2019

Accepted Date: 22 May 2019

Please cite this article as: Anjos, Ofé., Fernandes, R., Cardoso, S.M., Delgado, T., Farinha, N., Paula, V., Estevinho, Letí.M., Carpes, S.T., Bee pollen as a natural antioxidant source to prevent lipid oxidation in black pudding, *LWT - Food Science and Technology* (2019), doi: <https://doi.org/10.1016/j.lwt.2019.05.105>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Bee pollen as a natural antioxidant source to prevent lipid oxidation in black pudding**

Ofélia Anjos<sup>1,2,3</sup>, Rodrigo Fernandes<sup>1</sup>, Susana M. Cardoso<sup>4</sup>, Teresa Delgado<sup>3</sup>, Nelson Farinha<sup>3</sup>,  
Vanessa Paula<sup>6</sup>, Letícia M. Estevinho<sup>5</sup>, Solange Teresinha Carpes<sup>7</sup>

**Corresponding author** [Ofélia Anjos, [ofelia@ipcb.pt](mailto:ofelia@ipcb.pt)]

<sup>1</sup>Instituto Politécnico de Castelo Branco, Portugal

<sup>2</sup>Centro de Biotecnologia de Plantas da Beira Interior, Castelo Branco, Portugal

<sup>3</sup>Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Portugal

<sup>4</sup>QOPNA & LAQV-REQUIMTE, Department of Chemistry, University of Aveiro, Aveiro, Portugal

<sup>5</sup>Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Portugal

<sup>6</sup>School of Agricultural Sciences, Polytechnic Institute of Bragança, Bragança, Portugal

<sup>7</sup>Department of Chemistry, Federal University of Technology – Paraná (UTFPR), Pato Branco, PR, Brazil

**Abstract:** The antioxidant activity of bee pollen (mainly composed by *Cistus ladanifer pellets*) was explored in the context of black pudding production. For this purpose, three black pudding formulations comprising varying antioxidant compounds (sodium ascorbate, bee pollen and bee pollen extract) were produced.

Bee pollen was characterized according to the botanical origin, antioxidant activity, total phenol and flavonoid contents and phenolic profile. Black pudding was characterized by the microbiological safety, lipid oxidation, pH, water activity and humidity for 1, 10, 21, 30 and 37 days. Sensory acceptance was evaluated on the four first periods of storage. *Salmonella* spp., *Escherichia coli* and *Listeria monocytogenes* were absent in all samples. Small variations on humidity and pH were observed during the black pudding's storage. Regarding lipid oxidation, it increased, on average, from 1.36 mg to 2.11 mg malondialdehyde/kg meat. Differences among the three formulations were only significant on the first days of storage. The sensory assessment did not differ between products. This study suggests that bee pollen may be used as a natural antioxidant in meat products, yet a careful labelling is essential to alert allergic consumers.

**Key words:** black pudding, pollen, antioxidant, lipid oxidation

37

38 **Introduction**

39 The safety and quality of food products are some of the main concerns of health agencies and  
40 consumers worldwide. Also, the consumers are increasingly demanding for a diverse range of  
41 food options particularly those containing biologically active ingredients with health promoting  
42 capacities and free of food additives. However, for many food products, like those containing  
43 animal derivatives, the lipid oxidation is an important source of quality deterioration, reducing  
44 their shelf lifetime and impairing its consumption (Jayawardana et al., 2011; Morrissey,  
45 Sheehy, Galvin, Kerry, & Buckley, 1998; Shah, Bosco, & Mir, 2014).

46 Black pudding is a meatless sausage containing pork blood as a main ingredient and is a  
47 product of excellence in the traditional Portuguese charcuterie called “morcela de assar”. Blood  
48 sausages are produced and consumed throughout Europe and each Region reveal their own  
49 specificity and tradition. However, all black pudding are based on pork blood as its main raw  
50 material. In this case, the raw material is considered an important source of nutrients because  
51 meat derived products contain high amounts of proteins, vitamins (A, B12, and folic acid),  
52 essential minerals such as iron, zinc and selenium (Fellendorf, O’Sullivan, & Kerry, 2017). In  
53 addition, the blood also provides an important source of proteins and lysine (Fellendorf et al.,  
54 2017).

55 In the central region of Portugal, the black pudding is manufactured with pork fat, pork blood,  
56 bread, onion, coriander, sugar, olive oil and salt. The shelf-life of this product commonly ranges  
57 from 20 to 30 days, although for some specific formulations it can be increased to 90 days  
58 (Silva et al., 2014).

59 As far as the authors know the information available in the literature regarding the black  
60 pudding produced in Portugal is scarce. However, the physicochemical and sensory  
61 characterisation of *Morcilla de Burgos* a traditional Spanish blood sausage were studied by  
62 Santos et al (Santos, González-Fernández, Jaime, & Rovira, 2003). Ramos et al. (Ramos et al.,  
63 2013) provided an important study concerning the composition and quality of different blood

64 sausages from different countries with diverse raw materials and composition, while also  
65 plotting the importance of the mineral content in this kind of food product.

66 All blood sausages include antioxidants in its additive list, which allow to minimize  
67 lipid oxidation levels. However, both producers and consumers are looking for products where  
68 the synthetic antioxidants are replaced by natural ones derived from plants. In this sense, one  
69 must highlight bee pollen as a functional food product, since this is rich in proteins, lipids, free  
70 sugars, carbohydrates, minerals, phenolic acids, flavonoids, sterols, terpenoids, carotenoids and  
71 vitamins (Bogdanov, 2011). In fact bee pollen has gained widespread attention due it's  
72 purported antioxidant (Estevinho, Dias, & Anjos, 2019), anti-inflammatory (Maruyama,  
73 Sakamoto, Araki, & Hara, 2010), antimutagenic (Tohamy, Abdella, Ahmed, & Ahmed, 2014)  
74 and antimicrobial (Morais, Moreira, Feás, & Estevinho, 2011) properties. Indeed, new  
75 applications for bee pollen are currently being developed (Almeida et al., 2017; Krystyjan,  
76 Gumul, Ziobro, & Korus, 2015) mainly due to its use as a free radical scavenger and as lipid  
77 peroxidation inhibitor.

78 The aim of this study was to evaluate the shelf life of black pudding using pollen as  
79 natural antioxidant. Therefore, different formulations of black pudding with bee pollen, bee  
80 pollen extract and synthetic antioxidant were prepared to determine oxidative stability and  
81 sensory acceptability of the final product.

82

## 83 **1. Material and Methods**

### 84 **1.1. Chemicals**

85 Folin Ciocalteu phenol reagents, 2,2-diphenyl-1-picryl-hydrazyl (DPPH), gallic acid, 1,1,3,3  
86 tetramethoxypropane (TMP) and trichloroacetic acid were obtained from Sigma Aldrich  
87 (Sternheim, Germany). Aluminum chloride, sodium carbonate, sodium erythorbate (SE),  
88 potassium acetate, ethanol, ethylenediaminetetraacetic acid (EDTA), thiobarbituric acid (TBA)  
89 and chloroform were purchased from Sigma-Aldrich (Germany) and their purities were all over  
90 99%. Absolute alcohol was obtained from Sigma-Aldrich (Germany). All reagents used were of  
91 analytical grade.

## 92 **1.2. Bee pollen samples**

93 The bee pollen samples were collected directly from local beekeepers in the spring of 2017 in  
94 Castelo Branco, Portugal and stored frozen at -15 °C until further analysis.

95 The percentage of pollen grains belonging to each botanical family was determined based on the  
96 observation of 500 pollen grains in slides prepared according the acetolise method. The  
97 observation of pollens was carried out with a Leitz microscope (Leica, DML, Wetzlar,  
98 Germany) at x400 and an image analysis system Qwin 500 (Leica, England).

### 99 **1.2.1. Preparation of bee pollen extract**

100 For the bee pollen extraction 11 g of fresh bee pollen was stirred in a digital shaker (VWR  
101 15000-1 Advanced Orbital Digital Shaker) with 200 mL of 80% ethanol-water (v/v) at room  
102 temperature and at 4 x g during 24 hours in the dark. After this, samples were centrifuged at  
103 4080 x g, during 10 minutes and the supernatant was reserved. The extracts were evaporated at  
104 40 °C and then were frozen and lyophilized. After that the samples were stored at -20 °C until  
105 further analysis.

### 106 **1.2.2. Total phenolic and flavonoid compounds**

107 The total phenolic content (TPC) of the bee pollen extracts was determined using the Folin–  
108 Ciocalteu method as described by Moreira et al. (Moreira, Dias, Pereira, & Estevinho, 2008)  
109 and expressed as mg of gallic acid equivalents per g of bee pollen (GAE/g pollen).

110 For total flavonoids contents (TFC) determination in bee pollen the aluminium chloride method  
111 was used. Total flavonoids content was expressed as mg of quercetin equivalents per g of bee  
112 pollen (QE/g pollen) (Serra Bonvehí, Soliva Torrentó, & Centelles Lorente, 2001).

### 113 **1.2.3. Identification of the phenolic compounds in bee pollen**

114 The major phenolic compounds of the bee pollen extracts were identified by UHPLC-DAD-  
115 ESI-MS<sup>n</sup> analysis, using a Ultimate 3000 (Dionex Co., San Jose, CA, USA) apparatus with an  
116 ultimate 3000 Diode Array Detector (Dionex Co., San Jose, CA, USA) coupled to a Thermo  
117 LTQ XL (Thermo Scientific, San Jose, CA, USA) ion trap mass spectrometer equipped with an  
118 ESI source. The chromatographic column was an Hypersil Gold (Thermo Scientific, San Jose,  
119 CA, USA) C18 column (100 mm length; 2.1 mm i.d.; 1.9 µm particle diameter, end-capped) and

120 the general chromatographic conditions corresponded to those previous described (Wasli, Jelali,  
121 Silva, Ksouri, & Cardoso, 2018).

#### 122 **1.2.4. Antioxidant activity of the extracts**

123 In order to determine the antioxidant activity of the extract of bee pollen were tested two  
124 different methods, namely DPPH and reducing power assays.

#### 125 **Free-radical-scavenging (DPPH) assay**

126 The capacity to scavenge the 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical was monitored  
127 according to a method previously described by Morais et al (Morais et al., 2011).

128 The extract concentrations providing 50% scavenging ( $EC_{50}$ ) were calculated from the graph of  
129 scavenging effect percentage against extract concentration and the results were expressed as  
130 mg/mL.

#### 131 **Reducing power assay**

132 Reducing power in the extracts were determined by the procedure described by Berker et al  
133 (Berker, Güçlü, Tor, & Apak, 2007). In this procedure, the extract concentration that providing  
134 0.5 of absorbance ( $C_{0.5}$ ) was calculated from the graph of absorbance registered at 700 nm  
135 against the correspondent extract concentration and the results were expressed as mg GAE/mL.

#### 136 **1.3. Black pudding samples**

137 The black pudding preparation was conducted under formulation and traditional procedures  
138 used in the local factory named "Salsicharia Rebolosa". Regarding producer confidentiality  
139 issues the quantity of each ingredient is not publicised in this work, as well the quantity of  
140 pollen added.

141 The basic mixture of ingredients, without the commercial antioxidant, was divided into three  
142 lots. Several black puddings were performed containing three different antioxidant sources,  
143 namely: 1- fresh bee pollen; 2- lyophilized ethanolic extract of bee pollen; 3- sodium ascorbate  
144 (E301, a commercial antioxidant commonly used in the industrial process that in this study was  
145 used as control). The bee pollen (fresh and lyophilized ethanolic extract) was added dissolved in  
146 the volume of olive oil necessary for the black pudding preparation.

147 The black pudding for three treatments (different antioxidant sources) at five times (0, 10, 21,  
148 30 and 37 days) and in three replicates was made, totalizing 45 samples. All samples were  
149 divided in sealed polyethylene bags under vacuum and stored at 4 °C in a refrigerator. Because  
150 the legal shelf life of the black pudding is 30 days, and because is impossible to have the results  
151 of microbiological analysis at the same data of the sensory analysis, the evaluators do not made  
152 the sensory evaluation for the 37<sup>th</sup> days to avoid possible health risk.

### 153 **1.3.1. Microbiological analysis**

154 For the microbiological analysis of *L. monocytogenes*, 25 g of sample was homogenised for 2  
155 min in 225 mL of Half Fraser Base CM0895 (Oxoid, Hampshire, UK), using a Stomacher 400  
156 homogenizer (Seward, Basingstoke, England). The enumeration was performed according to the  
157 ISO 11290-2:1998/Amd. 1:2004(E) procedure (ISO, 1998). After incubation of the initial  
158 suspension for 1 h at 20 °C, a 0.1 mL volume was surface-inoculated on Oxoid Chromogenic  
159 Listeria Agar Base CM1084 (OCLA, Oxoid) and incubated at 37 °C for 48 h. The detection of  
160 *L. monocytogenes* was according to the ISO 11290-1:1996/Amd. 1:2004(E) procedure (ISO,  
161 1996). The initial suspension was supplemented with SR0166G selective supplement (Oxoid),  
162 incubated at 30 °C for 24 h. To the primary-enriched sample, 0.1 mL was streaked on OCLA  
163 and incubated at 37 °C for 48 h, for the secondary-enriched sample, 0.1 mL of the same initial  
164 supplemented suspension was transferred into 10 mL Fraser Broth supplemented with SR0156E  
165 (Oxoid), incubated at 37 °C for 48 h. If no growth was detected in primary-enriched sample, 0.1  
166 mL of the secondary-enriched sample was streaked on OCLA and incubated at 37 °C for 48 h.  
167 The colonies *L. monocytogenes* that grew on OCLA was green-blue surrounded by an opaque  
168 halo. The determinations per sample were carried out in duplicate and the results were  
169 expressed in CFU/g.

### 170 **1.3.2. Physicochemical analysis**

171 The samples were analysed for physicochemical composition (moisture, pH and water activity  
172 ( $a_w$ )) using standard procedures, along the storage time. Moisture content of samples, along the  
173 storage time, was quantified directly, according to the loss of mass after drying at 105 °C in an

174 oven (Thermo Scientific, Heratherm IMH 180) until constant weight, using AOAC procedures  
175 (AOAC, 1995). The results were expressed in percentage.

176 The pH of samples was determined weighing 10 g of black pudding and mixed with 100 mL of  
177 ultrapure water until a homogeny solution. The measurements were performed at room  
178 temperature (around 24 °C).

179 Water activity was determined by means of a Rotronic (HygroskopDT, Swiss) coupled with a  
180 Julabo (F35) thermostated Baths.

### 181 **1.3.3. Oxidative stability - thiobarbituric acid reactive substances content (TBARS)**

182 In order to determine the oxidative stability of the black pudding the method of thiobarbituric  
183 acid reactive substances (TBARS) was performed according Almeida et al. (Almeida et al.,  
184 2017). Measurements were made on the day of their production and over the storage time (1, 10,  
185 21, 30 and 37 days). Concentrations of 0.6 and 3.0 mmol/L of 1,1,3,3 tetramethoxypropane  
186 (TMP) were used as the standards. The results were expressed as mg of MDA/kg of sample  
187 (MDA: malondialdehyde). All measurement was carried out in triplicate.

### 188 **1.3.4. Sensory analysis**

189 The sensory acceptance test was performed using 32 untrained assessor's usual consumers of  
190 black pudding (14 women and 8 men with ages ranging between 23 and 54 years) performed the  
191 sensory evaluation.

192 The sensory analyses were performed at a room temperature and the samples were presented to  
193 the panel cut as 1 cm thick slices of roasted black pudding, under white natural lighting  
194 (according to the International Standards (ISO, 1988). Water and apple was provided for mouth  
195 rinsing between samples.

196 It was made a ranking descriptive analysis (RDA) (Richter, de Almeida, Prudencio, & de  
197 Toledo Benassi, 2010), in which the samples were presented at the same time to the panelists  
198 who had to rank the samples for the attribute aroma quality and the flavor, according to a proof  
199 sheet prepared for this specific purpose.

### 200 **1.4. Statistical analysis**



201 All tests were performed in triplicate and the results were presented as mean  $\pm$  standard  
202 deviation. A factorial variance analysis was performed to assess the effects of the different  
203 antioxidant used as well the shelf life period.

204 For each significant factor or interaction, the variance percentage was calculated and a Scheffé  
205 post-hoc test with 95% confidence was applied to the corresponding variables. For the statistical  
206 analysis of the sensory data resulting from the ranking test, the Friedman's test was performed  
207 based on the sum of the ordinations assigned by the tasters. All the calculations were performed  
208 using Statistica from Statsoft (vs 7.09) (Tulsa, OK, USA).

209

## 210 **2. Results and discussion**

### 211 **2.1. Bee pollen characterization**

212 It is well known that the chemical composition of bee pollen varies depending on the plant  
213 sources, growth conditions and storage conditions (Anjos, Paula, Delgado, & Estevinho, 2019;  
214 Atrouse, Oran, & Al-Abbadi, 2004; Bogdanov, 2011; Elamine et al., 2019; Letícia M.  
215 Estevinho, Dias, & Anjos, 2018; Leticia M. Estevinho, Rodrigues, Pereira, & Feás, 2012;  
216 Komosinska-Vassev, Olczyk, Kaźmierczak, Mencner, & Olczyk, 2015; Serra Bonvehí et al.,  
217 2001).

218 Palynological analysis found as predominant pollen *Cistus ladanifer* (42.6 %) followed by  
219 *Echium* spp. (13.6%) and *Apiaceae* (13.2%). 8.6 % of pollen of *Cistaceae* family were also  
220 founded. *Cistus ladanifer* pollen is very usual in Mediterranean regions, and in particular in  
221 the region of the study which was well characterized previously by Raimundo et al. (Raimundo  
222 et al., 2018).

223 The others pollen founded in the mixtures were: *Brassicaceae* spp. (10.1%); *Cichorieae* spp.  
224 (8.0%); *Asteraceae* spp. (1.9%); *Lavandula* spp. (1%); *Plantago* spp. (0.5%); *Silene* spp. (0.5%).

225 The values of TPC and TFC of bee pollen were  $35.05 \pm 0.5$  mg GAE/g of pollen and  $6.81 \pm 0.08$   
226 mg QE/g of pollen, respectively (Table 2). Our results showed a TPC higher than observed by  
227 Morais et al (Morais et al., 2011) that studied the honeybee-collected pollen from five  
228 Portuguese Natural Parks. They are also superior to those described by Campos et al. (Campos,

229 Webby, Markham, Mitchell, & da Cunha, 2003) who studied pollens from New Zealand and  
230 Portugal. Furthermore, the present values are comparable of TPC and TFC of bee pollen  
231 collected in Portugal with similar amount of *Cistus ladanifer* pollen (Anjos et al., 2019). The  
232 TPC of this pollen mixture were higher that the results founded for the pollen mixtures used by  
233 Almeida et al. (Almeida et al., 2017) that studied the use of lyophilized bee pollen extract as a  
234 natural antioxidant source in refrigerated sausages.

235 Because different antioxidant agents present different mechanism for their antioxidant capacities  
236 in this work was evaluated the antioxidant activity by two methods (DPPH and reducing power  
237 assay). On the other hand, and as say before, the antioxidant activity of bee pollen is well  
238 knowing as well the properties of *Cystus ladanifer* pollen. In this work the evaluation of this  
239 property is only to calculate the quantity of bee pollen that must be added in the black pudding  
240 formulation, in order to allow a similar antioxidant power of that of the commercial one.

241 The results of antioxidant activity of bee pollen assessed by free-radical-scavenging (DPPH)  
242 assay, expressed in terms of EC<sub>50</sub> value, and reducing power assay are summarized in Table 2.  
243 The EC<sub>50</sub> values of bee pollen sample is 2.62±0.09 mg/mL. This value indicates a good  
244 antioxidant activity and higher that the values reported by some authors (Negri, Barreto, Sper,  
245 Carvalho, & Campos, 2018; Suriyatem R., Auras R. A., Intipunya P., 2017)

246 Concerning the values obtained for reducing power assay they are also higher than those  
247 reported for Rape Bee Pollen (Sun, Guo, Zhang, & Zhuang, 2017).

248 The chromatographic profile at 280 nm of bee pollen extract is represented in Figure 1, while  
249 Table 1 summarizes the retention time, UV-vis and MSn spectral data of the identified  
250 compounds. Globally, the bee pollen extract was mainly rich in myricetin and quercetin O-  
251 derivatives (Table 1). Please note that the presence of flavonoids such as quercetin derivatives  
252 in the bee pollen has been previously related to the biological quality of the pollen, including its  
253 high antioxidant function (Lv, Wang, He, Wang, & Suo, 2015; Serra Bonvehí et al., 2001),  
254 which is one of the main claimed advantages to the use of bee pollen as an healthy product.

## 255 **2.2. Black pudding characterization**

256 In all samples, *Salmonella* spp., *Escherichia coli* and *Listeria monocytogenes* ATCC 19117.  
257 were analysed, according the Portuguese legislation and were absent in all of them.

258 The results obtained for the pH, moisture content, water activity and lipid oxidation by TBARS  
259 analysis during the storage period of black pudding are presented in Table 3. Overall, the pH of  
260 black pudding samples were similar to those studied by Santos et al. (Santos et al., 2003) and  
261 higher to those founded by Diez et al. (Diez, Santos, Jaime, & Rovira, 2008) that studied blood  
262 sausages produced with rice. The pH of the different formulations of black puddings was  
263 influenced by the antioxidant added and the storage period (Table 3). The higher values are  
264 founded for the black puddings produced with bee pollen as antioxidant, and the variations  
265 during the time is different for the different formulation ( $TxD = 44.0^{***}$ , Table 4). These  
266 variations could be explained by the fact that the bee pollen have a lower pH than black  
267 pudding. According to Anjos et al (Anjos et al., 2019) the pH of pollen ranging between 3.4 and  
268 5.9. The pH of the pollen used in the present study is  $4.70 \pm 0.47$ .

269 Moisture content of black pudding depends on the fat content and the final preservation process:  
270 cooked, drying or smoking (Ramos et al., 2013). The moisture content of studied products  
271 ranging, on average, between 46.33% and 49.73%. The different treatments and storage days  
272 were significant factors explaining 48% and 7% of the total variance, respectively, regardless  
273 variability between samples also had a high impact (explaining 11.6% of the total variance). The  
274 lower moisture content is observed for the black pudding made with pollen and the higher  
275 values for the black pudding made with pollen extract. Our values are lower than observed by  
276 Fellendorf et al (Fellendorf et al., 2017) that studied black puddings usually consumed in  
277 Ireland and the United Kingdom. Differences can however be due to the distinct list of  
278 ingredients among the formulations.

279  $a_w$  is a feature of great importance in food products preservation and particularly in black  
280 pudding that was produced with meat and blood. During the manufacturing process of these  
281 products they were subjected to high temperatures and, as expected, vegetative cells do not  
282 survive, but after the high temperatures process, post-contamination of the product may occur.  
283 The higher values of  $a_w$  for these kind of products are always higher (Santos et al., 2003) and

284 because of that it is very important to performed a restrict quality control. Our values for  $a_w$   
285 ranging between 0.90 and 0.92 was lower than observed in other studies (Santos et al., 2003).  
286 For this parameter, no significant differences were found among the sausages formulations  
287 neither along the storage period (Table 4).

288 TBARS is generally used as an indicator of the degree of lipid oxidation for pork meat and pork  
289 meat sausages, that reflects the content of MDA formed during the oxidation of polyunsaturated  
290 fatty acids (Tang, Sheehan, Buckley, Morrissey, & Kerry, 2001).

291 Concerning the TBARS, Selani et al. (Selani et al., 2011) refer that values lower than 3 mg of  
292 MDA/kg sample can be considered in good condition. All the black pudding samples analysed  
293 could be considered in good condition during all storage periods (values lower 2.56 mg  
294 MDA/kg sample) (Table 3). In the first day, the sample prepared with pollen extract had a value  
295 of 3.04 mg MDA/kg of black pudding. The black pudding prepared with pollen as natural  
296 antioxidant had values similar to those observed for the black pudding prepared with the  
297 commercial antioxidants, except for the first 15 days after production. Further studies may be  
298 performed in order to evaluate the optimum quantity of pollen.

299 For TBARS all factors are highly significant, but the stored days are the most important and  
300 explain 44.7 % of the total variance. The variation between days is also different for the  
301 different antioxidants used. The bee pollen presents a similar antioxidant effect than the  
302 commercial product.

### 303 **2.3. Sensory evaluation**

304 The sensory evaluation of the 12 black pudding products is plot in Figure 2.

305 The black pudding was sensory analysed only until 30 days, because is the legal self-life in the  
306 factory. According Silva et al (Silva et al., 2014) after 30 days of storage, the over-wrap packed  
307 blood sausages present mould and yeast. Nevertheless, for the vacuum-packed blood sausages  
308 the mould and yeast appears only after 45 days (Silva et al., 2014). In our study we use the  
309 vacuum-packed system but because no studies were performed in this product to extend the  
310 self-life we consider only the legal limit (30 days) established for this product for sensory  
311 analysis, excluding for this propose the samples with 37 day of self-life. In fact, the

312 microbiological results confirm that no mould or yeast have been developed in the samples, so it  
313 was need future research in other to establish better the shelf-life, is not an aim of this work.

314 The ANOVA made for all samples and for the appearance and flavor revel that no significant  
315 difference exists for the storage period (appearance:  $p=1.000$ ; flavor:  $p=0.999$ ) and for the  
316 different formulation period (appearance:  $p=0.328$ ; flavor:  $p=0.235$ ). These results confirm that  
317 the new additives, pollen or pollen extract do not affect the preference of the consumers.

318 Many of the tasters referred that the forced choice required by the triangular test was very  
319 difficult because the tree samples are very similar (24 % of the tasters). Other comments given  
320 by the tasters that help they to identify some differences are: homogeneity of the product (18%);  
321 visible pieces of onion (5%) and more fat quantity in a specific sample (3%). However, this  
322 kinds of comments are all related to manufacture process of this product. The different raw  
323 materials are cut in small pouches and mixed but not crushed.

324

### 325 **3. Conclusion**

326 The inclusion of bee pollen as an antioxidant could be a natural alternative to prevent the lipid  
327 oxidation in black pudding. These products could be added dissolved in the olive oil that will be  
328 used in the preparation of the sausage and have the advantage to be a recognized healthy food  
329 product.

330 Additionally, the use of bee pollen as antioxidant improve product quality and consumer  
331 acceptance and do not affect their traditional flavor.

332 Furthermore, it is important to note that the use of bee pollen must be very well mentioned in  
333 the label, to prevent allergic risks. More studies will be need in order to identify the more  
334 appropriate concentration of bee pollen to use as well the influence of botanical origin of bee  
335 pollen.

336

### 337 **Acknowledgments**

338 Centro de Estudos Florestais that is a research unit funded by FCT (UID/AGR/00239/2019).

339 The author Estevinho M. L. wishes to thank to the strategic programme  
340 UID/BIA/04050/2013(POCI-01-0145-FEDER-007569) funded by national funds through the  
341 Fundação para a Ciência e a Tecnologia (FCT, Portugal) and by the European Regional  
342 Development Fund (ERDF) through the COMPETE2020-Programa Operacional  
343 Competitividade e Internacionalização (POCI).

344 Thanks are also due to FCT/MEC for the financial support to the research unit QOPNA (FCT  
345 UID/QUI/00062/2019), through national funds and where applicable co-financed by FEDER,  
346 within PT2020 Partnership Agreement. Susana Cardoso thanks the research contract under the  
347 project AgroForWealth (CENTRO-01-0145-FEDER-000001), funded by Centro2020, through  
348 FEDER and PT2020.

349

350 **Declarations of interest:** none

## 351 **References**

- 352 Almeida, J. de F., Reis, A. S. dos, Heldt, L. F. S., Pereira, D., Bianchin, M., Moura, C. de, ...  
353 Carpes, S. T. (2017). Lyophilized bee pollen extract: A natural antioxidant source to  
354 prevent lipid oxidation in refrigerated sausages. *LWT - Food Science and Technology*, *76*,  
355 299–305. <http://doi.org/10.1016/j.lwt.2016.06.017>
- 356 Anjos, O., Paula, V., Delgado, T., & Estevinho, L. (2019). Influence of the storage conditions  
357 on the quality of bee pollen. *Zemdirbyste-Agriculture*, *106*(1), 87–94.  
358 <http://doi.org/10.13080/z-a.2019.106.012>
- 359 AOAC. Official Methods of Analysis, 16th edn. Arlington, VA, USA: Association of Official  
360 Analytical Chemists (1995).
- 361 Atrouse, O. M., Oran, S. A., & Al-Abadi, S. Y. (2004). Chemical analysis and identification of  
362 pollen grains from different jordanian honey samples. *International Journal of Food  
363 Science and Technology*, *39*(4), 413–417. [http://doi.org/10.1111/j.1365-  
364 2621.2004.00798.x](http://doi.org/10.1111/j.1365-2621.2004.00798.x)
- 365 Berker, K. I., Güçlü, K., Tor, İ., & Apak, R. (2007). Comparative evaluation of Fe(III) reducing  
366 power-based antioxidant capacity assays in the presence of phenanthroline, batho-  
367 phenanthroline, tripyridyltriazine (FRAP), and ferricyanide reagents. *Talanta*, *72*(3),  
368 1157–1165. <http://doi.org/10.1016/j.talanta.2007.01.019>
- 369 Bogdanov, S. (2011). Pollen: Nutrition, Functional Properties, Health: A Review. In B. P.  
370 Science (Ed.), *Bee Product Science*. Retrieved from <http://www.bee->

- 371 hexagon.net/files/file/fileE/Health/PollenBook2Review.pdf
- 372 Campos, M. G., Webby, R. F., Markham, K. R., Mitchell, K. A., & da Cunha, A. P. (2003).  
373 Age-Induced Diminution of Free Radical Scavenging Capacity in Bee Pollens and the  
374 Contribution of Constituent Flavonoids. *Journal of Agricultural and Food Chemistry*,  
375 *51*(3), 742–745. <http://doi.org/10.1021/jf0206466>
- 376 Diez, A. M., Santos, E. M., Jaime, I., & Rovira, J. (2008). Application of organic acid salts and  
377 high-pressure treatments to improve the preservation of blood sausage. *Food*  
378 *Microbiology*, *25*(1), 154–161. <http://doi.org/10.1016/j.fm.2007.06.004>
- 379 Elamine, Y., Lyoussi, B., Anjos, O., Estevinho, L. M., Aazza, S., Carlier, J. D., ... Miguel, M.  
380 G. (2019). Zantaz honey “monoflorality”: Chemometric applied to the routinely assessed  
381 parameters. *LWT*. <http://doi.org/10.1016/j.lwt.2019.02.039>
- 382 Estevinho, L. M., Dias, T., & Anjos, O. (2018). Influence of the storage conditions (frozen vs  
383 dried) in health-related lipid indexes and antioxidants of bee pollen. *European Journal of*  
384 *Lipid Science and Technology*, 1800393. <http://doi.org/10.1002/ejlt.201800393>
- 385 Estevinho, L. M., Rodrigues, S., Pereira, A. P., & Feás, X. (2012). Portuguese bee pollen:  
386 palynological study, nutritional and microbiological evaluation. *International Journal of*  
387 *Food Science & Technology*, *47*(2), 429–435. [http://doi.org/10.1111/j.1365-](http://doi.org/10.1111/j.1365-2621.2011.02859.x)  
388 [2621.2011.02859.x](http://doi.org/10.1111/j.1365-2621.2011.02859.x)
- 389 Fellendorf, S., O’Sullivan, M. G., & Kerry, J. P. (2017). Effect of different salt and fat levels on  
390 the physicochemical properties and sensory quality of black pudding. *Food Science &*  
391 *Nutrition*, *5*(2), 273–284. <http://doi.org/10.1002/fsn3.390>
- 392 ISO. Sensory analysis. General guidance for the design of test rooms. Ref. no. International  
393 Organization for Standardization, Genève, Switzerland, ISO 8589:1988. (1988).
- 394 ISO. (1996). 11290-1:1996/Amd 1:2004. Modification of the isolation media and the  
395 haemolysis test, and inclusion of precision data. *Modification of the Isolation Media and*  
396 *the Haemolysis Test, and Inclusion of Precision Data*.
- 397 Jayawardana, B. C., Hirano, T., Han, K. H., Ishii, H., Okada, T., Shibayama, S., ... Shimada, K.  
398 I. (2011). Utilization of adzuki bean extract as a natural antioxidant in cured and uncured  
399 cooked pork sausages. *Meat Science*, *89*(2), 150–153.  
400 <http://doi.org/10.1016/j.meatsci.2011.04.005>
- 401 Komosinska-Vassev, K., Olczyk, P., Kaźmierczak, J., Mencner, L., & Olczyk, K. (2015). Bee  
402 Pollen: Chemical Composition and Therapeutic Application. *Evidence-Based*  
403 *Complementary and Alternative Medicine*, *2015*, 1–6. <http://doi.org/10.1155/2015/297425>
- 404 Krystijan, M., Gumul, D., Ziobro, R., & Korus, A. (2015). The fortification of biscuits with bee  
405 pollen and its effect on physicochemical and antioxidant properties in biscuits. *LWT -*  
406 *Food Science and Technology*, *63*(1), 640–646. <http://doi.org/10.1016/j.lwt.2015.03.075>
- 407 Lv, H., Wang, X., He, Y., Wang, H., & Suo, Y. (2015). Identification and quantification of

- 408 flavonoid aglycones in rape bee pollen from Qinghai-Tibetan Plateau by HPLC-DAD-  
409 APCI/MS. *Journal of Food Composition and Analysis*, 38, 49–54.  
410 <http://doi.org/10.1016/j.jfca.2014.10.011>
- 411 Maruyama, H., Sakamoto, T., Araki, Y., & Hara, H. (2010). Anti-inflammatory effect of bee  
412 pollen ethanol extract from *Cistus* sp. of Spanish on carrageenan-induced rat hind paw  
413 edema. *BMC Complementary and Alternative Medicine*, 10(1), 30.  
414 <http://doi.org/10.1186/1472-6882-10-30>
- 415 Morais, M., Moreira, L., Feás, X., & Estevinho, L. M. (2011). Honeybee-collected pollen from  
416 five Portuguese Natural Parks: Palynological origin, phenolic content, antioxidant  
417 properties and antimicrobial activity. *Food and Chemical Toxicology*, 49(5), 1096–1101.  
418 <http://doi.org/10.1016/j.fct.2011.01.020>
- 419 Moreira, L., Dias, L. G., Pereira, J. A., & Estevinho, L. (2008). Antioxidant properties, total  
420 phenols and pollen analysis of propolis samples from Portugal. *Food and Chemical*  
421 *Toxicology*, 46(11), 3482–3485. <http://doi.org/10.1016/j.fct.2008.08.025>
- 422 Morrissey, P. A., Sheehy, P. J. A., Galvin, K., Kerry, J. P., & Buckley, D. J. (1998). Lipid  
423 stability in meat and meat products. *Meat Science*, 49, S73–S86.  
424 [http://doi.org/10.1016/S0309-1740\(98\)90039-0](http://doi.org/10.1016/S0309-1740(98)90039-0)
- 425 Negri, G., Barreto, L. M. R. C., Sper, F. L., Carvalho, C. de, & Campos, M. das G. R. (2018).  
426 Phytochemical analysis and botanical origin of *Apis mellifera* bee pollen from the  
427 municipality of Canavieiras, Bahia State, Brazil. *Brazilian Journal of Food Technology*,  
428 21(2), e57566. <http://doi.org/10.1590/1981-6723.17616>
- 429 Raimundo, J. R., Frazão, D. F., Domingues, J. L., Quintela-Sabarís, C., Dentinho, T. P., Anjos,  
430 O., ... Delgado, F. (2018). Neglected Mediterranean plant species are valuable resources:  
431 the example of *Cistus ladanifer*. *Planta*, 248(6), 1351–1364.  
432 <http://doi.org/10.1007/s00425-018-2997-4>
- 433 Ramos, D. D., Villalobos-delgado, L. H., Cabeza, E. A., Caro, I., Fernández-diez, A., & Mateo,  
434 J. (2013). Mineral Composition of Blood Sausages – A Two-Case Study, (January), 93–  
435 111. <http://doi.org/10.5772/53591>
- 436 Richter, V. B., de Almeida, T. C. A., Prudencio, S. H., & de Toledo Benassi, M. (2010).  
437 Proposing a ranking descriptive sensory method. *Food Quality and Preference*, 21(6),  
438 611–620. <http://doi.org/10.1016/j.foodqual.2010.03.011>
- 439 Santos, E. M., González-Fernández, C., Jaime, I., & Rovira, J. (2003). Physicochemical and  
440 sensory characterisation of Morcilla de Burgos, a traditional Spanish blood sausage. *Meat*  
441 *Science*, 65(2), 893–898. [http://doi.org/10.1016/S0309-1740\(02\)00296-6](http://doi.org/10.1016/S0309-1740(02)00296-6)
- 442 Selani, M. M., Contreras-Castillo, C. J., Shirahigue, L. D., Gallo, C. R., Plata-Oviedo, M., &  
443 Montes-Villanueva, N. D. (2011). Wine industry residues extracts as natural antioxidants  
444 in raw and cooked chicken meat during frozen storage. *Meat Science*, 88(3), 397–403.



- 445 <http://doi.org/10.1016/j.meatsci.2011.01.017>
- 446 Serra Bonvehí, J., Soliva Torrentó, M., & Centelles Lorente, E. (2001). Evaluation of  
447 polyphenolic and flavonoid compounds in honeybee-collected pollen produced in Spain.  
448 *Journal of Agricultural and Food Chemistry*, 49(4), 1848–53. Retrieved from  
449 <http://www.ncbi.nlm.nih.gov/pubmed/11308335>
- 450 Shah, M. A., Bosco, S. J. D., & Mir, S. A. (2014). Plant extracts as natural antioxidants in meat  
451 and meat products. *Meat Science*, 98(1), 21–33.  
452 <http://doi.org/10.1016/j.meatsci.2014.03.020>
- 453 Silva, F. A. P., Amaral, D. S., Guerra, I. C. D., Arcanjo, N. M. O., Bezerra, T. K. A., Ferreira,  
454 V. C. S., ... Madruga, M. S. (2014). Shelf life of cooked goat blood sausage prepared with  
455 the addition of heart and kidney. *Meat Science*, 97(4), 529–533.  
456 <http://doi.org/10.1016/j.meatsci.2014.03.018>
- 457 Silva, F. A. P., Amaral, D. S., Guerra, I. C. D., Dalmás, P. S., Arcanjo, N. M. O., Bezerra, T. K.  
458 A., ... Madruga, M. S. (2013). The chemical and sensory qualities of smoked blood  
459 sausage made with the edible by-products of goat slaughter. *Meat Science*, 94(1), 34–38.  
460 <http://doi.org/10.1016/j.meatsci.2013.01.004>
- 461 Sun, L., Guo, Y., Zhang, Y., & Zhuang, Y. (2017). Antioxidant and Anti-tyrosinase Activities  
462 of Phenolic Extracts from Rape Bee Pollen and Inhibitory Melanogenesis by  
463 cAMP/MITF/TYR Pathway in B16 Mouse Melanoma Cells. *Frontiers in Pharmacology*,  
464 8. <http://doi.org/10.3389/fphar.2017.00104>
- 465 Suriyatem R., Auras R. A., Intipunya P., R. P. (2017). Predictive mathematical modeling for  
466 EC50 calculation of antioxidant activity and antibacterial ability of Thai bee products.  
467 *Journal of Applied Pharmaceutical Science*. <http://doi.org/10.7324/JAPS.2017.70917>
- 468 Tang, S., Sheehan, D., Buckley, D. J., Morrissey, P. A., & Kerry, J. P. (2001). Anti-oxidant  
469 activity of added tea catechins on lipid oxidation of raw minced red meat, poultry and fish  
470 muscle. *International Journal of Food Science and Technology*, 36(6), 685–692.  
471 <http://doi.org/10.1046/j.1365-2621.2001.00497.x>
- 472 Tohamy, A. A., Abdella, E. M., Ahmed, R. R., & Ahmed, Y. K. (2014). Assessment of anti-  
473 mutagenic, anti-histopathologic and antioxidant capacities of Egyptian bee pollen and  
474 propolis extracts. *Cytotechnology*, 66(2), 283–297. <http://doi.org/10.1007/s10616-013-9568-0>
- 475
- 476 Wasli, H., Jelali, N., Silva, A. M. S., Ksouri, R., & Cardoso, S. M. (2018). Variation of  
477 polyphenolic composition, antioxidants and physiological characteristics of dill (*Anethum*  
478 *graveolens* L.) as affected by bicarbonate-induced iron deficiency conditions. *Industrial*  
479 *Crops and Products*, 126, 466–476. <http://doi.org/10.1016/j.indcrop.2018.10.007>
- 480
- 481

482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492

**Table 2.** TPC, TFC and antioxidant activity of bee pollen extract.

Parameters	Bee pollen
TPC (mg GAE/g of pollen)	35.05±0.5
TFC (mg QE/g pollen)	6.99±0.33
EC <sub>50</sub> (mg/mL)	2.62±0.09
Reducing power assay (mg GAE/mL)	6.51±0.30

493  
494  
495

**Table 3.** Chemical analysis of different formulation of black pudding.

	Days	Treatment		
		E301	Pollen	Pollen extract
pH	1	6.75±0.02 <sup>bB</sup>	6.70±0.01 <sup>aA</sup>	6.68±0.01 <sup>aA</sup>
	10	6.72±0.01 <sup>bB</sup>	6.75±0.01 <sup>bC</sup>	6.72±0.00 <sup>aA</sup>
	21	6.60±0.02 <sup>cA</sup>	6.80±0.01 <sup>cB</sup>	6.59±0.04 <sup>bA</sup>
	30	6.55±0.02 <sup>aA</sup>	6.77±0.02 <sup>bC</sup>	6.61±0.02 <sup>bB</sup>
	37	6.51±0.01 <sup>aA</sup>	6.71±0.02 <sup>aC</sup>	6.66±0.01 <sup>aB</sup>
Moisture content (%)	1	48.53±0.22 <sup>aA</sup>	48.87±0.14 <sup>bA</sup>	48.63±0.19 <sup>abA</sup>
	10	49.07±0.28 <sup>abB</sup>	46.54±0.43 <sup>aA</sup>	48.66±0.28 <sup>abB</sup>
	21	47.93±0.73 <sup>abB</sup>	46.33±0.59 <sup>aA</sup>	49.73±0.08 <sup>bC</sup>
	30	48.39±0.48 <sup>abB</sup>	46.47±0.44 <sup>aA</sup>	48.10±0.89 <sup>aAB</sup>
	37	48.50±0.34 <sup>aA</sup>	47.16±0.30 <sup>abB</sup>	48.56±0.31 <sup>abA</sup>
Water activity	1	0.92±0.01 <sup>dA</sup>	0.92±0.01 <sup>bA</sup>	0.92±0.03 <sup>bA</sup>
	10	0.90±0.01 <sup>bA</sup>	0.92±0.01 <sup>bC</sup>	0.91±0.01 <sup>aB</sup>
	21	0.91±0.01 <sup>cA</sup>	0.91±0.01 <sup>aA</sup>	0.91±0.01 <sup>aA</sup>
	30	0.89±0.01 <sup>aA</sup>	0.91±0.02 <sup>aA</sup>	0.92±0.05 <sup>bA</sup>
	37	0.92±0.02 <sup>cA</sup>	0.92±0.01 <sup>bA</sup>	0.92±0.01 <sup>bA</sup>
TBARS (mg of MDA/kg of black pudding)	1	1.30±0.08 <sup>aA</sup>	2.56±0.30 <sup>abB</sup>	3.02±0.13 <sup>aC</sup>
	10	1.28±0.02 <sup>aA</sup>	2.21±0.23 <sup>abB</sup>	2.46±0.24 <sup>bB</sup>
	21	1.24±0.15 <sup>aA</sup>	1.33±0.16 <sup>bA</sup>	1.32±0.06 <sup>cA</sup>
	30	1.15±0.07 <sup>aA</sup>	1.34±0.10 <sup>bA</sup>	1.27±0.07 <sup>cA</sup>
	37	1.13±0.06 <sup>aA</sup>	1.30±0.15 <sup>bA</sup>	1.50±0.06 <sup>cB</sup>

E301- sodium ascorbate. Different lower-case letter in the same column indicate significant difference ( $P < 0.05$ ) by Scheffe test. Different capital letters in the same row indicate significant difference ( $P < 0.05$ ) by Tukey's test.

496

497  
 498  
 499  
 500  
 501  
 502  
 503  
 504  
 505  
 506  
 507


**Table 4.** Component variance analysis for the measured parameter of black pudding considered the three treatment and the 5 storage period

	Variance origin	DF	F	p	Variance percentage
pH	Treatment (T)	2	234.1	0.0000***	37.8
	Days (D)	4	59.4	0.0000***	15.8
	TxD	8	55.2	0.0000***	44.0
	Residual	30			2.4
Moisture	Treatment (T)	2	63.2	0.0000***	48.0
	Days (D)	4	6.4	0.0007***	7.0
	TxD	8	9.7	0.0000***	33.4
	Residual	30			11.6
Water activity	Treatment (T)	2	276.3	0.157 <sup>n.s.</sup>	--
	Days (D)	4	196.7	0.064 <sup>n.s.</sup>	--
	TxD	8	116.8	0.281 <sup>n.s.</sup>	--
	Residual	30			
TBARS	Treatment (T)	2	149.8	0.0000***	25.7
	Days (D)	4	156.1	0.0000***	44.7
	TxD	8	32.3	0.0000***	27.0
	Residual	30			2.6

DF – degrees of freedom; n.s. – not significant,  $p > 0,05$ ; \* Significant,  $0,01 < p < 0,05$ ; \*\* very significant,  $0,001 < p < 0,01$ ; \*\*\* highly significant,  $p < 0,001$

508  
 509  
 510  
 511  
 512  
 513  
 514  
 515  
 516

517  
518  
519  
520  
521  
522  
523  
524  
525  
526



527  
528  
529  
530  
531  
532  
533  
534  
535  
536

**Figure 1.** Chromatographic profile at 280 nm of bee pollen extract. Numbers in the figure correspond to the eluted UHPLC peaks for which UV and MS data is summarized in **Table 1**.

537


538

539

540


541

542



543

544



545

546

547 **Figure 2.** Appearance and flavor evaluation by sensory analysis of black pudding self-life

**Table 1.** UHPLC-DAD-ESI-MS<sup>n</sup> data for bee pollen.

PN	t <sub>R</sub> (min)	λ <sub>max</sub> (nm)	(m/z)	MS <sup>n</sup> ions (m/z)	Probable compound
1	10	312	337	MS <sup>2</sup> [337]: 173, 162, 191	coumaroyl quinic acid
2	10.3	258, 356	625	MS <sup>2</sup> [625]: 316, 317, 271, 461, 479, 609	myricetin- <i>O</i> -rutinoside
3	10.7	264, 351	609	MS <sup>2</sup> [609]: 447, 285	luteolin- <i>O</i> -dihexoside
			625	MS <sup>2</sup> [625]: 301, 463, 445	quercetin- <i>O</i> -dihexoside
4	10.9	262, 353	479	MS <sup>2</sup> [479]: 316, 317	myricetin- <i>O</i> -hexoside
5	11	261, 357	711	MS <sup>2</sup> [711]: 667, 316, 317	myricetin- <i>O</i> -(malonyl)rutinoside
6	11.2	270, 355	639	MS <sup>2</sup> [639]: 459, 315	isorhamnetin- <i>O</i> -dihexoside
			595	MS <sup>2</sup> [595]: 301, 463	quercetin- <i>O</i> -hexosyl-pentoside
7	11.4	256, 308, 354	609	MS <sup>2</sup> [609]: 301, 463	quercetin- <i>O</i> -rutinoside
			565	MS <sup>2</sup> [565]: 521, 316, 317	myricetin- <i>O</i> -(malonyl)hexoside
8	11.7	266, 353	609	MS <sup>2</sup> [609]: 301	quercetin- <i>O</i> -rutinoside
9	12	266, 351	755	MS <sup>2</sup> [755]: 609, 593, 573, 285, 255	luteolin-di- <i>O</i> -hexosyl-rhamoside
10	12.1	257, 353	695	MS <sup>2</sup> [695]: 661, 609, 301	Quercetin- <i>O</i> -(malonyl)rutinoside
11	12.2	255, 354	623	MS <sup>2</sup> [623]: 315, 459	Isorhamnetin- <i>O</i> -rutinoside
12	12.5	250sh, 297, 308	437	MS <sup>2</sup> [437]: 317	hydroxybenzoyl myricetin
13	12.6	256, 354	549	MS <sup>2</sup> [549]: 505, 301, 463	quercetin- <i>O</i> -(malonyl)hexoside
14	13.0	265, 350	679	MS <sup>2</sup> [679]: 635, 301, 575, 255	quercetin derivative
15	13.2	257, 351	447	MS <sup>2</sup> [447]: 301	quercetin-3- <i>O</i> -rhamnoside
16	13.6	271, 351	563	MS <sup>2</sup> [563]: 315, 519, 545	isorhamnetin- <i>O</i> -(malonyl)hexoside
17	13.8	256, 354	533	MS <sup>2</sup> [533]: 489, 285	luteolin- <i>O</i> -(malonyl)hexoside
18	13.8	mix	317	MS <sup>2</sup> [317]: 179, 151	myricetin
19	14.0	255, 353	563	MS <sup>2</sup> [563]: 519, 315, 359	isorhamnetin- <i>O</i> -(malonyl)hexoside
20	14.5	245, 296sh, 319	631	MS <sup>2</sup> [631]: 495, 317	myricetin- <i>O</i> -dihydroferuloyl protocatechuic acid
21	15.2	245, 296, 310	615	MS <sup>2</sup> [615]: 479; MS <sup>3</sup> [479]: 359; MS <sup>4</sup> [359]: 317	myricetin- <i>O</i> -acetyl hydroxybenzoyl protocatechuic acid

22	15.4	245, 296, 310	615	MS <sup>2</sup> [615]: 479; MS <sup>3</sup> [479]: 359; MS <sup>4</sup> [359]: 317	myricetin- <i>O</i> -acetyl hydroxybenzoyl <i>protocatechuic acid</i>
23	16.1	240, 295, 308	599	MS <sup>2</sup> [599]: 463; MS <sup>3</sup> [463]: 343; MS <sup>4</sup> [343]: 301	quercetin- <i>O</i> -acetyl hydroxybenzoyl <i>protocatechuic acid</i>
24	16.3	240, 295, 309	599	MS <sup>2</sup> [599]: 479; MS <sup>3</sup> [479]: 359; MS <sup>4</sup> [359]: 317	myricetin- <i>O</i> -acetyl hydroxybenzoyl <i>hydrobenzoic acid</i>
25	17.0	240, 295, 312	583	MS <sup>2</sup> [583]: 463; MS <sup>3</sup> [463]: 343; MS <sup>4</sup> [343]: 301	quercetin- <i>O</i> -acetyl hydroxybenzoyl <i>hydrobenzoic acid</i>
26	17.2	240, 295, 308	583	MS <sup>2</sup> [583]: 463; MS <sup>3</sup> [463]: 343; MS <sup>4</sup> [343]: 301	quercetin- <i>O</i> -acetyl hydroxybenzoyl <i>hydrobenzoic acid</i>
27- 29	17.8-18.4	242, 270-294	785	MS <sup>2</sup> [785]: 665; MS <sup>3</sup> [665]: 545; MS <sup>4</sup> [545]: 503, 459, 399	<i>O</i> -dihydroxybenzoyl acetyl malonyl coumaric acid flavonoid derivative

Peak numbers (PN) correspond to those depicted in Figure 1

## Highlights

- Bee pollen is a healthy product.
- Chemical and sensory characterization of Black pudding with pollen.
- The work suggest that bee pollen could be a natural alternative to prevent the lipid oxidation in black pudding.



## Conflict of Interest and Authorship Conformation Form

Please check the following as appropriate:

- All authors have participated in (a) conception and design, or analysis and interpretation of the data; (b) drafting the article or revising it critically for important intellectual content; and (c) approval of the final version.
- This manuscript has not been submitted to, nor is under review at, another journal or other publishing venue.
- The authors have no affiliation with any organization with a direct or indirect financial interest in the subject matter discussed in the manuscript
- The following authors have affiliations with organizations with direct or indirect financial interest in the subject matter discussed in the manuscript:

Author's name

Affiliation

Ofélia Anjos<sup>1,2,3</sup>, Rodrigo Fernandes<sup>1</sup>, Susana M. Cardoso<sup>4</sup>, Teresa Delgado<sup>3</sup>, Nelson Farinha<sup>3</sup>, Vanessa Paula<sup>6</sup>, Letícia M. Estevinho<sup>5</sup>, Solange Teresinha Carpes<sup>7</sup>

1 Instituto Politécnico de Castelo Branco, Portugal

2 Centro de Biotecnologia de Plantas da Beira Interior, Castelo Branco, Portugal

3 Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Portugal

4 QOPNA & LAQV-REQUIMTE, Department of Chemistry, University of Aveiro, Aveiro, Portugal

5 Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Portugal

6 School of Agricultural Sciences, Polytechnic Institute of Bragança, Bragança, Portugal

7 Department of Chemistry, Federal University of Technology – Paraná (UTFPR), Pato Branco, PR, Brazil