

Journal Pre-proof

Biochar amendment increases bacterial diversity and vegetation cover in trace element-polluted soils: A long-term field experiment

Paloma Campos, Ana Z. Miller, Sergio A. Prats, Heike Knicker, Nikolas Hagemann, José M. De la Rosa



PII: S0038-0717(20)30310-2

DOI: <https://doi.org/10.1016/j.soilbio.2020.108014>

Reference: SBB 108014

To appear in: *Soil Biology and Biochemistry*

Received Date: 18 June 2020

Revised Date: 5 September 2020

Accepted Date: 13 September 2020

Please cite this article as: Campos, P., Miller, A.Z., Prats, S.A., Knicker, H., Hagemann, N., De la Rosa, José.M., Biochar amendment increases bacterial diversity and vegetation cover in trace element-polluted soils: A long-term field experiment, *Soil Biology and Biochemistry* (2020), doi: <https://doi.org/10.1016/j.soilbio.2020.108014>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

1 **Biochar amendment increases bacterial diversity and vegetation cover in trace**
2 **element-polluted soils: a long-term field experiment**

3

4 **Authors:** Paloma Campos¹, Ana Z. Miller^{2*}, Sergio A. Prats³, Heike Knicker¹, Nikolas
5 Hagemann^{4,5}, José M. De la Rosa¹

6

7 **Affiliations:** 1) Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS-
8 CSIC), Av. Reina Mercedes 10, 41012, Seville, Spain; 2) Laboratorio Hercules,
9 University of Évora, Largo Marquês de Marialva 8, 7000-809, Évora, Portugal; 3)
10 Centre for Environmental and Marine Studies (CESAM), Department of Environment
11 and Planning, University of Aveiro, Aveiro, Portugal; 4) Agroscope Zurich,
12 Reckenholzstr. 191, Zurich, Switzerland; 5) Ithaka Institute, Arbaz, Switzerland

13

14 *corresponding author: anamiller@uevora.pt

15

16 **Abstract**

17 Application of biochar has been widely suggested as a remediation tool for trace
18 element-polluted soils, but the impact of biochar on microbial communities and on
19 native plants remain largely unknown. To overcome this knowledge gap, biochar
20 produced from rice husk and olive pit were applied at a rate of 8 t ha⁻¹ into a soil with
21 two contrasting levels of trace elements (high and moderate) to study their effects on
22 soil microbial community composition, vegetation cover and soil properties after 1, 6,
23 12 and 20 months under field conditions. Differences in bacterial community
24 composition were studied using the Illumina Miseq technology of the 16S rRNA gene.

25 Although variations in soil properties and ecological function were seasonal and soil-
26 type dependent, biochar application enhanced soil properties and vegetation cover in
27 the moderately polluted soil (MPS), and increased microbial diversity as well as
28 vegetation cover in the highly polluted soil (HPS). Enzymatic activities and soil
29 respiration rates were not modified with the application of biochar, but increased total
30 carbon content of soils. The application of biochar from crop residues to trace-element
31 contaminated soils provided environmental benefits, including plant diversity and
32 growth, as well as the increase of bacterial diversity and carbon sequestration.

33

34 **Keywords:** Soil remediation; pyrogenic carbon; heavy metals; soil microbial
35 community; plant diversity

36 1. Introduction

37 Trace element-polluted soils is a worldwide concern comprising 37 % of the degraded
38 soils in the European Union (EEA, 2007). Ex-situ decontamination of polluted soils is
39 generally unfeasible due to land size and soil contamination levels, which are difficult to
40 effectively and economically reduce with conventional soil remediation procedures
41 (Tack et al., 2018). Biochar, the C-rich porous solid residue produced by the thermal
42 conversion of biomass under the partial or total absence of oxygen (pyrolysis, e.g.
43 Hagemann et al. 2018), has the ability to immobilize trace elements and increase the
44 pH of acidic soils reducing trace element mobility and bioavailability. Karer et al. (2015)
45 reported a decrease in NH_4NO_3 -extractable fraction of Pb, Zn and Cd with biochar
46 amendment, but an increase of Cu. Beesley et al. (2010) also reported an
47 immobilization of Cd and Zn and a mobilization of Cu after biochar application.
48 Oustriere et al. (2017) showed long-term Cu stabilization due to biochar addition into a
49 contaminated soil, whereas Uchimiya et al. (2012) reported Cu immobilization but
50 mobilization of Sb. These discrepancies are probably due to the complexity of
51 immobilization mechanisms and different biochar compositions and properties, but also
52 due to differences in the soil properties, e.g. in pH. In fact, previous studies already
53 demonstrated that the efficacy of biochar as a soil amendment greatly depends on its
54 pyrolysis conditions and feedstock (Campos et al., 2020; De la Rosa et al., 2014). For
55 instance, Kammann et al. (2012) showed a significant increase in biomass yield after
56 applying 50 Mg ha^{-1} of peanut hull biochar to a Luvisol. Gascó et al. (2016) reported
57 that β -glucosidase, phosphomonoesterase and phosphodiesterase activities were
58 lower when a sandy loam soil was incubated with 8 % (w/w) of biochar produced from
59 pig manure at $500 \text{ }^\circ\text{C}$ whereas the biochar produced at $300 \text{ }^\circ\text{C}$ increased
60 dehydrogenase activity. The study of Shen et al. (2019) demonstrated that biochar
61 produced at $500 \text{ }^\circ\text{C}$ was more effective in the removal of lead from soil solution than
62 the biochar produced at $300 \text{ }^\circ\text{C}$. Generally, biochar produced at $500 \text{ }^\circ\text{C}$ has high pH
63 and water holding capacity, and high degree of aromatization (Campos et al., 2020).
64 The effects of biochar on the physical and chemical properties of agricultural soils have
65 been profoundly studied and during the last years special attention has been paid to
66 the study of biochar as soil amendment for the retention of contaminants (Uchimiya et
67 al., 2011; Kumar et al., 2018; De la Rosa et al., 2019). Within this context, effects of
68 biochar addition on soil microbiota, which play a vital role in soil ecosystem stability,
69 soil quality and soil nutrient cycle (Lehmann et al., 2011), are highly relevant. Li et al.
70 (2019) reported a decrease in *Actinobacteria* with biochar application into purple soil,
71 whereas Ali et al. (2019) reported an increase for pesticide-contaminated soil. Most of
72 these studies on soil microbial diversity in polluted soils after biochar application are

73 pot-based experiments (Jiang et al., 2017; Han et al., 2017), which are likely to be
74 important in the quest to constrain the numerous influencing factors, but are less
75 realistic than field studies.

76 An aspect also worth further researching is the biodegradability of biochar in soils.
77 Biochar has traditionally been considered a material of high chemical and biochemical
78 stability, which predominantly contains C in the form of condensed aromatic rings. This
79 fraction of C is hardly decomposed by soil biota due to its recalcitrant nature (Kuzyakov
80 et al., 2009, 2014). Nevertheless, recent studies indicate a much lower biochemical
81 stability (Knicker et al., 2013, De la Rosa et al., 2018). Thus, the effects of biochar
82 application on soil CO₂ emissions are often ambiguous and previous studies reported
83 increases, decreases or no changes (Bamminger et al., 2014; Kolb et al., 2009; Paz-
84 Ferreira et al., 2012). Hence and considering that changes on soil properties promoted
85 by biochar application may affect soil microbial communities (Luo et al., 2013; Su et al.,
86 2015; Xu et al., 2016) and soil CO₂ emissions, their assessment deserve further
87 attention.

88 The application of low degradability of biochar in trace element-polluted soils would
89 allow an effective *in situ* remediation by enhancing soil quality and improving its
90 capability to perform soil ecological functions. To test this hypothesis, we applied rice
91 husk (RH) and olive pit (OP) biochar into two trace element-polluted acidic soils under
92 field conditions to study their effects on soil physicochemical properties, soil CO₂
93 emissions and enzymatic activities, as well as soil microbial community composition
94 and vegetation cover after 1, 6, 12 and 20 months of biochar application.

95

96 **2. Materials and methods**

97 **2.1. Biochar samples**

98 Rice husk (RH) and olive pit (OP) were used as feedstock to produce biochar due to
99 their great abundance in Mediterranean countries. RH is a siliceous-rich raw material
100 with relatively low C content, while OP is a hard-wood biomass, mainly composed of
101 cellulose and lignin. The company Orivarzea S.A. (Portugal) provided the RH biomass,
102 whereas OP was provided by *Cooperativa Nuestra Señora de los Ángeles* (Montellano,
103 Spain).

104 Prior to pyrolysis process, feedstock was dried at 40 °C during 48 h, homogenized and
105 stored in sealed plastic bags at 4 °C. The RH and OP biochar (RHB and OPB) were
106 produced in a continuously feed pyrolysis reactor with a screw conveyor (PYREKA,
107 Pyreg GmbH, Dörth, Germany, cf. Hagemann et al. 2020) under N₂ flux at Agroscope
108 Zurich (Switzerland). The pyrolysis temperature was 500 °C and the residence time

109 was 12 minutes. Biochar was not quenched with water and stored in plastic bags.
110 Biochar characteristics are shown in Table 1.

111

112 **2.2. Area of study and experimental design**

113 The field experiment was conducted at 'Las Doblas' site (37° 23' 7.152"N, 6° 13'
114 43.175") over a period of 20 months. This place is located close to the Guadiamar river,
115 10 km from the former mine "Los Frailes" close to Aznalcollar, Southern Spain. On the
116 25th of April 1998, after a major mining accident, a huge amount of toxic sludge spilled
117 out from a tailing reservoir of this large open-pit mine, causing high levels of heavy
118 metals to leach into the soil and groundwater. Figure 1 shows the location of the field
119 experiment. The area belongs to a typical dry Mediterranean climate region, with hot
120 and extended summers, soft winters and a very pronounced variation in the
121 precipitation rate (AEMET, 2018).

122 The sandy loam soil of the area is classified as Fluvisol (IUSS Working Group WRB,
123 2015). In this study, two nearby sites were selected according to their contamination
124 level and acidity, comprising a highly polluted soil (HPS) and a moderately polluted soil
125 (MPS). HPS is a bare soil with high acidity and concentrations of heavy metals, as
126 previously described in Cabrera et al. (1999) and Martín-Peinado et al. (2015). These
127 bare spots account over 200 ha of lands affected by the accumulation of residual toxic
128 sludge of the spill. In contrast, MPS areas were subjected to a decontamination
129 programme by the Andalusian regional government which included the removal of the
130 toxic sludge (Arenas et al., 2008). Despite the decontamination efforts, MPS also
131 shows relatively high concentration of Ba, Cu, Fe, Pb and Zn (Campos and De la Rosa,
132 2020). Soil pH, total carbon (TC), total nitrogen (TN) and total hydrogen (TH) contents
133 of HPS and MPS are shown in Table 1.

134 In April 2018, 12 plots of 1 m × 1 m each were randomly established in HPS and MPS
135 sites (6 plots per site). RHB and OPB were applied as produced and mixed into the first
136 10 cm of soil at a dose of 8 t ha⁻¹ (Plots ID: RHB_HPS, OPB_HPS, RHB_MPS and
137 OPB_MPS). In addition, control plots without amendment were established for both
138 areas (C_MPS and C_HPS) but received the same mechanical treatment. For all the
139 plots, ground vegetation (shrub and grass) was manually removed; the soil was then
140 homogenised using a manual rake.

141 Four sampling campaigns were performed after 1, 6, 12 and 20 months of biochar
142 incorporation into soils (hereafter: t₁, t₆, t₁₂ and t₂₀, respectively). For each plot, five
143 samples of soil were taken randomly from the first 10 cm depth to create a composite
144 sample per plot. An aliquot of the composite sample was immediately used for
145 enzymatic analyses, other aliquot was stored in sterile Whirl-pak® bags at -80 °C for

146 DNA-based analysis and the remaining material was dried at 40 °C during 48 h, sieved
147 (<2 mm) and stored in sealed bags at 4 °C.

148

149 **2.3. Chemical and biochemical analysis**

150 The pH was measured in triplicates in the supernatant of a 1:5 (w/v) soil:0.01 M CaCl₂
151 solution ratio mixture after 30 minutes shaking and 30 minutes resting, using a pH
152 meter (CRISON pH Basic 20).

153 The soil moisture (%) was determined on the dry weight basis: 20 g of moist soil was
154 weighed, dried at 40 °C during 24h and re-weighed. Total soil moisture (%) was
155 determined for soil samples dried at 105 °C for 24 h.

156 Total C (TC) was obtained by dry combustion (1050 °C) using an elemental analyzer
157 (TRUSPEC CHNS MICRO, LECO, St. Joseph, MI, USA).

158 The water holding capacity (WHC) was measured following the procedure and formula
159 described in Campos et al. (2020). The WHC is expressed as the percentage relatively
160 to the total dry weight of the sample:

161

$$WHC (\%) = \frac{\text{Water retained weight}}{\text{Initial weight of the dry sample}} \cdot 100 \text{ (Eq. 1)}$$

162

163 For elucidating microbial oxidative activities in soil, dehydrogenase activity was
164 determined according to the method of Trevors (1986). Briefly, soil samples were
165 incubated for 20 h with 1 M TRIS–HCl buffer (pH 7.5) and 2(p-iodophenyl)-3-(p-
166 nitrophenyl) 5-phenyl tetrazolium chloride (INT), that was used as the electron
167 acceptor. After adding methanol, the idonitrotetrazolium formazan (INTF) produced
168 was measured spectrophotometrically at 490 nm.

169 In addition, soil β -glucosidase activity was measured according to the method of
170 Tabatabai (1982). Briefly, 1 g of soil was incubated 1 h at 37 °C with p-nitrophenyl- β -D-
171 glucopyranoside. After addition of CaCl₂, the p-nitrophenol was extracted by filtration
172 and measured using a spectrophotometer (Jenway, model 6315, UK) at 400 nm. β -
173 glucosidase and dehydrogenase activities were measured in both unamended and
174 biochar-amended soils at t_1 , t_6 , t_{12} and t_{20} .

175 All chemical and biochemical analyses of the samples were performed in triplicate.

176

177 **2.4. Measurement of soil CO₂ efflux (Soil respiration)**

178 Soil respiration (carbon decomposition by microorganisms and ground root respiration)
179 was determined by measuring the CO₂ effluxes and expressed as $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$.

180 For each plot, 3 PVC collars (10 cm diameter and 5 cm high) were installed 3 cm into

181 the soil and measurements were conducted in triplicate using the soil CO₂ flux chamber
182 LI-COR 6400-09 (LI-COR, Nebraska, USA) at t₁, t₆, t₁₂ and t₂₀. Soil temperature was
183 monitored using a thermocouple probe (Li6000-09 TC, LiCor Inc) inserted to a depth of
184 5 cm near the soil CO₂ flux chamber.

185

186 **2.5. Effects on vegetation**

187 The vegetation species were carefully identified and the number of individuals per plot
188 were accounted at t₁₂. Subsequently, the total plant biomass was determined by
189 harvesting and measuring the fresh weight per plot.

190 In order to determine the percentage of vegetation cover at time t₂₀, high resolution
191 photographs were taken for each plot using a digital camera (Canon Inc., Canon 7D,
192 Japan) installed on a tripod at a height of 1.5 m. Digital images were then analysed
193 using the open source image-processing software Image J. The area covered by green
194 plants was selected by adjusting the hue levels in the colour threshold tool. The
195 percentage of vegetation cover was determined by using the following equation 2:

196

$$197 \quad \% \text{ vegetation cover} = \frac{\text{green area}}{\text{total area}} \cdot 100 \quad (\text{Eq. 2})$$

198

199 **2.6. Soil DNA isolation and sequencing**

200 Total DNA was extracted from soil samples using the DNeasy PowerSoil DNA isolation
201 kit (Qiagen, Hilden, Germany), according to the manufacturer's instructions. DNA
202 quality and quantity were tested. As a standard procedure, 1.5% agarose gel
203 electrophoresis was performed with 1 µL of gDNA of each sample to test the integrity
204 and purity. DNA concentrations were verified using a Qubit 2.0 fluorometer (Life
205 Technologies, Carlsbad, CA, USA). Qubit broad-range reagent was used for
206 determining the DNA concentration of MPS samples, whereas Qubit high-sensitivity
207 reagent was needed for HPS samples due to their low amount of DNA.

208 Library construction was performed according to the Illumina 16S Metagenomic
209 Sequencing Library preparation protocol by STAB VIDA Sequencing Services
210 (Portugal). MiSeq Reagent Kit v3 in the Illumina MiSeq platform was used for
211 sequencing new generated DNA fragments. 300 bp paired-end sequencing reads were
212 used.

213 The microbial community composition and diversity (alpha and beta-diversity) were
214 determined after bioinformatics processing of the 16S rRNA gene sequences.
215 Sequence quality control was performed using QIIME2 v2019.1.0 (Bolyen et al., 2019).
216 The reads were denoised using the DADA2 plugin, organized in operational taxonomic

217 units (OTUs) and classified by taxon using the SILVA database, with a clustering
218 threshold of 97% similarity. OTUs were considered as significant only when they
219 contained at least 10 sequence reads. This procedure results in an abundance table
220 with taxonomy information, which were further analysed and visualized using the online
221 web-tool Calypso (Zakrzewski et al., 2017).

222 The raw reads were deposited in NCBI Sequence Read Archive (SRA) database
223 (<https://submit.ncbi.nlm.nih.gov/about/sra/>) under the accession number
224 PRJNA637319.

225

226 **2.7. Data analysis**

227 Data of soil and biochar characteristics are expressed as mean \pm standard error (SE)
228 of triplicate measurements. Data of the samplings are expressed as mean \pm SE of the
229 five composite samples per treatment. Number of species and number of individuals
230 are expressed as median.

231 Shapiro-Wilk test was used to verify normality and Levene test was used to test
232 homoscedasticity of the data. Normal distributed response variables were analysed by
233 one-way ANOVA followed by the Tukey's Honestly Significant Difference test. The level
234 of significance used was 0.05. When response variables were non-normal, Kruskal
235 Wallis followed by Mann Whitney U tests were conducted. Pearson correlation ($p <$
236 0.05) was conducted to examine relationships between soil properties, soil microbial
237 community and vegetation data. Statistical analyses were carried out using IBM SPSS
238 Statistics 26.0 (SPSS, Chicago, USA) for Windows.

239

240 **3. Results**

241 **3.1. Soil pH and moisture**

242 Soil pH of HPS samples were more acidic than MPS (3.57–3.77 and 4.18–5.11,
243 respectively) (Table 2). Biochar addition did not significantly enhance soil pH of HPS.
244 For MPS, biochar amendment clearly increased soil pH, but this increase was
245 mitigated over the time span of the experiment.

246 Soil moisture determined by drying at 40 °C was greater for MPS than HPS samples
247 with comparable treatments (Table S1). Biochar addition augmented soil moisture (40
248 °C) at t_6 and to a lesser extent at t_{20} . As expected, seasonal changes modified the soil
249 moisture at 40 °C and 105 °C, with a significant drop at t_{12} , followed by a considerable
250 increase during the autumn (t_{20}). MPS showed a greater WHC (%) than HPS, and were
251 affected by seasonal changes, showing in general a similar trend as the soil moisture
252 content (Table 2).

253

254 **3.2. Total carbon content and soil respiration**

255 Biochar addition caused a non-significant increase ($p > 0.05$) of the C content of the
 256 amended soils when compared with control plots (Table 2). At t_{12} , the C content
 257 significantly increased due to OPB amendment in HPS samples (9.6 to 16.6 g kg⁻¹).

258 The CO₂ emission rates were always higher in MPS than in HPS. The latter showed
 259 very low respiration rates. Significant differences were not observed with biochar
 260 application into the soils, nor differences between both biochar samples for the same
 261 soil type (Table 2).

262

263 **3.3. Soil enzymatic activities**

264 β -glucosidase activity at t_1 of MPS was greater than HPS (0.69-1.45 vs 0.17-0.39 μmol
 265 PNF g⁻¹ h⁻¹) (Fig. 2). At this time, the HPS control soil showed a greater β -glucosidase
 266 activity than biochar amended soils. Nevertheless, this difference disappeared at t_6 and
 267 t_{12} . This enzymatic activity was greater for both amended treatments than for control in
 268 MPS at t_1 . Nevertheless, at t_6 only OPB addition maintained a greater β -glucosidase
 269 activity than control soils and at t_{20} no significant differences were observed.

270 MPS plots showed greater dehydrogenase activity than in HPS in all the cases.
 271 Similarly to the trend observed for β -glucosidase activity, dehydrogenase activity
 272 showed seasonal changes and during the first 6 months of the experiment MPS soils
 273 amended with biochar showed lower values than control soils.

274

275 **3.4. Effects on vegetation development**

276 A total of 14 different species were observed in MPS plots which were not found in
 277 HPS plots (*Echium gaditanum*, *Lotus parviflorus*, *Trifolium arvense*, *Ornithopus*
 278 *compresus*, *Anagallis arvensis*, *Bartsia trixago*, *Trifolium* sp., *Vulpia ciliata*, *Trifolium*
 279 *vesiculosum*, *Trifolium striatum*, *Hypochaeris glabra*, *Astragalus pelecinus*, *Trifolium*
 280 *campestre*, *Spergularia media*, *Silene sclerocarpa* and *Petrorhagia nanteuillii*). In
 281 contrast, solely one plant species (*Sonchus oleraceus*) was found in HPS plots which
 282 was not found in MPS (Table S2). *Rosmarinus officinalis*, *Chamaemelum mixtum*,
 283 *Agrostis truncatula*, *Spergularia rubra*, *Logfia minima* and *Cynodon dactylon* were
 284 found in HPS and MPS plots. Furthermore, biochar application enhanced vegetation
 285 diversity, as *Trifolium campestre*, *Spergularia media*, *Silene sclerocarpa* and
 286 *Petrorhagia nanteuillii* solely grew in MPS biochar-amended plots. *Logfia minima* was
 287 strictly found in OPB plots, but not in the unamended ones (Table S2).

288 Figure 3a and b shows the average number of different plant species and individuals,
 289 respectively, in biochar-amended and unamended plots 12 months after the setup of
 290 the experiment. A greater diversity of vegetation species was observed in MPS than in

291 HPS plots (Fig. 3a), as also occurred for the number of individuals (Fig. 3b). One-way
292 ANOVA showed that the number of individuals per square meter (Fig. 3b) and
293 vegetation cover (Fig. 3c) in HPS were significantly ($p < 0.05$) lower than in MPS. The
294 application of OPB in HPS plots significantly increased the area of vegetation cover in
295 comparison with the control plots (C_HPS; Fig. 3c). In contrast, although an increase in
296 vegetation cover was observed for HPS amended with RHB, it was not statistically
297 significant (Fig. 3c).

298 Concerning the fresh weight per plot (Fig. 3d), it increased significantly due to OPB
299 application into HPS, but no significant differences were found for MPS plots. However,
300 the increase of plant fresh weight in OPB_HPS was statistically similar to OPB_MPS.
301 For MPS plots, the application of biochar did not promote statistical differences ($p >$
302 0.05) among samples for all parameters (Fig. 3 b–d).

303

304 **3.5. Pearson correlations of soil properties**

305 Table S3 shows that pH was positively correlated with soil moisture measured at 40 °C
306 and soil respiration after 6 months ($p < 0.05$; Pearson coefficients were 0.877 and
307 0.917, respectively) and 20 months after setup ($p < 0.05$; Pearson coefficients were
308 0.903 and 0.964, respectively). Soil moisture correlated with WHC at month 6 of the
309 experiment ($p < 0.05$; Pearson coefficient 0.835) and with soil respiration after 20
310 months ($p < 0.05$; Pearson coefficient 0.862). Nevertheless, 12 months after biochar
311 application, only the pH was negatively correlated with soil moisture measured at 40 °C
312 ($p < 0.05$; Pearson coefficient -0.984), indicating variability of soil properties with time
313 and climate conditions.

314 Pearson correlations were performed between vegetation results and between
315 vegetation results and soil properties (Table S4). Fresh weight correlated positively
316 with dry weight, number of species and number of individuals ($p < 0.05$; Pearson
317 coefficients between 0.839–0.877). Plots with greater number of species also showed
318 greater number of individuals ($p < 0.05$; Pearson coefficient 0.959). Positive correlation
319 was found between pH and fresh weight, number of individuals and number of species
320 ($p < 0.05$, Pearson coefficients 0.851, 0.867 and 0.949, respectively). In addition,
321 positive correlation was found between WHC and fresh weight ($p < 0.05$; Pearson
322 coefficient 0.870). Negative correlations were found between soil moisture and
323 vegetation results.

324

325 **3.6. Bacterial community composition**

326 **3.6.1 Sequence data**

327 The number of raw sequence reads ranged from 113072 to 237356 for samples
328 collected at t_6 , from 377786 to 704600 for t_{12} and from 291568 to 415478 for t_{20} . After
329 quality filtering and denoising, a total of 2354783 paired-end sequences were obtained
330 for all samples. These sequences were clustered into 16964 OTUs at 97% similarity,
331 containing both assigned and non-identified bacteria. Samples from t_{12} showed the
332 highest number of OTUs (8000), followed by t_{20} (4863) and t_6 with 4101 OTUs. For all
333 the samples, the rarefaction curves reached the plateau (data not shown), suggesting
334 that we obtained a good representation of the microbial communities from both soil
335 types (HPS and MPS).

336

337 **3.6.2. Differences between HPS and MPS**

338 The microbial communities from all the samples were almost exclusively composed of
339 bacteria, with the exception of the control sample HPS (C_HPS), where Archaea
340 accounted for 0.47%, being represented by the *Thaumarchaeota* and *Euryarchaeota*
341 phyla.

342 Differences in the taxonomic composition were clearly observed between both types of
343 soil and between sampling campaigns, particularly between t_6 and t_{12} (Fig. 4 and Table
344 S5). MPS plots at t_6 showed to be more diverse at the phylum level than HPS. At t_6 , the
345 most abundant phyla found in MPS plots were *Proteobacteria*, *Planctomycetes*,
346 *Actinobacteria*, *Acidobacteria*, *Bacteroidetes*, *Gemmatimonadetes* and
347 *Verrucomicrobia*, whereas in HPS *Chloroflexi*, *Proteobacteria*, *Actinobacteria*,
348 *Acidobacteria*, *Firmicutes* and *Saccharibacteria* were the most abundant (Fig. 4a).

349 The *Planctomycetes* phylum was mostly observed in MPS and in very low relative
350 abundance in HPS. *Bacteroidetes*, *Gemmatimonadetes* and *Verrumicrobia* were solely
351 found in MPS (amended and control plots), whereas *Firmicutes* was mostly found in
352 HPS (7% in C_HPS and 2-3% in amended HPS samples). Interestingly,
353 *Saccharibacteria* was solely found in the amended HPS plots.

354 After 1 year (t_{12}), the relative abundance of bacterial phyla changed considerably in
355 comparison with t_6 (Fig. 4b). *Actinobacteria*, *Chloroflexi*, *Proteobacteria*, *Acidobacteria*,
356 *Firmicutes* and *Planctomycetes* contributed to 80% and 90% of the total bacterial
357 sequences in MPS and HPS, respectively. *Verrumicrobia* and *Gemmatimonadetes*
358 were solely found in MPS (amended and control plots). *Patescibacteria* was found in
359 the control and amended MPS samples, as well as in amended HPS.

360 Differences in microbial community composition at the phylum level were not
361 significantly observed between t_{12} and t_{20} (Fig. 4b and c). At t_{20} the predominant phyla
362 were *Actinobacteria*, *Chloroflexi*, *Proteobacteria*, *Acidobacteria*, *Firmicutes*,
363 *Planctomycetes* and *Bacteroidetes* for both soils (Fig. 4c).

364 Remarkable differences in microbial community composition at the order level were
365 noticeable between HPS and MPS plots, as the bacterial sequences in MPS were
366 almost absent in HPS samples (Fig. S1).

367 Principal components analysis (PCA) was computed to explain differences between
368 samples (Fig. 5). At t_6 , the first two components explained 95% of the variation
369 observed (Fig. 5a). The plot of the loadings of PC-1 vs PC-2 defined two clusters,
370 corresponding to each soil type. This showed that microbial diversity from HPS
371 samples is significantly different from MPS samples. However, within cluster 1 (HPS
372 samples), PC2 significantly separates the HPS control sample from the biochar-treated
373 HPS samples along the projected plane (Fig. 5a). In contrast, no significant differences
374 of bacterial community composition were noted between amended and unamended
375 MPS plots at t_6 .

376 At t_{12} (Fig. 5b), the plot also displays clear discrimination between both types of soils,
377 but in addition contains separation within cluster 1 among biochar treatments.

378 At t_{20} (Fig. 5c), PC-1 (70%) vs PC-2 (12%) scores of the control and biochar-treated
379 soils also define two clusters, reinforcing that microbial diversity from both soil types is
380 significantly different along the time span of the field experiment. However, PC-2
381 discriminates samples within each cluster. Specifically, samples treated with OPB for
382 both types of soils were separated from their corresponding control and RHB-treated
383 samples (Fig. 5c), revealing changes in the microbial diversity for OPB-treated soils
384 after 20 months of incubation.

385 Venn diagrams were plotted to calculate the number of unique and shared OTUs
386 among the HPS and MPS samples at t_6 , t_{12} and t_{20} (Fig. S2). Interestingly, the number
387 of shared taxa between treatments (control and amended plots) was remarkably higher
388 than the number of unique taxa, at t_6 , t_{12} and t_{20} , especially in MPS.

389 The number of shared taxa in HPS increased over time (Fig. S2c and e). The largest
390 OTUs numbers were shared between all the MPS samples (260 at t_6 , 504 at t_{12} and
391 330 at t_{20}), whereas the unique taxa ranged between 5 (at t_6 for C_MPS) and 104 (at t_{12}
392 for C_MPS). At t_6 , 34 OTUs (50%) were uniquely present in HPS plots, while in MPS
393 56 OTUs (16%) corresponded to unique taxa (Fig. S2a). At t_{20} (Fig. S2e and f), the
394 percentage of unique taxa was 45% in HPS against 34% in MPS, suggesting that the
395 bacterial communities became in general more similar between the two soil types over
396 time. Focusing on the biochar treatments, there were more overlapped OTUs among
397 the amended plots than between biochar-treated plots and the controls at t_6 (Fig. S2a
398 and b) and t_{12} (Fig. S2c and d).

399

400 **3.6.3. Impact of biochar amendment on HPS and MPS**

401 At month 6 (Fig. 4a), the relative abundance of soil microbiota from the HPS control
402 plot (C_HPS) differed markedly from those treated with biochar (RHB_HPS and
403 OPB_HPS). The *Chloroflexi* phylum was the most abundant in the C_HPS plot, but
404 decreased notably from 83% to 50% in the biochar-amended HPS samples, as well as
405 *Firmicutes*. In contrast, the relative abundance of *Proteobacteria* in HPS increased
406 from 2 to 16% with biochar application, as well as *Acidobacteria* (Fig. 4a and Table
407 S5).

408 For MPS plots, no significant changes were noted on the relative abundance of soil
409 microbiota between biochar-amended and unamended plots.

410 After 1 year of biochar application into HPS (Fig. 4b), the relative abundance of
411 *Actinobacteria* slightly decreased with biochar amendment from 47 to 30-37%.
412 *Chloroflexi* increased from 20% in the control to 40% in the biochar-amended HPS
413 samples.

414 At t_{20} (Fig. 4c), an increase was observed on the relative abundance of *Proteobacteria*
415 (from 14 to 17-23%) and *Bacteroidetes* (from 0.4 to 4-14%) for HPS plots amended
416 with biochar. In contrast, biochar application reduced the relative abundance of
417 *Chloroflexi* (from 35 to 27-28%), *Acidobacteria* (from 12 to 5%) and *Firmicutes* (from 7
418 to 3-5 %) in the HPS samples.

419 In MPS plots at t_{20} , the application of biochar increased the relative abundance of
420 *Actinobacteria* (from 14 to 17-18%) and reduced the abundance of cyanobacteria (from
421 9 to 0.4-1%).

422 At the order level, the most abundant taxa, representing *Chloroflexi* in C_HPS at t_6 ,
423 belonged to the order *Ktedonobacterales* (71%), followed by the enigmatic phylotypes
424 JG30-KF-AS9 and B12-WMSP1 also within the class *Ktedonobacteria*, both
425 contributing to 9% of the total bacterial sequences (Fig. S1a). The relative abundance
426 of this Ktedonobacterial community was almost reduced by half (from 80% to 46%) with
427 the incorporation of biochar. However, the relative contribution of B12-WMSP1 and
428 JG30-KF-AS9 increased significantly from 4% to 32-37% and from 5% to 12-16%,
429 respectively. In contrast, a sharp decrease was observed for members of the order
430 *Ktedonobacterales* (from 71% to 2% in RHB_HPS and 0.5% in OPB_HPS). The
431 abundances of *Rhodospirillales* (within the *Proteobacteria* phylum) and
432 *Acidobacteriales* (within *Acidobacteria*) were also higher in biochar-amended HPS. The
433 order *Bacillales*, belonging to the phylum *Firmicutes* and solely represented by the
434 genus *Alicyclobacillus* in C_HPS, decreased from 7% to 3% in OPB-amended HPS
435 and to 1.6% in RHB-amended HPS.

436 The most abundant orders found across all treatments in MPS plots at t_6 (Fig. S1a)
437 were *Tepidisphaerales* (within *Planctomycetes*), *Sphingomonadales* (within

438 *Alphaproteobacteria*), *Shingobacteriales* (within *Bacteroidetes*), *Burkholderiales* (within
439 *Betaproteobacteria*) and *Rhizobiales* (within *Alphaproteobacteria*).

440 At t_{12} and t_{20} (Fig. S1b and c), a greater bacterial diversity at the order level is observed
441 for all the treatments in comparison with t_6 , and bacterial communities in biochar-
442 amended plots became more similar to their corresponding control plots, particularly at
443 t_{20} (Fig. S1c). Nevertheless, at t_{12} , the relative abundance of B12-WMSP1, representing
444 the Ktedonobacterial community, was higher in the biochar-amended HPS plots (Fig.
445 S1b), as also observed at t_6 . The order *Frankiales*, belonging to the *Actinobacteria*
446 phylum, increased markedly at t_{12} across all treatments, independently of biochar
447 application. It is also worth noting the increase of *Acetobacterales*, representing
448 *Alphaproteobacteria*, with biochar amendments in HPS plots. At t_{20} , communities in
449 biochar-amended plots became in general more similar to their corresponding control
450 plots (Fig. S1c). However, after 20 months differences in the relative abundance of
451 bacterial taxa within MPS samples were noticed for OPB-treated MPS (Fig. S1c).

452

453 **3.7. Bacterial diversity**

454 The diversity of microbial community structure in the HPS and MPS samples was
455 estimated by alpha diversity and richness indices, revealing values significantly
456 different between the samples. MPS samples (control and amended) showed higher
457 alpha diversity (Shannon and Simpson) and richness (Chao1 and OTU count) than
458 HPS samples (Table 3). Shannon index values ranged from 2.00 to 4.24 in HPS
459 samples, with an average of 3.23, and from 4.47 to 5.33 in MPS samples, with an
460 average of 4.73.

461 The observed Simpson index of diversity ranged from 0.71 to 0.97, with an average of
462 0.91 for HPS samples, and from 0.97 to 0.99, with an average of 0.98 for MPS (Table
463 3).

464 Shannon and Simpson index values increased in the HPS due to biochar addition at t_6 .
465 This increase in alpha diversity indices was also observed for MPS plots at t_6 .
466 Regardless of the presence or absence of biochar, alpha diversity increased through
467 the time span of the experiment (Table 3).

468

469 **3.8 Correlation between soil properties and microbial community composition**

470 Soil physicochemical properties and bacterial abundance variables were used to
471 generate correlation heatmaps for t_6 , t_{12} and t_{20} (Fig. 6). At t_6 , pH, WHC and soil
472 respiration were significantly ($p < 0.05$) and positively correlated with the most
473 abundant bacterial phyla found in MPS plots (*Gemmatimonadetes*, *Planctomycetes*,
474 *Verrucomicrobia*, *Bacteroidetes*), and negatively correlated with *Chloroflexi* and

475 *Firmicutes*, which were the most abundant phylum in HPS plots (Fig. 6a and Table S6).
476 Soil moisture measured at 40 °C and total moisture (measured at 105 °C) were also
477 positively correlated with most of bacterial phyla commonly found in MPS samples,
478 particularly *Proteobacteria* and *Acidobacteria*. Soil TC and dehydrogenase activity
479 showed weak correlation (positive or negative) with the most abundant phyla retrieved
480 in the soil samples. Similarly, glucosidase activity showed almost no correlation with
481 soil microbial communities (Fig. 6a).

482 At t_{12} , the most abundant phyla detected in MPS plots were strongly and positively
483 correlated with soil pH, WHC, dehydrogenase and glucosidase activities, as well as
484 with the botanical variables measured at t_{12} (fresh weight, number of plant species and
485 individuals), but negatively correlated with *Actinobacteria* (Fig. 6b and Table S6).
486 *Chloroflexi* was negatively correlated with pH, whereas it was positively correlated with
487 soil moistures measured at 40 and 105 °C ($p < 0.05$). Soil respiration and TC showed
488 no significant relationship with soil microbial communities.

489 At t_{20} , all the soil physicochemical parameters measured in this study were positively
490 correlated with MPS microbial communities, and negatively correlated with *Chloroflexi*,
491 which was almost exclusively found in HPS samples (Fig. 6c and Table S6).

492

493 **4. Discussion**

494 Physical, chemical and biological parameters were monitored for 6, 12 and 20 months
495 in biochar-amended soils with two different levels of trace-element contamination under
496 field conditions. These parameters (pH, carbon content, WHC, soil moisture, enzymatic
497 activities, soil respiration, vegetation cover and microbial diversity) were selected to
498 integrate the three types of soil quality indicators, which allow assessing the capability
499 of a soil to perform its ecological functions (Arias et al., 2005).

500 Soil properties and, consequently, their plant and microbial diversity were very different
501 in HPS and MPS, independently of biochar addition. The soil properties of HPS plots
502 measured before biochar application indicated a very degraded soil with extreme
503 difficulties to sustain ecological functions. Although biochar application induced
504 changes on soil properties, climatic conditions need to be considered, as changes
505 between samplings were notable.

506 The dehydrogenase activity (DHA) has been also proposed as a good indicator of the
507 toxicity of trace elements (Dick et al., 1996). Under acidic conditions, this enzymatic
508 activity can be inhibited due to the destruction of ion and hydrogen bonds in the
509 enzyme active centre and the alteration of its three-dimensional shape (Frankenberger
510 and Johanson, 1982). This explains the greater values of dehydrogenase activity
511 observed for the less acidic MPS plots, in comparison with HPS, and the positive

512 correlation between soil pH and dehydrogenase activity (Fig. S3). The low β -
513 glucosidase activity measured for all HPS plots, regardless of biochar addition, can be
514 related to soil pH (Eivazi and Tabatai, 1990), and the low abundance of labile organic
515 matter (Ferraz de Almeida et al., 2015). This low enzymatic activity indicates a high
516 recalcitrance of the applied biochar in HPS, as biochar has condensed aromatic
517 structures that make them less available to microbial degradation (Elzobair et al., 2016;
518 Günal et al., 2018; Sohi et al., 2010). Despite of the increase of C content in soils
519 caused by biochar addition, respiration measurements showed that the application of
520 OPB or RHB to the Fluvisol did not modify CO₂ emissions rates. This is similar to the
521 findings previously reported by other authors (Sun et al., 2014; Phongthep et al., 2017).
522 In this study, no priming effect is found and a high stability of both sorts of biochar can
523 be predicted. Considering that soil metal pollution is a significant environmental issue,
524 the use of biochar is worthwhile for the remediation of trace element-polluted soils.

525 As expected, MPS plots showed greater diversity and abundance of vegetation species
526 than HPS (Fig. 3). Comparing both biochar treatments, the application of OPB
527 enhanced not only plant diversity but also the primary productivity in HPS.

528 The combination of digital image analysis, for measuring the total area of soil covered
529 by the vegetation canopy, and the plant fresh weight approach, which provides
530 information on the plant yield, gave a rather good presentation of the effect of biochar
531 addition on the vegetation production (Fig. 3c and d). It is interesting to note that the
532 application of OPB in HPS plots promoted a significant increase of fresh weight,
533 reaching values similar to those observed for MPS (Fig. 3d), while the vegetation cover
534 of OPB-HPS was five times lower than the unamended and amended MPS plots (Fig.
535 3c). This is explained by the presence of different plant species in OPB-HPS and MPS
536 plots (Table S2). In MPS, the plant species greatly covered the soil surface (high
537 vegetation coverage area), but their stem diameter and height were much smaller than
538 the species found in HPS plots. In OPB-treated HPS plots, few plants were found but
539 they displayed greater height and stem diameter, and less vegetation coverage.

540 The positive correlations obtained between plant data (number of species and
541 individuals, and fresh weight) with soil pH (Table S4), demonstrate that biochar is able
542 to enhance the properties of acidic soils, favouring the recovery of degraded polluted
543 soils due to the spill of heavy metals.

544 Changes in soil microbial community were also assessed in the biochar-amended and
545 untreated soils to inform about soil quality and biochar potential to restore soil
546 functionality. Monitoring microbial diversity by 16S rRNA gene NGS-based analyses
547 after 6, 12 and 20 months of biochar addition showed changes in the soil microbial
548 community structure, particularly in HPS plots after 6 months of soil amendment with

549 biochar. However, after 12 and 20 months, we did not find consistent phylum or order-
550 level responses to biochar amendments (Fig. 4 and Fig. S1), as the treated plots
551 showed higher similarity over control soils, as also reported by Song et al. (2017).
552 Similarly, Shannon and Simpson index values indicated that the addition of RH and OP
553 biochar solely promoted soil bacterial diversity in HPS at t_6 (Table 3). These findings
554 suggest that the type and dosages of biochar applied into HPS had a short-term effect
555 on the distribution of microbial communities, which was dissipated over time.

556 From the microbial community structure displayed in Figure 4, we drew the conclusion
557 that biochar addition significantly decreased the relative abundance of members of the
558 *Chloroflexi* phylum in HPS-amended plots at t_6 , probably due to changes in soil pH and
559 elements immobilization as *Chloroflexi* have preference for extreme environments (Soo
560 et al., 2009; Yabe et al., 2017). This phylum was mainly represented by the order
561 *Ktedonobacterales* (with 71% of relative abundance in C_HPS) from the class
562 *Ktedonobacteria*, which are filamentous bacteria that inhabit forest and garden soils at
563 low abundances, as well as extreme environments such as geothermal areas and
564 caves (Yabe et al., 2017). The relative abundance of the Ktedonobacterial community
565 (80%) in the HPS control plots abruptly declined (from 80% to 46%) with the
566 incorporation of biochar (Fig. S1a), probably due to changes in pH, which explains the
567 negative correlation between pH and *Chloroflexi* in HPS (Fig. 6a). This decline in
568 *Chloroflexi* abundances after biochar application was also previously reported by
569 several authors (Nielsen et al., 2014; Xu et al., 2014; Ali et al., 2019; Li et al., 2019).
570 However, Chen et al. (2019) showed an increase in *Chloroflexi* with the application of
571 10% of biochar to calcareous soils.

572 The relative abundance of *Firmicutes* was also reduced in HPS after 6 months of
573 biochar application. *Firmicutes* can adapt to low nutrient environments and thrive in
574 extreme conditions by forming spores (Bai et al., 2017; Li et al., 2014). Cole et al.
575 (2019) also found a decline in relative abundance of *Firmicutes* with biochar
576 application. However, Ali et al. (2019) reported an increase when a biochar produced
577 from sewage sludge was applied.

578 Conversely, the increase of *Proteobacteria* observed in HPS-amended samples at t_6 is
579 probably explained by their heterotrophic nature, as biochar increases soil carbon
580 content and nutrient conditions of poor-nutrient soils as HPS. Ali et al. (2019), Cole et
581 al. (2019), Li et al. (2019) and Su et al. (2015) also reported greater abundances of
582 *Proteobacteria* in amended soils than in control soils, obtaining good correlation
583 between *Proteobacteria* and labile C content.

584 In addition to *Proteobacteria*, the relative abundance of *Acidobacteria* slightly increased
585 after 6 months of biochar addition into HPS, as also reported by Cole et al. (2019).

586 However, Li et al. (2019) and Fan et al. (2020) reported a decrease of *Acidobacteria*
587 after biochar application, but Jenkins et al. (2017) found an increase of *Acidobacteria*
588 even in control soils without biochar treatment, indicating variations by weather
589 conditions. This is in accordance with our results for control HPS over time.

590 In this study, the relative abundance of *Planctomycetes*, *Bacteroidetes*,
591 *Gemmatimonadetes* and *Verrucomicrobia* did not depend on biochar application but on
592 soil type and seasonal changes. *Planctomycetes* were more abundant in MPS plots
593 than in HPS and their relative abundance varied with different seasons. Rice husk
594 biochar only slightly reduced *Planctomycetes* after 12 months of application into HPS,
595 which is in accordance with the findings of Noyce et al. (2016) when low pH soil was
596 amended with wood chips biochar. However, Ali et al. (2019) showed an increase in
597 *Planctomycetes* abundance in a contaminated-agricultural soil after the application of
598 rice straw biochar.

599 Chen et al. (2019) observed that the relative abundance of *Bacteroidetes* was higher in
600 the control soil than in the biochar-amended soil, attributing these changes to the initial
601 high pH and nutrient levels in the studied calcareous soils. However, Hu et al. (2014)
602 solely detected *Bacteroidetes* in the biochar amended soil. In this study, *Bacteroidetes*
603 were found in control and amended MPS plots, but not in HPS. It could be due to their
604 copiotrophic nature and capability for living in rhizosphere conditions (Shen et al.,
605 2018), as plant growth was solely observed in MPS plots at t_6 . Khodadad et al. (2011)
606 reported an increase of *Gemmatimonadetes* in soils with natural or added pyrogenic
607 carbon, suggesting an active role of these microorganisms in soil pyrogenic C
608 metabolism. We observed a small increase when RHB was applied, which could
609 indicate that this biochar could be more accessible than OPB for these group of
610 bacteria.

611 *Verrucomicrobia* was only found in MPS plots, suggesting that its presence was
612 dependent on the soil type, instead of biochar application. In fact, Chen et al. (2019)
613 observed that *Verrucomicrobia* was greater in control than in biochar-amended soils.
614 Nevertheless, Fan et al. (2020) reported an increase in *Verrucomicrobia* phylum in soils
615 amended with biochar.

616 *Actinobacteria* are possibly involved in the redistribution of consumed C or in the
617 degradation of more recalcitrant compounds (Blagodatskaya and Kuzyakov, 2008).
618 Cole et al. (2019) and Khodadad et al. (2011) reported an increase in *Actinobacteria* in
619 soils with natural or added pyrogenic carbon. However, Li et al. (2019) reported a
620 decrease in the relative abundance of *Actinobacteria* after biochar addition to soil. Our
621 results are more in accordance with this decline, particularly in RHB-amended MPS
622 plots at t_6 and in HPS plots at t_{12} . Jenkins et al. (2017) found an increase of

623 *Actinobacteria* even in control soils without biochar, indicating variations due to weather
624 conditions. In this study, the relative abundance of *Actinobacteria* also seemed to be
625 related to seasonal changes particularly in the case of HPS plots (Fig. 4).

626 In summary, the effects of biochar on soil bacterial communities are not unanimously
627 explained, as numerous other factors, such as soil type, pH, moisture and biochar
628 feedstock are likely to structure microbial communities (Chen et al., 2019; Jenkins et
629 al., 2017). In addition, environmental conditions and long-term biochar application may
630 have more influence in soil microbial communities than biochar types. It is worth
631 mentioning that this variability in soil microbial communities is mostly find in field
632 experiments, whereas in pot incubation experiments parameters are constrained (Hu et
633 al., 2014; Xu et al., 2014). Overall, the changes in the soil bacterial richness and
634 diversity after soil amendment application were correlated with changes in soil pH (Fig.
635 6), as the incorporation of biochar increased pH, and bacterial diversity, as well as
636 plant growth.

637

638 **5. Conclusions**

639 This field study conducted on polluted acidic soils has shown that the addition of
640 biochar allowed the recovery of plant cover and increased plant biodiversity,
641 particularly in moderately contaminated soils (MPS). Biochar application did not modify
642 soil CO₂ emissions, nor significantly increase enzymatic activity beyond the first six
643 months of biochar application, which points to a great stability of the tested olive pit and
644 rice husk biochar (OPB and RHB) and their ability to be used for carbon sequestration
645 in degraded soils. Findings from 16S rRNA gene next-generation sequencing revealed
646 that the incorporation of biochar modified the soil microbial community in the highly
647 polluted soil (HPS). Bacterial diversity was found to be site-specific as the properties
648 differed among the studied soils. We conclude that the application of biochar from crop
649 residues to trace-element polluted soils participated in soil conditioning, promoting
650 plant development, increasing bacterial diversity and soil carbon stabilization. This
651 suggested that the application of biochar is important in the ecological restoration of
652 these degraded soils. Our results showed that long-term experiments under field
653 conditions are essential in the quest to investigate the performance of biochar without
654 constraining environmental parameters, as seasonal changes were remarkable in this
655 study. This knowledge could help to fully understand the impact of biochar on global
656 nutrient cycles and on the recovery of soil ecological functions.

657

658 **Declaration of competing interest**

659 The authors declare that they have no conflict of interest to this work.

660

661 **Acknowledgments**

662 The Spanish Ministry of Economy, Industry and Competitiveness (MINEICO) and
663 AEI/FEDER UE are thanked for funding the project CGL2016-76498-R. P. Campos
664 thanks the “Fundación Tatiana Pérez de Guzmán el Bueno” for funding her PhD.
665 MINEICO is also acknowledged for funding the “Ramón y Cajal” post-doctoral contract
666 of José M. De la Rosa [grant number RYC2014-16338]. Ana Z. Miller acknowledges
667 the support from the Portuguese “Fundação para a Ciência e a Tecnologia” (FCT)
668 [grant number CEECIND/01147/2017]. S. Prats thanks the Portuguese FCT for his
669 research contract [CDL-CTTRI-88-ARH/2018 REF.-138-88-ARH/2018], funded in the
670 scope of Law 57/2017 and for the financial support to CESAM (UID/AMB/50017/2019).

671

672 **References**

- 673 AEMET. 2020. Available at: <https://datosclima.es/Aemet2013/Tempestad2013.php>
674 (accessed on 20 May 2020).
- 675 Ali, N., Khan, S., Li, Y., Zheng, N., Yao, H., 2019. Influence of biochars on the
676 accessibility of organochlorine pesticides and microbial community in contaminated
677 soils. *Science of the Total Environment* 647, 551–560.
- 678 Arias, M.E., González-Pérez, J.A., González-Vila, F.J., Ball A.S., 2005. Soil health-a
679 new challenge for microbiologists and chemists. *International Microbiology* 8, 13–21.
- 680 Bai, R., Wang, J.-T., Deng, Y., He, J.-Z., Feng, K., Zhang, L.-M., 2017. Microbial
681 community and functional structure significantly varied among distinct types of
682 paddy soils but responded differently along gradients of soil depth layers. *Frontiers*
683 *in Microbiology* 8, 945, 1–16.
- 684 Bamminger, C., Marschner, B., Juschke, E., 2014. An incubation study on the stability
685 and biological effects of pyrogenic and hydrothermal biochar in two soils. *European*
686 *Journal of Soil Science* 65, 72–82.
- 687 Beesley, L., Moreno-Jiménez, E., Gomez-Eyles, J.L., 2010. Effects of biochar and
688 greenwaste compost amendments on mobility, bioavailability and toxicity of
689 inorganic and organic contaminants in a multi-element polluted soil. *Environmental*
690 *Pollution* 158, 2282–2287.
- 691 Blagodatskaya, E., Kuzyakov, Y., 2008. Mechanisms of real and apparent priming
692 effects and their dependence on soil microbial biomass and community structure:
693 critical review. *Biology and Fertility of Soils* 45, 115–131.
- 694 Bolyen, E., Rideout, J.R., Dillon, M.R. et al. 2019. Reproducible, interactive, scalable
695 and extensible microbiome data science using QIIME 2. *Nature Biotechnology* 37,
696 852–857.

- 697 Cabrera, F., Clemente, L., Díaz Barrientos, E., López, R., Murillo, J.M., 1999. Heavy
698 metal pollution of soils affected by the Guadiamar toxic flood. *Science of the Total*
699 *Environment* 242, 117–129.
- 700 Campos, P., De la Rosa, J.M., 2020. Assessing the effects of biochar on the
701 immobilization of trace elements and plant development in a naturally contaminated
702 soil. *Sustainability*, 12, 6025–6044.
- 703 Campos, P., Miller, A.Z., Knicker, H., Costa-Pereira, M.F., Merino, A., De la Rosa, J.M.,
704 2020. Chemical, physical and morphological properties of biochars produced from
705 agricultural residues: Implications for their use as soil amendment. *Waste*
706 *Management* 105, 256–267.
- 707 Chen, J., Lee, X., Tang, Y., Zhang, Q., 2019. Long-term effects of biochar amendment
708 on rhizosphere and bulk soil microbial communities in a karst region, southwest
709 China. *Applied Soil Ecology* 140, 126–134.
- 710 Cole, E.J., Zandvakili, O.R., Blanchard, J., Xing, B., Hashemi, M., Etemadi, F., 2019.
711 Investigating responses of soil bacterial community composition to hardwood
712 biochar amendment using high-throughput PCR sequencing. *Applied Soil Ecology*
713 136, 80–85.
- 714 Dick, R.P., Breakquill, D., Turco, R., 1996. Soil enzyme activities and biodiversity
715 measurements as integrating biological indicators. In Doran, J.W., Jones, A.J.,
716 *Handbook of Methods for Assessment of Soil Quality*. Soil Science Society of
717 America Specific Publications, Madison, WI, 242–272.
- 718 De la Rosa, J.M., Paneque, M., Miller, A.Z., Knicker, H., 2014. Relating physical and
719 chemical properties of four different biochars and their application rate to biomass
720 production of *Lolium perenne* on a Calcic Cambisol during a pot experiment of 79
721 days. *Science of the Total Environment* 499, 175–184.
- 722 De la Rosa, J.M., Rosado, M., Paneque, M., Miller, A.Z., Knicker, H., 2018. Effects of
723 aging under field conditions on biochar structure and composition: Implications for
724 biochar stability in soils. *Science of the Total Environment* 613-614, 969–976.
- 725 De la Rosa, J.M., Sánchez-Martín, A.M., Campos, P., Miller, A.Z., 2019. Effect of
726 pyrolysis conditions on the total contents of polycyclic aromatic hydrocarbons in
727 biochars produced from organic residues: Assessment of their hazard potential.
728 *Science of the Total Environment* 667, 578–585.
- 729 EEA; 2007. *Progress in Management of Contaminated Sites*. CSI 015. Copenhagen,
730 Denmark: European Environmental Agency.
- 731 Eivazi, F., Tabatai, M., 1990. Factors affecting glucosidase and galactosidase in soils.
732 *Soil Biology and Biochemistry* 22, 145–152.

- 733 Elzobair, K.A., Stromberger, M.E., Ippolito, J.A., Lentz, R.D., 2016. Contrasting effects
734 of biochar versus manure on soil microbial communities and enzyme activities in an
735 Aridisol. *Chemosphere* 142, 145–152.
- 736 Fan, S., Zuo, J., Fond, H., 2020. Changes in Soil Properties and Bacterial Community
737 composition with Biochar Amendment after Six Years. *Agronomy* 10, 746, 1–15.
- 738 Ferraz de Almeida, R., Rezende Naves, E., Pinheiro da Mota, R., 2015. Soil quality:
739 Enzymatic activity of soil β -glucosidase. *Global Journal of Agricultural Research and*
740 *Reviews* 3, 146–150.
- 741 Frankenberger, W., Johanson, J., 1982. Effect of pH On Enzyme Stability in Soils. *Soil*
742 *Biology and Biochemistry* 14, 433–437.
- 743 Gascó, G., Paz-Ferreiro, J., Cely, P., Plaza, C., Méndez, A., 2016. Influence of pig
744 manure and its biochar on soil CO₂ emissions and soil enzymes. *Ecological*
745 *Engineering* 95, 19–24.
- 746 Günal, E., Erdem, H., Demirbaş, A., 2018. Effects of three biochar types on activity of
747 β -glucosidase enzyme in two agricultural soils of different textures. *Archives of*
748 *Agronomy and Soil Science*, 64, 14, 1963–1974.
- 749 Hagemann, N., Schmidt, H.-P., Kagi, R., Bohler, M., Sigmund, G., Maccagnan, A.,
750 McArdell, C.S., Bucheli, T.D., 2020. Wood-based activated biochar to eliminate
751 organic micropollutants from biologically treated wastewater. *Science of the Total*
752 *Environment* 730, 138417, 1–11.
- 753 Hagemann, N., Spokas, K., Schmidt, H.-P., Kägi, R., Böhler, M.A., Bucheli, T.D., 2018.
754 Activated Carbon, Biochar and Charcoal: Linkages and Synergies across Pyrogenic
755 Carbon's ABCs. *Water* 10, 182, 1–19.
- 756 Han, G., Lan, J., Chen, Q., Yu, C., Bie, S., 2017. Response of soil microbial community
757 to application of biochar in cotton soils with different continuous cropping years.
758 *Scientific Reports* 7, 10184, 1–11.
- 759 Hu, L., Cao, L., Zhang, R., 2014. Bacterial and fungal taxon changes in soil microbial
760 community composition induced by short-term biochar amendment in red oxidized
761 loam soil. *World Journal of Microbiology and Biotechnology* 30, 1085–1092.
- 762 IUSS Working Group WRB, 2015. World Reference Base for Soil Resources 2014,
763 update 2015. International soil classification system for naming soils and creating
764 legends for soil maps. *World Soil Resources Reports* 106. FAO, Rome, 1–203.
- 765 Jenkins, J. R., Viger, M., Arnold, E.C., Harris, Z.M., Ventura, M., Miglietta, F., Girardin,
766 C., Edwards, R.J., Rumpel, C., Fornasier, F., Zavalloni, C., Tonon, G., Alberti, G.,
767 Taylor, G., 2017. Biochar alters the soil microbiome and soil function: results of next-
768 generation amplicon sequencing across Europe. *Global Change Biology Bioenergy*
769 9, 591–612.

- 770 Jian, L.-L., Han, G.-M., Lan, Y., Liu, S.-N., Gao, J.-P., Yang, X., Meng, J., Chen, W.-F.,
771 2017. Corn cob biochar increases soil culturable bacterial abundance without
772 enhancing their capacities in utilizing carbon sources in Biolog Eco-plates. *Journal*
773 *of Integrative Agriculture* 16, 712–724.
- 774 Kammann, C., Ratering, S., Eckhard, C., Müller, C., 2012. Biochar and hydrochar
775 effects on greenhouse gas (carbon dioxide, nitrous oxide, methane) fluxes from
776 soils. *Journal of Environmental Quality* 41, 1052–1066.
- 777 Karer, J., Wawra, A., Zehetner, F., Dunst, G., Wagner, M., Pavel, P.-B., Puschenreiter,
778 M., Friesl-Hanl, W., Soja, G., 2015. Effects of Biochars and Compost Mixtures and
779 Inorganic Additives on Immobilisation of Heavy Metals in Contaminated Soils.
780 *Water, Air, & Soil Pollution* 226, 342, 1–12.
- 781 Khodadad, C.L.M., Zimmerman, A.R., Green, S.J., Uthandi, S., Foster, J.S., 2011.
782 Taxa-specific changes in soil microbial community composition induced by
783 pyrogenic carbon amendments. *Soil Biology and Biochemistry* 43, 385–392.
- 784 Knicker, H., Hilscher, A., De la Rosa, J.M., González-Pérez, J.A., González-Vila, F.J.,
785 2013. Modification of biomarkers in pyrogenic organic matter during the initial phase
786 of charcoal biodegradation in soils. *Geoderma* 197–198, 43–50.
- 787 Kolb, S.E., Fermanich, K.J., Dornbush, M.E., 2009. Effect of charcoal quantity on
788 microbial biomass and activity in temperate soils. *Soil Science Society of America*
789 *Journal* 73, 1173–1181.
- 790 Kumar A, Joseph S, Tsechansky L, Privat, K., Schreiter, I.J., Schüth, C., Graber, E.R.,
791 2018. Biochar aging in contaminated soil promotes Zn immobilization due to
792 changes in biochar surface structural and chemical properties. *Science of the Total*
793 *Environment* 626, 953–961.
- 794 Kuzyakov, Y., Bogomolova, I., Glaser, B., 2014. Biochar stability in soil: Decomposition
795 during eight years and transformation as assessed by compound-specific ^{14}C
796 analysis. *Soil Biology and Biochemistry* 70, 229–236.
- 797 Kuzyakov, Y., Subbotina, I., Chen, H., Bogomolova, I., Xu, X., 2009. Black carbon
798 decomposition and incorporation into soil microbial biomass estimated by ^{14}C
799 labelling. *Soil Biology and Biochemistry* 41, 210–219.
- 800 Lehmann, J., Rillig, M.C., Thies, J., Masiello, C.A., Hockaday, W.C., Crowley, D., 2011.
801 Biochar effects on soil biota - a review. *Soil Biology and Biochemistry* 43, 1812–
802 1836.
- 803 Li, C., Yan, K., Tang, L., Jia, Z., Li, Y., 2014. Change in deep soil microbial
804 communities due to long-term fertilization. *Soil Biology and Biochemistry* 75, 264-
805 272.

- 806 Li, Y., Yang, Y., Shen, F., Tian, D., Zeng, Y., Yang, G., Zhang, Y., Deng, S., 2019.
807 Partitioning biochar properties to elucidate their contributions to bacterial and fungal
808 community composition of purple soil. *Science of the Total Environment* 648, 1333–
809 1341.
- 810 Luo, Y., Durenkamp, M., De Nobili, M., Lin, Q., Devonshire, B.J., Brookes, P.C., 2013.
811 Microbial biomass growth, following incorporation of biochars produced at 350 °C or
812 700 °C, in a silty-clay loam soil of high and low pH. *Soil Biology and Biochemistry*
813 57, 513–523.
- 814 Martín-Peinado, F. J., Romero-Freire, A., García-Fernández, I., Sierra Aragón, M.,
815 Ortiz-Bernad, I., Simón Torres, M., 2015. Long-term contamination in a recovered
816 area affected by a mining spill. *Science of the Total Environment* 514, 219–223.
- 817 Nielsen, S., Minchin, T., Kimber, S., van Zwieten, L., Gilbert, J., Munroe, P., Joseph,
818 S., Thomas, T., 2014. Comparative analysis of the microbial communities in
819 agricultural soil amended with enhanced biochars or traditional fertilisers.
820 *Agriculture, Ecosystems and Environment* 191, 73–82.
- 821 Noyce, G.L., Winsborough, C., Fulthorpe, R., Basiliko, N., 2016. The microbiomes and
822 metagenomes of forest biochars. *Scientific Reports* 6, 26425, 1–12.
- 823 Oustriere, N., Marchand, L., Lottier, N., Motelica, M., Mench, M., 2017. Long-term Cu
824 stabilization and biomass yields of Giant reed and poplar after adding a biochar,
825 alone or with iron grit, into a contaminated soil from a wood preservation site.
826 *Science of the Total Environment* 579, 620–627.
- 827 Paz-Ferreiro, J., Gascó, G., Gutiérrez, B., Méndez, A., 2012. Soil biochemical activities
828 and the geometric mean of enzyme activities after application of sewage sludge and
829 sewage sludge biochar to soil. *Biology and Fertility of Soils* 48, 511–517.
- 830 Phongthep, H., Jiranut, W., Tanakit, S., Sathaporn, J., Sukanya, T., 2017. Soil
831 respiration in rubber tree plantation applied with biochar. *Research Journal of*
832 *Chemistry and Environment* 21 (10), 27–34.
- 833 Shen, G., Zhang, S., Liu, X., Jiang, Q., Ding, W., 2018. Soil acidification amendments
834 change the rhizosphere bacterial community of tobacco in a bacterial wilt affected
835 field. *Applied Microbiology and Biotechnology* 102, 9781–9791.
- 836 Shen, Z., Hou, D., Jin, F., Shi, J., Fan, X., Tsang, D.C.W., Alessi, D.S., 2019. Effect of
837 production temperature on lead removal mechanisms by rice straw biochars.
838 *Science of the Total Environment* 655, 751–758.
- 839 Sohi, S.P., Krull, E., Lopez-Capel, E., Bol, R., 2010. A review of biochar and its use
840 and function in soil. *Advances in Agronomy* 105, 47–82.
- 841 Song, Y., Bian, Y., Wang, F., Xu, M., Ni, N., Yang, X., Gu, C., Jiang, X., 2017. Dynamic
842 Effects of Biochar on the Bacterial Community Structure in Soil Contaminated with

- 843 Polycyclic Aromatic Hydrocarbons. *Journal of Agricultural and Food Chemistry* 65,
844 6789–6796.
- 845 Soo, R.M., Wood, S.A., Grzymiski, J.J., McDonald, I.R., Cary, S.C., 2009. Microbial
846 biodiversity of thermophilic communities in hot mineral soils of Tramway Ridge,
847 Mount Erebus, Antarctica. *Environmental Microbiology* 11, 715–728.
- 848 Su, P., Lou, J., Brookes, P.C., Luo, Y., He, Y., Xu, J., 2015. Taxon-specific responses
849 of soil microbial communities to different soil priming effects induced by addition of
850 plant residues and their biochars. *Journal of Soils and Sediments* 17 (3), 674–684.
- 851 Sun, Z., Bruun, E.W., Arthur, E., Wollesen de Jonge, L., Moldrup, P., Hauggaard-
852 Nielsen, H., Elsgaard, L., 2014. Effect of biochar on aerobic processes, enzyme
853 activity and crop yields in two sandy loam soils. *Biology and Fertility of Soils* 50,
854 1087–1097.
- 855 Tabatabai, M.A., 1982. Soil enzymes. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.),
856 Method of Soil Analysis, Part 2. Chemical and Microbiological Properties. American
857 Society of Agronomy, Madison, 903–948.
- 858 Tack, F., Rinklebe, J., Ok, Y.S., 2018. Interactions between biochar and trace elements
859 in the environment. *Science of The Total Environment* 649, 792–793.
- 860 Trevors, J.T., 1984. Dehydrogenase activity in soil: a comparison between the INT and
861 TTC assay. *Soil Biology and Biochemistry* 16 (6), 673–674.
- 862 Uchimiya, M., Bannon, D.H., Wartelle, L.H., Lima, I.M., Klasson, K.T., 2012. Lead
863 retention by broiler litter biochars in small arms range soil: impact of pyrolysis
864 temperature. *Journal of Agricultural and Food Chemistry* 60, 5035–5044.
- 865 Uchimiya, M., Klasson, K.T., Wartelle, L.H., Lima, I.M., 2011. Influence of soil
866 properties on heavy metal sequestration by biochar amendment: 2. Copper
867 desorption isotherms. *Chemosphere* 82, 1438–1447.
- 868 Xu, H.J., Wang, X.H., Li, H., Yao, H.-Y.Y., Su, J.Q., Zhu, Y.G., 2014. Biochar impacts
869 soil microbial community composition and nitrogen cycling in an acidic soil planted
870 with rape. *Environmental Science Technology* 48, 9391–9399.
- 871 Xu, N., Tan, G., Wang, H., Gai, X., 2016. Effect of biochar additions to soil on nitrogen
872 leaching, microbial biomass and bacterial community structure. *European Journal of*
873 *Soil Biology* 74, 1–8.
- 874 Yabe, S., Sakai, Y., Abe, K., Yokota, A., 2017. Diversity of Ktedonobacteria with
875 Actinomycetes-Like morphology in terrestrial environments. *Microbes and*
876 *Environment* 32, 61–70.
- 877 Zakrzewski, M., Proietti, C., Ellis, J.J., Hasan, S., Brion, M.J., Berger, B., Krause, L.,
878 2017. Calypso: a user-friendly web-server for mining and visualizing microbiome-
879 environment interactions. *Bioinformatics* 33, 782–783.

Journal Pre-proof

881 **FIGURE CAPTIONS:**

882

883 **Fig. 1.** Location of the field experiment, Aznalcóllar mine and Guadiamar Green
884 Corridor.

885

886 **Fig. 2.** Enzymatic activities in control and biochar amended soils. a) β -Glucosidase
887 activity in Highly Polluted Soil, b) β -Glucosidase activity in Moderately Polluted Soil, c)
888 Dehydrogenase activity in Highly Polluted Soil and d) Dehydrogenase activity in
889 Moderately Polluted Soil. Different letters for each sampling period indicate significant
890 differences between treatments ($p < 0.05$) based on one-way ANOVA test followed by
891 the Tukey HSD test.

892

893 **Fig. 3.** a) Number of different vegetation species per plot at t_{12} . b) Number of plants per
894 m^2 in control and biochar amended plots at t_{12} . c) Average of vegetation cover (%) per
895 plot. d) Average fresh weight of plants per plot. Different letters indicate significant
896 differences between treatments ($p < 0.05$) based on one-way ANOVA test followed by
897 the Tukey HSD test.

898

899 **Fig. 4.** Relative abundance of the OTUs at the phylum level in the control (C_HPS and
900 C_MPS) and biochar-amended soils (RHB_HPS, OPB_HPS, RHB_MPS and
901 OPB_MPS) at: a) t_6 (6 months after biochar application into soils); b) t_{12} (12 months),
902 and c) t_{20} (20 months after biochar application).

903 **Fig. 5.** Assessment of bacterial diversity using principal component analysis (PCA) of
904 highly polluted soil (HPS) and moderately polluted soil (MPS) at: a) t_6 ; b) t_{12} , and c) t_{20} .
905 RHB and OPB represent biochars derived from rice husk and olive pit, respectively,
906 whereas C_HPS and C_MPS correspond to control soils.

907

908 **Fig. 6.** Correlation heatmaps between soil physicochemical properties (pH, WHC, TC,
909 moisture, soil respiration, β -Glucosidase, Dehydrogenase) and bacterial abundance for
910 a) t_6 ; b) t_{12} , and c) t_{20} .

Table 1. pH, total carbon (TC), total nitrogen (TN) and trace elements contents of rice husk biochar (RHB), olive pit biochar (OPB), highly polluted soil (HPS) and moderately polluted soil (MPS). Values represent means ($n = 3$) and standard deviation.

		pH	TC	TN	Ba	Cd	Cu	Fe	Ni	Pb	Sr	Zn
			(g kg ⁻¹)	(g kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)
Biochars	RHB	10.17± 0.34	537± 1.0	1.6± 0.9	7.3	0.05	35.0	1224.2	8.5	1.7	11.6	42.6
	OPB	9.34± 0.19	927± 2.0	5.1± 2.4	<LOQ*	<LOQ	5.9	<LOQ	<LOQ	0.4	4.4	<LOQ
Soils	HPS	3.85± 0.14	7.2± 0.1	0.6± 0.1	47.1	1.28	240.6	53023.3	15.6	569.0	53.7	249.3
	MPS	4.82± 0.13	9.0± 0.6	0.8± 0.1	93.3	1.56	215.5	36945.7	15.6	156.5	38.6	293.5

*<LOQ: below limit of quantification.


Table 2. Changes in soil characteristics, soil total carbon (TC) and soil respiration during the field experiment (t_1 : 1 month-spring, t_6 : 6 months-autumn, t_{12} : 12 months-spring, t_{20} : 20 months-autumn).


Sample	pH			WHC (%)			TC (g kg ⁻¹)			Soil respiration ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)		
	t_6	t_{12}	t_{20}	t_6	t_{12}	t_{20}	t_6	t_{12}	t_{20}	t_6	t_{12}	t_{20}
C_HPS	3.57±0.13a	3.59±0.08a	3.52±0.20a	20±7a	14±5a	32±2a	7.9±0.6a	9.6±1.3a	12.3±2.7a	1.0±0.2a	1.0±0.2a	0.9±0.2a
RHB_HPS	3.64±0.18a	3.68±0.21a	3.77±0.17a	23±1a	20±2ab	26±4a	12.9±4.8a	14.9±4.7ab	16.7±1.5a	1.0±0.3a	0.8±0.2a	0.6±0.3a
OPB_HPS	3.63±0.03a	3.64±0.04a	3.69±0.17a	23±5a	17±5ab	30±4a	13.6±8a	16.6±4.5b	18.4±6.1a	1.2±0.4a	0.9±0.2a	1.2±0.6a
C_MPS	4.18±0.21a	4.8±0.10b	4.87±0.20b	32±2b	23±5ab	47±6ab	12.4±3.5a	9.5±0.3a	21.7±1.3a	2.9±0.2b	1.2±0.8a	3.9±0.2b
RHB_MPS	4.75±0.01b	5.02±0.22b	5.07±0.18b	39±1b	25±8b	52±2b	12.2±1.7a	14.3±3.2ab	19.9±9.2a	2.7±0.3b	1.3±0.7a	3.3±0.4b
OPB_MPS	4.74±0.54b	4.86±0.50b	5.11±0.20b	38±7b	19±7ab	67±3c	17.0±7.9a	12.7±0.3ab	20.3±10.2a	3.1±0.3b	0.5±0.1a	3.7±1.0b


WHC: Water holding capacity. Different letters within a column indicate significant differences between treatments ($p < 0.05$) based on a one-way ANOVA test followed by the Tukey HSD test.


Table 3. Alpha-diversity indices of microbial community structure in the unamended and biochar-amended HPS and MPS samples. The diversity indices (Shannon and Simpson index) and richness index (Chao1 and OTUs) were determined at 97% sequence similarity.


Sample ID	No. OTUs	Alpha-diversity								
		Shannon			Simpson			Chao1		
		t ₆	t ₁₂	t ₂₀	t ₆	t ₁₂	t ₂₀	t ₆	t ₁₂	t ₂₀
C_HPS	426	2.00	3.23	3.84	0.71	0.91	0.96	52	142	232
RHB_HPS	468	2.76	3.42	3.33	0.86	0.91	0.91	76	193	199
OPB_HPS	488	2.62	2.84	4.24	0.83	0.86	0.97	73	173	242
C_MPS	1475	4.47	5.33	4.72	0.97	0.99	0.98	291	702	482
RHB_MPS	1493	4.62	5.20	4.73	0.98	0.99	0.98	359	703	431
OPB_MPS	1530	4.61	5.18	5.16	0.98	0.99	0.99	341	646	543














Journal Pre-proof

Highlights:

- Biochar was applied in moderately and highly polluted soils under field conditions.
- Biochar application increased bacterial and plant diversity in highly polluted soil.
- Bacterial diversity slightly changed in moderately polluted soil with biochar.
- Soil pH was key factor of bacterial community structure.
- Seasonal changes affected microbial community, soil respiration and vegetation.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof