



Identifying knowledge gaps in understanding the effects of selective serotonin reuptake inhibitors (SSRIs) on fish behaviour

Carla S.S. Ferreira^{a,*}, Sandra C. Soares^{b,c}, Peter Kille^d, Miguel Oliveira^a

^a Centre for Marine and Environmental Studies (CESAM), Department of Biology, University of Aveiro, 3810-193, Aveiro, Portugal

^b William James Center for Research (WJRC), Department of Education and Psychology, University of Aveiro, 3810-193, Aveiro, Portugal


^c Center for Health Technology and Services Research (CINTESIS), Department of Education and Psychology, University of Aveiro, 3810-193, Aveiro, Portugal

^d School of Biosciences, Cardiff University, Cardiff, CF10 3AX, UK

HIGHLIGHTS

- The reported effects of SSRIs on fish fitness-related behaviours were reviewed.
- Most of the performed studies lack environmental relevance and consistency in the reported effects.
- Little is known on how personality influences experimental outcomes.
- Antidepressants exposure may lead to alteration on personality traits.
- Research gaps are highlighted.

GRAPHICAL ABSTRACT



ARTICLE INFO

Handling Editor: James Lazorchak

Keywords:
 Aquatic pollution
 Antidepressants
 Behaviour
 Personality

ABSTRACT

Selective serotonin reuptake inhibitors (SSRIs) are a class of antidepressants increasingly prescribed to treat patients with clinical depression. As a result of the significant negative impact of the COVID-19 pandemic on the population's mental health, its consumption is expected to increase even more. The high consumption of these substances leads to their environmental dissemination, with evidence of their ability to compromise molecular, biochemical, physiological, and behavioural endpoints in non-target organisms. This study aimed to provide a critical review of the current knowledge regarding the effects of SSRI antidepressants on fish ecologically relevant behaviours and personality-dependent traits. A literature review shows limited data concerning the impact of fish personality on their responses to contaminants and how such responses could be influenced by SSRIs. This lack of information may be attributable to a lack of widely adopted standardized protocols for evaluating behavioural responses in fish. The existing studies examining the effects of SSRIs across various biological levels overlook the intra-specific variations in behaviour and physiology associated with different personality patterns or coping styles. Consequently, some effects may remain undetected, such as variations in coping styles and the capacity to handle environmental stressors. This oversight could potentially result in long-term effects with ecological implications.

Data support the need for more studies to understand the impact of SSRIs on personality-dependent traits and how they may impair fitness-related behaviours. Given the considerable cross-species similarity in the personality dimensions, the collected data may allow new insights into the correlation between personality and animal fitness.

* Corresponding author.

E-mail address: csfia@ua.pt (C.S.S. Ferreira).

<https://doi.org/10.1016/j.chemosphere.2023.139124>

Received 2 March 2023; Received in revised form 1 June 2023; Accepted 2 June 2023

Available online 5 June 2023

0045-6535/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Pharmaceutical consumption is a global-scale practice, projected to increase further due to population growth and ageing (Saaristo et al., 2019; Moreira et al., 2022). This has spurred intensive research into the development of new and more potent drugs for preventive and symptomatic treatment of human ailments, including mental disorders (Mezzelani et al., 2018; Fekadu et al., 2019). Upon administration, pharmaceuticals undergo metabolism, exert their intended therapeutic effects by targeting specific biological pathways, and are excreted through urine and/or faeces, either in their original form or as metabolites, thereby entering the sewage system (Branchet et al., 2021; Madikizelaa et al., 2020; Klatte et al., 2017; Sehonova et al., 2018). However, current sewage treatment processes often fail to effectively remove these substances, leading to their dissemination in the aquatic environment (Saaristo et al., 2019; Madikizelaa et al., 2020; Klatte et al., 2017; Sehonova et al., 2018; Desbiolles et al., 2018; Hossain et al., 2021). This emergence of pharmaceuticals as environmental contaminants poses a growing challenge with potential risks to ecosystems and human health (Saaristo et al., 2019; Desbiolles et al., 2018; Puckowski et al., 2016). Consequently, pharmaceuticals have garnered increasing concern as emerging environmental contaminants (Moreira et al., 2022; Desbiolles et al., 2018; Hossain et al., 2021; Puckowski et al., 2016). Despite being detected at relatively low concentrations, typically ranging from nanograms per litre (ng/L) to micrograms per litre ($\mu\text{g/L}$), their continuous release into the environment requires careful attention (Puckowski et al., 2016). Even at these low concentrations, they can induce harmful effects on aquatic organisms due to the high degree of structural similarity between the primary targets of SSRIs in fish and humans (Hossain et al., 2021; Puckowski et al., 2016; Nguyen et al., 2021; Burkina et al., 2015). Pharmaceuticals have already been detected in aquatic environments at concentrations exceeding those known to induce toxic effects (Nguyen et al., 2021). Overall, available data indicate that the risks posed by pharmaceuticals to fish are influenced by multiple factors, including the nature and concentration of the drug and characteristics of the organisms (such as species, life stage, sex, and previous exposure to other environmental contaminants) (Burkina et al., 2015; Huang et al., 2019, 2020; Bachour et al., 2020; Sehonova et al., 2019).

Selective serotonin reuptake inhibitor (SSRI) antidepressants are one of the major classes of pharmaceuticals, widely and increasingly prescribed in the last decades to treat patients diagnosed with clinical depression, also being used to treat obsessive-compulsive disorder, panic disorders, social phobias, and attention-deficit disorder (Mole and Brooks, 2019; Silva et al., 2012). These pharmaceuticals have received particular attention as they can be particularly potent and hazardous since they are specifically designed to modulate human behaviour (Huang et al., 2019, 2020; Mole and Brooks, 2019). Therefore, they may induce behavioural alterations in non-target organisms, like fish, with severe fitness consequences (Martin et al., 2021). Several studies have already demonstrated that sublethal exposure to SSRIs can significantly alter fish behaviour, physiology, and gene expression (Hedgespeth et al., 2014; Kellner et al., 2015; Bisesi et al., 2016; Pelli and Connaughton, 2015; Weinberger and Klaper, 2014; Yang et al., 2021; Menon et al., 2020). This class of antidepressants has often been detected in water systems (Fekadu et al., 2019; Burkina et al., 2015; Menon et al., 2020; Fernandez-Rubio et al., 2019; Silva et al., 2015; Sumpter and Margiotta-Casaluci, 2022), and their environmental levels are expected to increase significantly due to the negative impact of the COVID-19 pandemic on mental health (Singh et al., 2022). SSRIs exert their therapeutic effect by blocking the serotonin reuptake transporter (SERT) in the presynaptic cell, thus inducing an increase in serotonin levels within the central nervous system (CNS) (Salahinejad et al., 2022; Kellner and Olsén, 2020; McDonald, 2017). Environmental risk assessments focusing on ecologically relevant sublethal endpoints related to chronic exposures, such as altered behaviour, have been considered a suitable and

reliable approach to characterize the ecotoxicological potential of these biologically active chemicals, ubiquitous at low concentrations (Mezzelani et al., 2018; Huang et al., 2019; Mole and Brooks, 2019; Martin et al., 2021).

Behaviour modification, resulting from complex and interconnected levels of biological organization (Saaristo et al., 2019; Martin et al., 2021; Bertram et al., 2022; Ford et al., 2021), has emerged as a sensitive indicator of environmental contaminants, carrying ecological significance. However, exposure to stress can elicit distinct intraspecific behavioural and physiological responses to environmental and toxicological stressors, known as coping styles (Demin et al., 2019; Wong et al., 2019; Bergmuller, 2020; Yuan et al., 2018). This individual sensitivity within a species can influence susceptibility to diseases and environmental contaminants (Demin et al., 2019; Wong et al., 2019). Extensive research involving humans and non-human animals has revealed variations in coping abilities when confronted with environmental challenges (Koolhaas, 2008; Xin et al., 2017; Santarnecchi et al., 2018). Humans, for instance, exhibit diverse coping styles when faced with stressful situations in their daily lives (Xin et al., 2017), representing unique sets of mental and behavioural strategies employed to navigate stress and traumatic experiences (Santarnecchi et al., 2018). Notably, individuals differ significantly in their responses to stress, with some displaying maladaptive reactions while others exhibit resilience (Xin et al., 2017), suggesting varying susceptibility to stress and, consequently, to diseases. Avoidance-based coping styles have been associated with a predisposition to developing psychiatric disorders like post-traumatic stress disorder (PTSD), anxiety, and major depression, whereas problem-oriented coping skills have shown positive correlations with well-being and a higher quality of life (Santarnecchi et al., 2018).

Similarly, it is recognized that organisms from many species, ranging from insects to humans, exhibit different coping styles, differing from their conspecifics in stress responsiveness to the environment when exposed to environmental stressors (Koolhaas, 2008; Demin et al., 2019; Wong et al., 2019). These individual coping styles occur within a population as a continuum axis between two extreme phenotypes, called bold (proactive) and shy (reactive), along with intermediate phenotypes (Ferrari et al., 2020; Tudorache et al., 2013, 2018). Nonetheless, there is a considerable lack of information about how coping styles/personality influence organisms' response to contaminant-induced stress, fitness, and health and vice versa. This can be studied in organisms like zebrafish (*Danio rerio*) that display two well-defined and distinct personality patterns. An additional advantage of using this species is its high genetic homology and significant conservation of monoaminergic systems with humans (Adhish and Manjubala, 2023; Xie et al., 2022; Gould et al., 2021).

Thus, this review aimed to provide an overview of the current knowledge on the sub-lethal effects induced by SSRIs on fish, focusing on behaviour and fish personality/coping styles, tested under ecologically relevant scenarios, and provide a critical discussion on the relevance of these findings.

2. SSRIs as emerging contaminants

The consumption of antidepressants significantly increased in the last few years, particularly in Europe. In 2000, consumption did not exceed 20 Defined Daily Doses (DDD)/1000 inhabitants/day, while in 2018 it almost reached 110 DDD/1000 inhabitants/day (Estrela et al., 2020). The recent pandemic of COVID-19 had a considerable impact on the population's mental health (Singh et al., 2022). According to the Organization for Economic Co-operation and Development (OECD), the incidence of anxiety and depression increased significantly from March 2020 onwards (Fig. 1) (Organization for Economic Co).

Therefore, the prescription and consumption of antidepressants are expected to increase even more, particularly of SSRIs, the most prescribed antidepressant drugs (Estrela et al., 2020; Zacarías et al., 2021),




Fig. 1. Prevalence of depression conditions in different countries, before the COVID-19 pandemic and in early 2020. (Data source: OECD, 2021).

representing an environmental and public health concern (Diaz-Camal et al., 2022), as some of these drugs are detected in drinking water (Zacarias et al., 2021).

Following oral ingestion, SSRIs are primarily subjected to hepatic metabolism, exerting their therapeutic effect by blocking the serotonin reuptake transporter (SERT) in the presynaptic neuron and inducing an increase in serotonin (5-HT; 5-hydroxytryptamine) extracellular concentrations within the CNS, thus leading to the relief of the symptomatology associated with depression and anxiety conditions (Sumpter and Margiotta-Casaluci, 2022; Salahinejad et al., 2022). These pharmaceuticals may be eliminated through urine and/or faeces as parent compounds and/or as their metabolites (Silva et al., 2012), resulting in their release to sewage systems (Sehonova et al., 2018). Due to incomplete removal during wastewater treatment, these active substances reach and spread along environmental matrices, such as water systems (Sumpter and Margiotta-Casaluci, 2022; Hughes et al., 2012) and, therefore, are frequently detected in surface waters (Mole and Brooks, 2019; Salahinejad et al., 2022). Once in the environment, they exhibit relatively low susceptibility to environmental degradation (e.g., hydrolysis, photolysis, biodegradation) which is translated.

Into their environmental persistence (Silva et al., 2012). They remain pharmaceutically active with the ability to cause detrimental effects on non-target organisms, even at low concentrations, due to the evolutionary conservation of the serotonergic system across vertebrates, along with their bioaccumulation potential (Silva et al., 2012; Martin et al., 2021; Sumpter and Margiotta-Casaluci, 2022; Salahinejad et al., 2022; Gould et al., 2021; Hughes et al., 2012; Kaushik and Thomas, 2019; Grabicova et al., 2017), associated with their lipophilic nature. pH plays an important role in the toxicokinetic and uptake of these antidepressants as it strongly influences their ionization state, with a general toxicity increase observed for higher pH values (Ford and Fong, 2016).

The most frequently detected SSRIs in surface waters are citalopram, fluoxetine, sertraline, and paroxetine, with reported levels up to 427, 330, 75 and 90 ng/L, respectively (Mole and Brooks, 2019; Silva et al., 2012; Sumpter and Margiotta-Casaluci, 2022). In this manuscript, the authors consider environmentally relevant those levels close to the levels reported in the environment.

Until the past decade, the predominant focus of studies examining the toxicity of SSRIs on organisms revolved around fluoxetine and sertraline. This emphasis was due to their significant consumption rates, physicochemical properties that affect their bioaccumulation and

metabolism in humans, as well as their capacity to influence behaviour (Brooks, 2014). The current environmental levels of SSRIs appear to reflect changes in consumption patterns as citalopram and paroxetine have also been increasingly and frequently detected in surface waters worldwide (Mole and Brooks, 2019; Silva et al., 2012; Sumpter and Margiotta-Casaluci, 2022) and, therefore, are also considered contaminants of emerging concern. Given that the class of antidepressants known as SSRIs encompasses substances with distinct pharmacokinetics and physicochemical properties, it is crucial to avoid generalizing scientific findings regarding their potential effects and toxicity. Variations in affinity for serotonin receptors have been observed among different SSRIs (Salahinejad et al., 2022; Esa-Pekka Palvimaki et al., 1999), and their interaction with fish detoxification systems may also differ. After uptake by fish, SSRIs, like other xenobiotics, are primarily subjected to a biotransformation process mediated by liver cytochrome P450 enzymes (CYP450) (Burkina et al., 2015). However, according to *in vitro* studies performed with fish cell fractions, the effects of SSRIs on the CYP450 system are distinct as they can act as inhibitors or inducers depending on the specific interaction between the SSRI itself and the enzymatic activity in question (Thibaut and Porte, 2008).

Therefore, we need to enhance our understanding of the pharmacokinetics and pharmacodynamics of SSRIs in fish. This knowledge will facilitate the improvement of environmental impact assessments related to these emerging contaminants.

3. Effects of SSRIs on fitness-related traits in fish

As SSRIs occur in the environment at very low or even at trace levels, they are more likely to cause sub-lethal chronic effects since they are specifically designed to elicit effects in their target organisms at low concentrations, and their biological targets are highly conserved across vertebrate taxa (Martin et al., 2021). SSRIs have already been identified in fish tissues, especially in the brain, highlighting their bioaccumulation potential (Moreira et al., 2022; Puckowski et al., 2016; Menon et al., 2020; Sumpter and Margiotta-Casaluci, 2022; Gould et al., 2021; Hughes et al., 2012). Arnok et al. (2017) (Arnok et al., 2017) reported high bioaccumulation of norsertaline (sertraline metabolite) in the liver and brain of fish. Grabicova et al. (2014) (Grabicova et al., 2014), observed that citalopram and sertraline were present in the brains and livers of most fish exposed to undiluted effluent from a sewage treatment plant, but not in blood, plasma or muscle. In human

patients, the brain-to-plasma ratio of fluoxetine is 2.6:1 (Karson et al., 1993). Long-term consumption of fish (including the entire body) may carry a potential risk to human health. Nonetheless, it should be highlighted that the only available study addressing human risk assessment related to SSRI exposure through food, concluded no evidence of considerable risk to human health associated with the consumption of marine mussels contaminated with SSRIs (Silva et al., 2017).

Current evidence highlights the ability of these environmental contaminants to cause maladaptive stress responses in fish, as they can also act and impact fish serotonergic system (Sehonova et al., 2018; Burkina et al., 2015; Sumpter and Margiotta-Casaluci, 2022; Sumpter et al., 2014) leading to impairment of diverse behavioural responses such as swimming, aggression, stress, fear responses and consequent fitness reduction (Sumpter and Margiotta-Casaluci, 2022; Salahinejad et al., 2022; Vinterstare et al., 2021). This is of particular concern since fishes play an ecological key role in aquatic ecosystems and, therefore, any SSRIs-induced behaviour changes can cause top-down effects on lower trophic levels affecting the whole ecosystem (Sehonova et al., 2018; Fernandez-Rubio et al., 2019).

Evidence suggests that due to the functional conservation of SSRI's therapeutical targets between humans and teleost fish, and SSRIs bioaccumulation potential, behavioural changes in fish are more likely to occur at plasma or tissue concentrations comparable to that inducing effects on humans (between 50 µg/L and 300 µg/L (Salahinejad et al., 2022; McDonald, 2017; Tanoue et al., 2019; Pan et al., 2018; Correia et al., 2023)). However, physiological, biochemical, and behavioural alterations have been reported on embryos/larvae of zebrafish exposed to fluoxetine levels as low as 0.1 µg/L, when the predicted plasma concentrations of fluoxetine were below human therapeutic doses (Salahinejad et al., 2022). Species with higher homology with humans, such as zebrafish, may be more sensitive to SSRIs (Salahinejad et al., 2022). Pan et al. (2018) (Pan et al., 2018) found evidence of tissue accumulation and concentration-dependent effects in two fish species (*D. rerio* and *Carassius auratus*) after short-term (6 days) exposure to the SSRI fluoxetine (0.1, 1, 10, 100 and 1000 µg/L). After long-term exposure, a more environmentally representative scenario, Pan et al. (2018) (Pan et al., 2018) detected the highest fluoxetine bioconcentration factor (BCF) in *C. auratus* at the lowest tested concentration (0.1 µg/L), suggesting that bioaccumulation levels may be higher in the natural environment where the concentrations are typically lower.

Among the SSRIs most detected in the environment, fluoxetine is considered to have the highest acute toxicity (Puckowski et al., 2016; Silva et al., 2012; Correia et al., 2023). This SSRI is the most investigated due to its environmental persistence and unique pharmacokinetics associated with a readily ionizable and toxic metabolite, norfluoxetine (Silva et al., 2012; Singh et al., 2022; Correia et al., 2023). Accordingly, extensive data concerning fluoxetine sub-lethal (acute and chronic) effects on fish are available (Menon et al., 2020; Fernandez-Rubio et al., 2019; Correia et al., 2023). Available data show that 7–28 days of exposure to environmentally relevant concentrations of fluoxetine (10–100 µg/L) induce anxiolytic responses in a variety of fish species (e.g., *D. rerio*, *Gasterosteus aculeatus*, *Oryzias latipes*, *Gambusia holbrooki*, *Poecilia reticulata*, *Pimephales promelas*) leading to maladaptive anxiety and fear responses which may impact ecologically relevant behaviours conditioning survival, such as feeding and predator escape (Pelli and Connaughton, 2015; Weinberger and Klaper, 2014; de Farias et al., 2020; Meijide et al., 2018; Ansai et al., 2016; Wong et al., 2013). Several studies associated acute and chronic exposures to fluoxetine (from 10 µg/L up to 10 mg/L) with altered (decreased) feeding behaviour in several fish species (e.g., *Cichlasoma dimerus*, *P. promelas*, *Pomatoschistus microps*) (Yang et al., 2021; Dorelle et al., 2020; Duarte et al., 2019). However, a lack of effects on feeding behaviour has also been reported. de Farias et al. (2020) found no effects on *D. rerio* feeding behaviour after 15 days of exposure to 100 µg/L fluoxetine (de Farias et al., 2020).

Changes in stress response, characterized by reduced anxiety and fear, are likely attributed to the effect of altered hormone levels, such as

cortisol and catecholamines. These effects may be related to interactions between the serotonergic and adrenergic systems and between the serotonergic system and the hypothalamus-pituitary-interrenal (HPI) axis (Sumpter and Margiotta-Casaluci, 2022; Kreke and Dietrich, 2008). Fluoxetine has been associated with effects on aggression and boldness across fish taxa (e.g., *D. rerio*, *Betta splendens*, *O. latipes*, *Neogobius melanostomus*), often with a negative correlation between the level of aggression and boldness and fluoxetine exposure concentrations (Eisenreich et al., 2017; Theodoridi et al., 2017; McCallum et al., 2017; Greene and Szalda-Petree, 2022; Dzieweczynski and Hebert, 2012). Nevertheless, the disruptive effect of fluoxetine on aggressive behaviour tends to be more consistent in exposures lasting between 5 and 21 days, with 75% of the studies demonstrating a reduction in aggression. In contrast, acute exposures (less than 5 days) show lower consistency, with 50% of the studies indicating decreased aggressiveness (McDonald, 2017).

An intraperitoneal administration of 25 and 50 µg fluoxetine/g of fish has been shown to affect processes such as branchial nitrogen excretion and intestinal osmoregulation in the teleost fish *Opsanus beta* (Morando et al., 2009). Richards et al. (2009) (Richards et al., 2009) demonstrated the ability of fluoxetine 100 and 1000 µg/L to constrict the arterio-arterial branchial vasculature, leading to impaired gas exchange and hypoxia, and ultimately to death, within the first 96 h of exposure to 1000 µg/L and after 35-day exposure to 100 µg/L fluoxetine. The ability of fluoxetine (50 ng/L) to modify circadian rhythm by reducing melatonin content in 5 days *D. rerio* larvae, has been reported with locomotor activity decrease in daytime associated with down-regulation of the clock genes *clock1a* and *bmal1b* (Wei et al., 2022). Nonetheless, fluoxetine ecotoxicity data focusing on ecologically relevant exposure conditions (e.g., concentrations and exposure duration) and ecologically relevant sublethal endpoints, such as altered behaviour, still lack consistency when it comes to correlating specific behaviours (e.g., predator avoidance, swimming, feeding) to risk assessment and management (Sumpter et al., 2014; Correia et al., 2023). Non-monotonic dose responses have been observed, with effects detected at low concentrations not being found at high concentrations, along with higher variations of investigated endpoints, mainly behavioural, at lower-level exposure and with species-specific and sex-dependent response (Pelli and Connaughton, 2015; Weinberger and Klaper, 2014; Yang et al., 2021; Salahinejad et al., 2022; Pan et al., 2018; Correia et al., 2023; de Farias et al., 2020; Meijide et al., 2018; Ansai et al., 2016; Kreke and Dietrich, 2008; Eisenreich et al., 2017; Theodoridi et al., 2017; McCallum et al., 2017; Greene and Szalda-Petree, 2022; Martin et al., 2017, 2019; Fursdon et al., 2019; Mennigen et al., 2010).

Compared to fluoxetine, the available data concerning the effects of citalopram, sertraline and paroxetine on fish are considerably limited. This fact is more evident for paroxetine. Short-term exposure to sertraline and citalopram seems to consistently result in a reduction of feeding behaviour in different fish species (Hedgspeth et al., 2014; Kellner et al., 2015; Hubena et al., 2021). A decrease in locomotor activity along with alterations in aggressiveness after 7–30 days of exposure to a wide range of citalopram and sertraline concentrations [0.38–100 µg/L] have also been reported (Yang et al., 2021; Kellner and Olsén, 2020; Hubena et al., 2021; Kania and Wronska, 2015) in different fish species. As for fluoxetine, these findings require further studies for their validation, as different sensitivity between species has been reported, along with a significant variation in exposure conditions. The available data for paroxetine suggest acute toxicity comparable to fluoxetine (e.g., paroxetine and fluoxetine 96 h median lethal concentration (LC₅₀) for the amphibian *Xenopus laevis* are 5.12 mg/L and 7.5 mg/L, respectively) (Silva et al., 2015). According to probabilistic environmental hazard assessments based on the calculation of therapeutic hazard values and fish plasma modelling approaches, paroxetine can be considered a contaminant of emerging concern and, thus, its ecotoxicity assessment is required to increase understanding of its potential environmental impacts (Mole and Brooks, 2019).

Overall, there are still considerable discrepancies in the available literature concerning behavioural effects on fish exposed to this class of antidepressants. Understanding the potential adverse impacts of SSRIs on fish behaviour is, undoubtedly, ecologically crucial as behavioural expression translates the interaction between an organism's internal status and its surrounding environment, and alterations in behaviour might impact fitness directly (Martin et al., 2019, 2021; Bertram et al., 2022; Ford et al., 2021). Indeed, behaviour can be considered a highly sensitive and useful endpoint in ecotoxicology assessment, being a multi-level indicator of biological effects and an early warning tool (Martin et al., 2021; Bertram et al., 2022; Ford et al., 2021; Orozco-Hernandez et al., 2022; Gaaied et al., 2020; Silva et al., 2020). However, behavioural assessment can itself be a source of variability as behaviour is not an easy endpoint to quantify, requiring a careful selection and execution of the tests considering the suitability and feasibility towards the behaviour it is intended to measure (Sumpter et al., 2014). This requires assessment criteria and standardized methods (e.g., US EPA, OECD) for fish toxicity studies with the inclusion of behavioural endpoints (Gould et al., 2021), as already done for other toxicological endpoints, to increase data consistency and thus improve reproducibility and reliability of the obtained results (Ford et al., 2021).

Future studies should focus on long-term and multigenerational exposures to increase understanding of sub-lethal SSRI-induced alterations under environmentally relevant scenarios. In humans, chronic SSRI treatment results in a profound reduction in serotonin transporter (SERT) expression and function, despite increased extracellular serotonin levels (Homborg et al., 2007). This complex pharmacokinetics of chronic SSRI exposure within humans suggests that to fully understand the implications of waterborne SSRI exposure on aquatic organisms, exposures should last at least for 21 days, as long-term compensatory mechanisms such as the desensitization of serotonin may occur (McDonald, 2017). Moreover, fish are more likely to be chronically exposed to complex mixtures of SSRIs which also requires further evaluation of the potential synergistic effects of combined SSRIs mixtures.

4. SSRIs and their impact on fish personality/coping styles

Different individuals cope with environmental challenges in different ways. This individual behavioural variance, like stress coping styles (SCS), has been associated with personality aspects (Tudorache et al., 2015; Fatsinia et al., 2020; Carbonara et al., 2019). Indeed, not all individuals within a species react the same way when subjected to the same conditions, even if they share the same age, size, sex, and place of origin (Grzesiuk and Pawelec, 2021). This variation in individual sensitivity may affect susceptibility to disease and/or environmental contaminants (Demin et al., 2019; Wong et al., 2019). However, the interaction between SCS, environmental stressors and disease susceptibility is still largely unknown.

Coping styles are suites of behavioural and physiological responses to stress that remain consistent over time and across contexts (Ferrari et al., 2020; Koolhaas et al., 2010a; Höglund et al., 2020). Such intra-specific behavioural variability is a well-documented phenomenon across vertebrates, including teleost fish and humans. Individuals of many species exhibit different SCS when subjected to stressful or risky situations, which may vary along a proactive-reactive continuum axis (Tudorache et al., 2015; Fatsinia et al., 2020; Carbonara et al., 2019; Thörnqvist et al., 2019). These behavioural patterns represent a relevant variable that may directly interfere with experimental outcomes in various behavioural tasks (Mackenzie et al., 2009). The existence of divergent SCS (proactive and reactive) among individuals and within a population is part of a fitness and survival strategy to optimize and ensure a correct adaptive response to changing environmental conditions (Ferrari et al., 2020). Different coping styles can already be present at a larval stage as shown by Tudorache et al. (2015) (Tudorache et al., 2015) in 8 days old zebrafish larvae.

Variations in coping styles are related and partially conditioned by heritable factors due to different gene expression patterns (Mackenzie et al., 2009; Fior et al., 2018a; Oswald et al., 2012), as well as consistent and divergent physiological and neuroendocrine responses to stress (Ferrari et al., 2020; Vinterstare et al., 2021; Höglund et al., 2020). Proactive fish are bolder, less plastic, more risk-taking, aggressive, territorial, and more predictable than shy fish. They are characterized by actively engaging stressors in a fight-flight strategy, exploring the environment quickly and superficially, with lower hypothalamic-pituitary-interrenal (HPI) axis reactivity and lower basal cortisol levels and higher sympathetic reactivity compared to reactive fish (Ferrari et al., 2020; Vinterstare et al., 2021; Höglund et al., 2020; Baker et al., 2018; Korte et al., 2005; Jolles et al., 2019). This behavioural pattern is better suited for predictable and stable environments. Conversely, reactive individuals are less aggressive and more cautious, exhibiting a conservation-withdrawal strategy that includes a freeze-hide response, with high HPI axis reactivity and baseline cortisol levels, and higher behavioural flexibility (Höglund et al., 2020; Korte et al., 2005), allowing them to cope with more unexpected and fluctuating environments (Carbonara et al., 2019; Baker et al., 2018). This different predisposition towards risk-prone behaviour in proactive-reactive individuals has proven effective in selecting fish, based on their personality, using standardized individual-based and/or group-based tests. Within this framework, various well-established protocols can be employed to differentiate proactive from reactive individuals and evaluate the consistency of coping styles over time. The group risk-taking test is one of the most successfully used tests when considering fish social species, that consists in measuring the time an individual takes to leave a group in the safety of a sheltered area to a new environment (area), thus exhibiting a risk-prone behaviour reflecting boldness level (Castanheira et al., 2013; Ferrari et al., 2015; Alfonso et al., 2019). The novel tank test is a valuable behavioural assay often utilized to distinguish between proactive and reactive individuals. This test shares a similar protocol with the previously described group risk-taking test, but with the notable difference that it is done with one individual at a time (Wong et al., 2012). The feeding latency test and the novel object test are other examples of individual-based assays that allow measuring fish boldness (Wong et al., 2012). Most of the performed studies in this scope have used different combinations of both individual and group-based tests to assess the repeatability and consistency of individual behavioural traits over short periods (from 1 to 14 days) (Castanheira et al., 2013; Ferrari et al., 2015; Alfonso et al., 2019; Wilson and Stevens, 2005; Øverli et al., 2007; Wilson and Godin, 2009). Indeed, the reproducibility of the outcomes seems to be higher for measurements that are repeated in short periods as individuals are more consistent in behaviour over short periods in comparison to long ones (Ferrari et al., 2015; Bell et al., 2009). Several factors, encompassing predation pressure, food availability, social interactions, temperature, hypoxia, and learning, have the potential to shape coping styles (Ferrari et al., 2015). Castanheira et al. (2016) (Castanheira et al., 2016) observed SCS behaviour consistency in *Sparus aurata* only before sexual maturation and highlighted the importance of considering life history. Thus, the behavioural aspect of coping styles is not stationary and may change according to life history events (Castanheira et al., 2016), besides appearing to be species-specific (Carbonara et al., 2019).

Current evidence demonstrates the association between behavioural, physiological, and neurological differences in proactive and reactive coping styles in fish and different expressions of central serotonin function (Koolhaas et al., 2010a; Coppens et al., 2010; Tran et al., 2016; Backström and Winberg, 2017). Additionally, brain serotonin is causally implicated in both aggression and behavioural flexibility, as lower prefrontal cortex serotonin levels have been linked to heightened aggression and impulsive behaviour (Coppens et al., 2010). Aggressive and impulsive control disorders in humans have been strongly correlated with low levels of brain serotonin and its metabolite 5-hydroxyindoleacetic acid (5-HIAA) along with up-regulation of the auto-receptors

(5-HT1A and 5-HT1B) located in presynaptic neurons that act as an additional self-regulatory mechanism of aggressive behaviour with inhibitory effects on the serotonin neurotransmission (Pavlov et al., 2012; Koolhaas et al., 2010b). Although it is well established that SSRIs can interfere with fish behaviour, data on differential responses to this type of antidepressant depending on fish personality, is very limited (Salahinejad et al., 2022). In humans, the suppression of certain personality traits due to SSRI treatment has been reported to occur in some cases (Grzesiuk and Pawelec, 2021).

As aggression and boldness are fish personality traits correlated with serotonin levels, it is expected that drugs that interfere with the expression of the serotonergic system, such as SSRIs, may lead to behavioural phenotype modulation (Fior et al., 2018a). Indeed, the modulation of bold (proactive) behaviour by the SSRI fluoxetine (0.5 µg–50 µg/L) into a shy phenotype has been recently reported in some fish species (*B. splendens*, *D. rerio*) (Fior et al., 2018b; Dziewieczynski et al., 2016a, 2016b). Nonetheless, Grzesiuk and Pawelec (2021) (Grzesiuk and Pawelec, 2021), in a study performed with two freshwater wild fish species (*N. fluviatilis* and *Gobio*), observed an increase in the proportion of bold individuals after 21 days of exposure to an environmentally relevant concentration of fluoxetine (360 ng/L). In a two-year multigenerational assay with *P. reticulata*, Polverino et al. (2021) (Polverino et al., 2021) reported that exposure to environmentally realistic levels of fluoxetine (40 and 366 ng/L) suppresses behavioural differences between individuals by diluting the individual activity level and risk-taking behaviour. Tan et al. (2020) (Tan et al., 2020) also observed a reduction in among-individual phenotypic variation in *P. reticulata* females after a long-term (15 months) exposure to environmentally relevant levels of fluoxetine (30 and 300 ng/L). These outcomes can be considered a first line of evidence that should guide further studies for a comprehensive insight into the ecological adverse implications that may arise from SSRIs-suppressive effects on individual coping strategies, by reducing intraspecific behavioural variance, which may ultimately compromise the ability of the whole population to adapt in an ever-changing environment. The potential implications are considerable if individuals experience increased shyness due to SSRI exposure. Shy individuals are typically less prone to explore and take risks, which could hinder their ability to venture into new environments with improved conditions and resources. Furthermore, increased susceptibility to disease may also be a consequence of their cautious nature. Data concerning other SSRIs of environmental concern such as citalopram, sertraline and paroxetine are even more scarce or non-existent. Kellner et al. (2020) (Kellner and Olsén, 2020) observed a suppressive effect of citalopram on *G. aculeatus* aggressiveness, with no effects on personality-dependent behaviours (no significant correlation between feeding behaviour and aggressive behaviour) found in response to citalopram exposure. To the best of our knowledge, there are no studies performed so far concerning the potential adverse effects of sertraline or paroxetine on fish personality.

Given the documented sex-dependent responses to SSRIs concerning behavioural traits such as feeding and aggression (Kellner and Olsén, 2020), it is plausible to hypothesize that a similar response pattern might extend to specific behaviours associated with personality traits. In this context, the involvement of estradiol in shaping female behavioural expression has been proposed (Dziewieczynski et al., 2016a). Studies by Mennigen et al. (2010) (Mennigen et al., 2010) and Lister et al. (2009) (Lister et al., 2009) revealed decreased estradiol levels in female goldfish (*C. auratus*) and zebrafish following fluoxetine exposure, which could potentially impact personality-dependent sexual behaviours and, subsequently, reproductive success. Clearly, there is still a long way ahead as we are only starting to understand to what extent SSRI's effects on personality-dependent traits may impair fitness-related behaviours leading to progressive physiological dysregulation that can ultimately result in disease and mortality (Christensen et al., 2019). Fish are more likely to be chronically exposed to contaminants throughout their life cycle. Thus, given that SSRI's therapeutical effects on people are

observed following 3–4 weeks of treatment, future studies should focus on exposures of not less than 21–28 days, particularly to concentrations often detected on surface waters.

A growing line of evidence highlights the importance of personality as a significant factor to be considered in farmed fish when to improve the management and welfare of aquaculture populations (Castanheira et al., 2017). Therefore, the potential economic impact of SSRI's effects on personality-dependent traits should not be neglected as farmed fish may be exposed to these environmental contaminants as they are commonly reared in natural water resources in several aquaculture farms. This fact may affect vulnerability and susceptibility to disease, making it difficult to optimize mitigation measures to minimise stressors that may act as triggers for disease.

Furthermore, there is also the need to increase knowledge concerning the potential transgenerational implications of SSRIs-suppressive effects on phenotypic variation in fish. As bold and aggressive personality traits are heritable with a significant maternal inheritance proportion, the potential interference of SSRIs on this heritable character should also be studied (Ariyomo et al., 2013).

5. Translational research: fish as a proxy for individual disease vulnerability

Current evidence highlights a relationship between personality traits and health (Zozulya et al., 2008; Murray and Booth, 2015). Under stressful conditions, both human personality traits and specific behavioural patterns in animals are prominently exhibited (Mehta and Gosling, 2008). Several studies have demonstrated that individuals from many species, including teleost fish and humans, deal with stress differently by adopting different behavioural strategies to overcome stress (Wong et al., 2019). This variation in individual sensitivity may affect disease susceptibility (McDonald, 2017; Demin et al., 2019). Despite the myriad differences in morphology, physiology, behaviour and biological complexity within humans and non-human animals, translational research has proven to be very helpful in understanding personality, immunity, and health. Humans share with other species many molecular and biochemical pathways involved in behavioural expression that are highly conserved across animal taxa (Ford et al., 2021; Wilson and Godin, 2009; Castanheira et al., 2016). The study of various behavioural models has shown that coping style determines, to a large degree, the reaction of the neuroendocrine system to stress, therefore influencing several aspects of the stress response such as activation of the sympathetic nervous system (SNS) and hypothalamic-pituitary-interrenal/adrenal (HPI/HPA) axis, both systems affecting the immune system (Wilson et al., 2019). In this context, zebrafish has emerged as a promising model for the study of behaviour and personality due to high genetic and physiological homology with humans along with two well-defined personality patterns (proactive/bold and reactive/shy) and the same modulatory neurotransmitters found in the mammalian brain (e.g., serotonergic, and dopaminergic systems) (Adhish and Manjubala, 2023; Xie et al., 2022; Gould et al., 2021). Therefore, assessing the effects of SSRIs on zebrafish personality may be very helpful in understanding the potential effects of these antidepressants on human personality, which is clinically and environmentally relevant. The silencing of personality traits is sometimes a consequence of SSRI's therapeutical treatment (Homborg et al., 2007). However, its potential influence on human response to environmental stressors (e.g., contaminants) and disease susceptibility is unknown. Additionally, effects related to unintentional exposure to these antidepressants should also be addressed as SSRIs have already been detected in some countries in drinking water (McDonald, 2017).

5. Final considerations

As current data highlight the potential of SSRIs to modulate personality by erasing intraspecific behavioural variance within a

population, it is environmentally relevant to understand how this can impact an organism's ability to cope with additional environmental stressors (e.g., temperature, chemicals, micro (nano)plastics and their combination) in such challenging and ever-changing environment. Considering the widespread consumption of antidepressants and the increasing exposure of humans to a diversity of environmental stressors, this approach could prove beneficial and potentially applicable to humans as well. The presence of non-monotonic responses in studies with SSRIs underscores the importance of prioritizing the development of environmentally relevant tests. These tests are essential for achieving a comprehensive understanding of the environmental consequences associated with SSRIs. Given that behaviour is influenced by integrated effects at multiple biological levels, it is crucial to include long-term and multigenerational assessments, incorporating various types of SSRIs, in behavioural assays due to their environmental relevance. By adopting this approach, we can accurately evaluate the long-term impacts of SSRIs on the environment.

Credit author statement

Carla Ferreira: Conceptualization, Investigation, Writing – Original Draft, Writing – Review & Editing; Sandra C. Soares: Review & Editing, Supervision; Peter Kille: Review & Editing, Supervision; Miguel Oliveira: Conceptualization, Writing – Review and Editing, Visualization, Supervision and Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgements

Thanks are due to FCT/MCTES for the financial support to the project NanoPlanet (2022.02340.PTDC) and CESAM (UIDP/50017/2020+UIDB/50017/2020 + LA/P/0094/2020) through national funds. Carla Ferreira was supported by FCT through a PhD Grant (2021.04580. BD).

References

- Adhish, M., Manjubala, I., 2023. Effectiveness of zebrafish models in understanding human diseases—a review of models. *Heliyon* 9 (3), e14557.
- Alfonso, S., Sadoul, B., Gesto, M., Joassard, L., Chatain, B., Geffroya, B., Bégout, M.L., 2019. Coping styles in European sea bass: the link between boldness, stress response and neurogenesis. *Physiol. Behav.* 207, 76–85.
- Ansai, S., Hosokawa, H., Maegawa, S., Kinoshita, M., 2016. Chronic fluoxetine treatment induces anxiolytic responses and altered social behaviours in medaka. *Oryzias latipes*. *Behav. Brain Res.* 303, 126–136.
- Ariyomo, T.O., Carter, M., Watt, P.J., 2013. Heritability of boldness and aggressiveness in the zebrafish. *Behav. Genet.* 43, 161–167.
- Arnnok, P., Singh, R.R., Burakham, R., Pérez-Fuentetaja, A., Aga, D.S., 2017. Selective uptake and bioaccumulation of antidepressants in fish from effluent-impacted niagara river. *Environ. Sci. Technol.* 51 (18), 10652–10662.
- Bachour, R.L., Golovko, O., Kellner, M., Pohl, J., 2020. Behavioural effects of citalopram, tramadol, and binary mixture in zebrafish (*Danio rerio*) larvae. *Chemosphere* 238, 124587.
- Backström, T., Winberg, S., 2017. Serotonin coordinates responses to social stress - what we can learn from fish. *Front. Neurol.* 11 (595), 1–10.
- Baker, M.R., Goodman, A.C., Santo, J.B., Wong, R.Y., 2018. Repeatability and reliability of exploratory behaviour in proactive and reactive zebrafish, *Danio rerio*. *Sci. Rep.* 8, 12114.
- Bell, A.M., Hankison, S.J., Laskowski, K.L., 2009. The repeatability of behaviour: a meta-analysis. *Anim. Behav.* 77, 771–783.
- Bergmüller, R., 2020. Animal personality and behavioural syndromes. *Anim. Behav.: Evolution and Mechanisms* 19, 587–621.

- Bertram, M.G., Martin, J.M., McCallum, E.S., Alton, L.A., Brand, J.A., Brooks, B.W., Cerveny, D., Fick, J., Ford, A.T., Hellström, G., Michelangeli, M., Nakagawa, S., Polverino, G., Saarisalo, M., Sih, A., Tan, H., Tyler, C.R., Wong, B.B.M., Brodin, T., 2022. Frontiers in quantifying wildlife behavioural responses to chemical pollution. *Biol. Rev.* 97, 1346–1364.
- Bisei, J.H., Sweet, L.E., van den Hurk, P., Klaine, S.J., 2016. Effects of an antidepressant mixture on the brain serotonin and predation behaviour of hybrid striped bass. *Environ. Toxicol. Chem.* 35, 938–945.
- Branchet, P., Arpin-Pont, L., Piram, A., Wong-Wah-Chung, P., Doumenq, P., Review, P., 2022. Pharmaceuticals in the marine environment: what are the present challenges in their monitoring? *Sci. Total Environ.* 766, 142644.
- Brooks, B.W., 2014. Fish on Prozac (and Zoloft): ten years later. *Aquat. Toxicol.* 151, 61–67.
- Burkina, V., Zlabeka, V., Zamaratskaia, G., 2015. Effects of pharmaceuticals present in aquatic environment on Phase I metabolism in fish (Review). *Environ. Toxicol. Pharmacol.* 40, 430–444.
- Carbonara, P., Alfonso, S., Zupa, W., Manfrin, A., Flocchi, E., Pretto, T., Spedicato, M.T., Lembo, G., 2019. Behavioural and physiological responses to stocking density in sea bream (*Sparus aurata*): do coping styles matter? *Physiol. Behav.* 212, 112698.
- Castanheira, M.F., Herrera, M., Costas, B., Conceição, L.E.C., Martins, C.I.M., 2013. Can we predict personality in fish? Searching for consistency over time and across contexts. *PLoS One* 8, e62037.
- Castanheira, M.F., Martínez-Páramo, S., Figueiredo, F., Cerqueira, M., Millot, S., Oliveira, C.C.V., Martins, C.I.M., Conceição, L.E.C., 2016. Are coping styles consistent in the teleost fish *Sparus aurata* through sexual maturation and sex reversal? *Fish Physiol. Biochem.* 42, 1441–1452.
- Castanheira, M.F., Conceição, L.E.C., Millot, S., Rey, S., Begout, M.L., Damsgard, B., Kristiansen, T., Høglund, E., Øverli, Ø., Martins, C.I.M., 2017. Coping styles in farmed fish: consequences for aquaculture. *Rev. Aquacult.* 9, 23–41.
- Christensen, D.S., Nadya Dich, N., Flensburg-Madsen, T., Garde, E., Hansen, A.M., Mortensen, E.L., 2019. Objective and subjective stress, personality, and allostatic load. *Brain and Behaviour* 9, 1–11.
- Coppens, C.M., Boer, S.F., Koolhaas, J.M., 2010. Coping styles and behavioural flexibility: towards underlying mechanisms. *Phil. Trans. R. Soc. B.* 365, 4021–4028.
- Correia, D., Domingues, I., Faria, M., Oliveira, M., 2023. Effects of fluoxetine on fish: what do we know and where should we focus our efforts in the future? *Sci. Total Environ.* 857 (Pt 2), 159486.
- de Farias, N.O., Oliveira, R., Moretti, P.N.S., e Pinto, J.M., Oliveira, A.C., Santos, V.L., Rocha, P.S., Andrade, T.S., Grisolia, C.K., 2020. Fluoxetine chronic exposure affects growth, behaviour and tissue structure of zebrafish. *Comp. Biochem. Physiol., Part C: Toxicol. Pharmacol.* 237, 108836.
- Demin, K.A., Lakstygala, A.M., Alekseeva, P.A., Sysoev, M., de, S., Abreue, M., Alpyshov, E.T., Serikulyc, N., Wang, D., Wang, M., Tang, Z., Yanc, D., Strekalovak, T.V., Volgind, A.D., Amstislavskaya, T.G., Wang, J., Song, C., Kalueff, A.V., 2019. The role of intraspecific variation in fish neurobehavioural and neuropharmacological phenotypes in aquatic models. *Aquat. Toxicol.* 210, 44–55.
- Desbiolles, F., Malleret, L., Tiliacos, C., Wong-Wah-Chung, P., Laffont-Schwob, I., 2018. Occurrence and ecotoxicological assessment of pharmaceuticals: is there a risk for the Mediterranean aquatic environment? *Sci. Total Environ.* 639, 1334–1348.
- Diaz-Camal, N., Cardoso-Vera, J.D., Islas-Flores, H., Gómez-Oliván, L.M., Mejía-García, A., 2022. Consumption and occurrence of antidepressants (SSRIs) in pre- and post- COVID-19 pandemic, their environmental impact and innovative removal methods: a review. *Sci. Total Environ.* 829, 154656.
- Dorelle, L.S., Da Cunha, R.H., Sganga, D.E., Rey Vázquez, G., López Greco, L., Lo Nostro, F. L., 2020. Fluoxetine exposure disrupts food intake and energy storage in the cichlid fish *Cichlasoma dimerus* (Teleostei, Cichliformes). *Chemosphere* 238, 124609, 2020.
- Duarte, I.A., Pais, M.P., Reis-Santos, P., Cabral, H.N., Fonseca, V.F., 2019. Biomarker and behavioural responses of an estuarine fish following acute exposure to fluoxetine. *Mar. Environ. Res.* 147, 24–31, 2019.
- Dziewieczynski, T.L., Hebert, O.L., 2012. Fluoxetine alters behavioural consistency of aggression and courtship in male Siamese fighting fish. *Betta splendens*. *Physiol. Behav.* 107 (1), 92–97.
- Dziewieczynski, T.L., Kane, J.L., Campbell, B.A., Lavin, L.E., 2016a. Fluoxetine exposure impacts boldness in female Siamese fighting fish. *Betta splendens*. *Ecotoxicology*. 25, 69–79.
- Dziewieczynski, T.L., Campbell, B.A., Kane, J.L., 2016b. Dose-dependent fluoxetine effects on boldness in male Siamese fighting fish. *J. Exp. Biol.* 219, 797–804.
- Eisenreich, B.R., Greene, S., Szalda-Petree, A., 2017. Of fish and mirrors: fluoxetine disrupts aggression and learning for social rewards. *Physiol. Behav.* 173, 258–262.
- Esa-Pekka Palvimäki, E.P., Kuoppamäki, M., Syvalähti, E., Hietala, J., 1999. Differential effects of fluoxetine and citalopram treatments on serotonin 5-HT_{2C} receptor occupancy in rat brain. *Int. J. Neuropsychopharmacol.* 2, 95–99.
- Estrela, M., Herdeiro, M.T., Ferreira, P.L., Roque, F., 2020. The use of antidepressants, anxiolytics, sedatives and hypnotics in Europe: focusing on mental health care in Portugal and prescribing in older patients. *Int. J. Environ. Res. Publ. Health* 17, 8612.
- Fatsinia, E., Reyc, S., Ibarra-Zatarain, Z., Boltana, S., Mackenzie, S., Duncan, N.J., 2020. Linking stress coping styles with brain mRNA abundance of selected transcripts for Senegalese sole (*Solea senegalensis*) juveniles. *Physiol. Behav.* 213, 112724.
- Fekadu, S., Alemayehu, E., Dewil, R., Bruggen, B.V., 2019. Review - pharmaceuticals in freshwater aquatic environments: a comparison of the African and European challenge. *Sci. Total Environ.* 654, 324–337.
- Fernandez-Rubio, J., Rodríguez-Gil, J., Postigo, C., Mastroianni, N., Alda, M., Barcel, D., Valcarlos, Y., 2019. Psychoactive pharmaceuticals and illicit drugs in coastal waters

- of North-Western Spain: environmental exposure and risk assessment. *Chemosphere* 224, 379–389.
- Ferrari, S., Millot, S., Leguaya, D., Chatain, B., Bégout, M.L., 2015. Consistency in European seabass coping styles: a life-history approach. *Appl. Anim. Behav. Sci.* 167, 74–88.
- Ferrari, S., Rey, S., Høglund, E., Øverli, Ø., Chatain, B., MacKenzie, S., Bégout, M.L., 2020. Physiological responses during acute stress recovery depend on stress coping style in European sea bass. *Dicentrarchus labrax*. *Physiology & Behaviour* 216, 112801.
- Fior, D., Dametto, F., Fagundes, M., Santos da Rosa, J.G., Sander de Abreu, M., Koalkoski, G., Idalencio, R., de Alcântara Barcellos, H.H., Piato, A., Gil Barcellos, L.J., 2018a. Divergent action of fluoxetine in zebrafish according to responsivity to novelty. *Sci. Rep.* 8, 13908.
- Fior, D., Dametto, F., Fagundes, M., et al., 2018b. Divergent action of fluoxetine in zebrafish according to responsivity to novelty. *Sci. Rep.* 8, 13908.
- Ford, A.T., Fong, P.P., 2016. The effects of antidepressants appear to be rapid and at environmentally relevant concentrations. *Environ. Toxicol. Chem.* 35 (4), 794–798.
- Ford, A.T., Ågerstrand, M., Brooks, B.W., Allen, J., Bertram, M.G., Brodin, T., Dang, Z., Dugesne, S., Sahm, R., Hoffmann, F., Hollert, H., Jacob, S., Klüver, N., Lazorchak, J., Ledesma, M., Melvin, S.D., Mohr, S., Padilla, S., Pyle, G., Scholz, S., Saaristo, M., Smit, E., Steevens, J.A., van den Berg, S., Kloas, W., Wong, B.B.M., Ziegler, M., Maack, G., 2021. The role of behavioral ecotoxicology in environmental protection. *Environ. Sci. Technol.* 55, 5620–5628.
- Fursdon, J.B., Martin, J.M., Bertram, M.G., Lehtonen, T.K., Wong, B.B.M., 2019. The pharmaceutical pollutant fluoxetine alters reproductive behaviour in a fish independent of predation risk. *Sci. Total Environ.* 650, 642–652.
- Gaaied, S., Oliveira, M., Domingues, I., Banni, M., 2020. 2, 4-Dichlorophenoxyacetic acid herbicide effects on zebrafish larvae: development, neurotransmission and behavior as sensitive endpoints. *Environ. Sci. Pollut. Control Ser.* 27 (4), 3686–3696.
- Gould, S.L., Winter, M.J., Norton, W.H., Tyler, C.R., 2021. The potential for adverse effects in fish exposed to antidepressants in the aquatic environment. *Environ. Sci. Technol.* 55 (24), 16299–16312.
- Grabicova, K., Lindberg, R.H., Östman, M., Grabic, R., Randak, T., Larsson, D.G.J., Fick, J., 2014. Tissue-specific bioconcentration of antidepressants in fish exposed to effluent from a municipal sewage treatment plant. *Sci. Total Environ.* 488–489, 46–50.
- Grabicova, K., Grabic, R., Fedorova, G., Fick, J., Cerveny, D., Kolarova, J., Turek, J., Zlabek, V., Randak, T., 2017. Bioaccumulation of psychoactive pharmaceuticals in fish in an effluent dominated stream. *Water Res.* 124, 654–662.
- Greene, S.M., Szalda-Petree, A.D., 2022. Fins of Fury or Painéant: fluoxetine impacts the aggressive behaviour of fighting fish (*Betta splendens*). *Behav. Process.* 194, 104544.
- Grzesiuk, M., Pawelec, A., 2021. Fluoxetine results in misleading conclusions on fish behaviour. *Ecol. Evol.* 11, 9707–9714.
- Hedgesepe, M.L., Nilsson, P.A., Berglund, O., 2014. Ecological implications of altered fish foraging after exposure to an antidepressant pharmaceutical. *Aquat. Toxicol.* 151, 84–87.
- Høglund, E., Moltesen, M., Castanheira, M.F., Thörnqvist, P.O., Silva, P.I.M., Øverli, Ø., Martins, C., Winberg, S., 2020. Contrasting neurochemical and behavioural profiles reflects stress coping styles but not stress responsiveness in farmed gilthead seabream (*Sparus aurata*). *Physiol. Behav.* 214, 112759.
- Homborg, J.R., Olivier, J.D.A., Smits, B.M.G., Mul, J.D., Mudde, J., Verheul, M., Nieuwenhuizen, O.F.M., Cools, A.R., Ronken, E., Cremers, T., Schoffemeer, A.N.M., Ellenbroek, B.A., Cuppen, E., 2007. Characterization of the serotonin transporter knockout rat: a selective change in the functioning of the serotonergic system. *Neuroscience* 146, 1662–1676.
- Hossain, M.S., Kubeck, J., Guo, W., Roje, S., Ložek, F., Grabicová, K., Randák, T., Koubal, A., Burič, M., 2021. A combination of six psychoactive pharmaceuticals at environmental concentrations alter the locomotory behaviour of clonal marbled crayfish. *Sci. Total Environ.* 751, 141383.
- Huang, J.J., Sirotkin, H.I., McElroy, A.E., 2019. Varying the exposure period and duration of neuroactive pharmaceuticals and their metabolites modulates effects on the visual motor response in zebrafish (*Danio rerio*) larvae. *Neurotoxicol. Teratol.* 72, 39–48.
- Huang, J.J., Dheilly, N.M., Sirotkin, H.I., McElroy, A.E., 2020. Comparative transcriptomics implicate mitochondrial and neurodevelopmental impairments in larval zebrafish (*Danio rerio*) exposed to two selective serotonin reuptake inhibitors (SSRIs). *Ecotoxicol. Environ. Saf.* 203, 110934.
- Hubena, P., Horký, P., Grabic, R., Grabicova, K., Douda, K., Slavík, O., Randak, T., 2021. Prescribed aggression of fishes: pharmaceuticals modify aggression in environmentally relevant concentrations. *Ecotoxicol. Environ. Saf.* 127, 112944.
- Hughes, S.R., Kay, P., Brown, L.E., 2012. Global synthesis and critical evaluation of pharmaceutical data sets collected from river systems. *Environ. Sci. Technol.* 47, 661–677.
- Jolles, J.W., Briggs, H.D., Araya-Ajoy, Y.G., Boogert, N.J., 2019. Personality, plasticity and predictability in sticklebacks: bold fish are less plastic and more predictable than shy fish. *Anim. Behav.* 154, 193–202.
- Kania, B.F., Wrońska, D., 2015. The selective serotonin reuptake inhibitor-sertraline diminishes conspecific aggression in male fighting *Betta splendens* fish. *J. Behav. Brain Sci.* 5, 578–585.
- Karson, C.N., Newton, J.E., Livingston, R., Jolly, J.B., Cooper, T.B., Sprigg, J., Komoroski, R.A., 1993. Summer. Human brain fluoxetine concentrations. *J. Neuropsychiatry Clin. Neurosci.* 5 (3), 322–329.
- Kaushik, G., Thomas, M.A., 2019. The potential association of psychoactive pharmaceuticals in the environment with human neurological disorders. *Sustain. Chem. Pharm.* 13, 100148.
- Kellner, M., Olsén, K.H., 2020. Divergent response to the SSRI citalopram in male and female three-spine sticklebacks (*Gasterosteus aculeatus*). *Arch. Environ. Contam. Toxicol.* 79, 478–487.
- Kellner, M., Porseryd, T., Porsch-Hallstrom, I., Hansen, S.H., Olsen, K.H., 2015. Environmentally relevant concentrations of citalopram partially inhibit feeding in the three-spine stickleback (*Gasterosteus aculeatus*). *Aquat. Toxicol.* 158, 165–170.
- Klatte, S., Schaefer, H.C., Hempel, M., 2017. Pharmaceuticals in the environment – a short review on options to minimize the exposure of humans, animals and ecosystems. *Sustain. Chem. Pharm.* 5, 61–66.
- Koolhaas, J.M., 2008. Coping style and immunity in animals: making sense of individual variation. *Brain Behav. Immun.* 22, 662–667.
- Koolhaas, J.M., de Boer, S.F., Coppens, C.M., Buwalda, B., 2010a. (Review) Neuroendocrinology of coping styles: towards understanding the biology of individual variation. *Front. Neuroendocrinol.* 31, 307–321.
- Koolhaas, J.M., de Boer, S.F., Coppens, C.M., Buwalda, B., 2010b. Neuroendocrinology of coping styles: towards understanding the biology of individual variation. *Front. Neuroendocrinol.* 31 (3), 307–321.
- Korte, S.M., Koolhaas, J.M., Wingfield, J.C., McEwen, B.S., 2005. The Darwinian concept of stress: benefits of allostasis and cost of allostatic load and the trade-offs in health and disease. *Neuroscience and Biobehavioural Rev.* 29, 3–38.
- Kreke, N., Dietrich, D.R., 2008. Physiological endpoints for potential SSRI interactions in fish. *Crit. Rev. Toxicol.* 37 (3), 215–247.
- Lister, A., Regan, C., Van Zwol, J., Van Der Kraak, G., 2009. Inhibition of egg production in zebrafish by fluoxetine and municipal effluents: a mechanistic evaluation. *Aquat. Toxicol.* 95, 320–329.
- Mackenzie, S., Ribas, L., Pilarczyk, M., Capdevila, D.M., Kadri, S., Huntingford, F.A., 2009. Screening for coping style increases the power of gene expression studies. *PLoS One* 4, 1–5.
- Madikizela, L.M., Ncube, S., Tutuc, H., Richards, H., Newman, B., Ndungue, K., Chimukac, L., 2020. Pharmaceuticals and their metabolites in the marine environment: sources, analytical methods and occurrence. *Trends Environ. Anal. Chem.* 28, e00104.
- Martin, J.M., Saaristo, M., Bertram, M.G., Lewis, P.J., Coggan, T.L., Clarke, B.O., Wong, B.B.M., 2017. The psychoactive pollutant fluoxetine compromises antipredator behaviour in fish. *Environ. Pollut.* 222, 592–599.
- Martin, J.M., Bertram, M.G., Saaristo, M., Ecker, T.E., Hannington, S.L., Tanner, J.L., Michelangeli, M., O'Bryan, M.K., Wong, B.B.M., 2019. Impact of the widespread pharmaceutical pollutant fluoxetine on behaviour and sperm traits in a freshwater fish. *Sci. Total Environ.* 650, 1771–1778.
- Martin, J.M., Bertram, M.G., Blanchfield, P., Brand, J.A., Brodin, T., Brooks, B.W., Cerveny, D., Lagisz, M., Ligocki, I.Y., Michelangeli, M., Nakagawa, S., Orford, J.T., Sundin, J., Tan, H., Wong, B.B.M., McCallum, E.S., 2021. Evidence of the impacts of pharmaceuticals on aquatic animal behaviour: a systematic map protocol. *Environ. Evid.* 10, 26.
- McCallum, E.S., Bose, A.P., Warriner, T.R., Balshine, S., 2017. An evaluation of behavioural endpoints: the pharmaceutical pollutant fluoxetine decreases aggression across multiple contexts in round goby (*Neogobius melanostomus*). *Chemosphere* 175, 401–410.
- McDonald, M.D., 2017. An AOP analysis of selective serotonin reuptake inhibitors (SSRIs) for fish. *Review. Comp. Biochem. Physiol.* 197, 19–31. Part C.
- Mehta, P.H., Gosling, S.D., 2008. Bridging human and animal research: a comparative approach to studies of personality and health. *Brain Behav. Immun.* 22, 651–661.
- Meijide, F.J., Da Cuña, R.H., Prieto, J.P., Dorelle, L.S., Babay, P.A., Nostro, F.L.L., 2018. Effects of waterborne exposure to the antidepressant fluoxetine on swimming, shoaling and anxiety behaviours of the mosquitofish *Gambusia holbrooki*. *Ecotoxicol. Environ. Saf.* 163, 646–655.
- Mennigen, J.A., Lado, W.E., Zamora, J.M., Duarte-Guterman, P., Langlois, V.S., Metcalfe, C.D., Chang, J.P., Moon, T.W., Trudeau, V.L., 2010. Waterborne fluoxetine disrupts the reproductive axis in sexually mature male goldfish, *Carassius auratus*. *Aquat. Toxicol.* 100, 354–364.
- Menon, N.G., Mohapatra, S., Padhye, L.P., Tatiparti, S.S.V., Mukherji, S., 2020. Review on occurrence and toxicity of pharmaceutical contamination in Southeast Asia. In: *Emerging Issues in the Water Environment during Anthropocene*. Springer, Singapore, pp. 63–91.
- Mezzelani, M., Gorbi, S., Regoli, F., 2018. Pharmaceuticals in the aquatic environments: evidence of emerged threat and future challenges for marine organisms. *Mar. Environ. Res.* 140, 41–60.
- Mole, R.A., Brooks, B.W., 2019. Global scanning of selective serotonin reuptake inhibitors: occurrence, wastewater treatment and hazards in aquatic systems. *Environ. Pollut.* 250, 1019–1031.
- Morando, M.B., Medeiros, L.R., McDonald, M.D., 2009. Fluoxetine treatment affects nitrogen waste excretion and osmoregulation in a marine teleost fish. *Aquat. Toxicol.* 95, 164–171.
- Moreira, D.G., Aires, A., de Lourdes Pereira, M., Oliveira, M., 2022. Levels and effects of antidepressant drugs to aquatic organisms. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 256, 109322.
- Murray, A.L., Booth, T., 2015. Personality and physical health. *Curr. Opin. Psychol.* 5, 50–55.
- Nguyena, P.Y., Carvalho, G., Reis, M.A.M., Oehmenc, A., 2021. A review of the biotransformations of priority pharmaceuticals in biological wastewater treatment processes. *Water Res.* 188, 116446.
- Organization for Economic Co-operation and Development (OECD) Available at: [Tackling the Mental Health Impact of the COVID-19 Crisis: an Integrated, Whole-Of-Society Response \(oecd.org\)](https://www.oecd.org/).
- Orozco-Hernandez, J.M., Gomez-Oliván, L.M., Elizalde-Velazquez, G.A., Rosales-Pérez, K.E., Cardoso-Vera, J.D., Heredia-García, G., Islas-Flores, H., García-

- Medina, S., Galar-Martínez, M., 2022. Fluoxetine-induced neurotoxicity at environmentally relevant concentrations in adult zebrafish *Danio rerio*. *Neurotoxicology* 90, 121–129.
- Oswald, M.E., Drew, R.E., Racine, M., Murdoch, G.K., Robison, B.D., 2012. Is behavioural variation along the bold-shy continuum associated with variation in the stress axis in zebrafish? *Physiol. Biochem. Zool.* 85, 718–728.
- Øverli, Ø., Sorensen, C., Pulman, K.G.T., Pottinger, T.G., Korzan, W., Summers, C.H., Nilsson, G.E., 2007. Evolutionary background for stress-coping styles: relationships between physiological, behavioural, and cognitive traits in non-mammalian vertebrates. *Neurosci. Biobehav. Rev.* 31, 396–412.
- Pan, C., Yang, M., Xu, H., Xu, B., Jiang, L., Wu, M., 2018. Tissue bioconcentration and effects of fluoxetine in zebrafish (*Danio rerio*) and red crucian carp (*Carassius auratus*) after short-term and long-term exposure. *Chemosphere* 205, 8–14.
- Pavlov, K.A., Chistiakov, D.A., Chekhonin, V.P., 2012. Genetic determinants of aggression and impulsivity in humans. *J. Appl. Genet.* 53, 61–82.
- Pelli, Marco, Connaughton, Victoria P., 2015. Chronic exposure to environmentally relevant concentrations of fluoxetine (Prozac) decreases survival, increases abnormal behaviours, and delays predator escape responses in guppies. *Chemosphere* 139, 202–209.
- Polverino, G., Martin, J.M., Bertram, M.G., Soman, V.R., Tan, H., Brand, J.A., Mason, R. T., Wong, B.B.M., 2021. Psychoactive pollution suppresses individual differences in fish behaviour. *Proc. R. Soc. A B.* 288, 20202294.
- Puckowski, A., Mioduszewska, K., Łukasiewicz, P., Borecka, M., Caban, M., Maszkowska, J., Stepnowski, P., 2016. Bioaccumulation and analytics of pharmaceutical residues in the environment: a review. *J. Pharmaceut. Biomed. Anal.* 127, 232–255.
- Richards, S.M., Wilson, C.J., Johnson, D.J., Castle, D.M., Lam, M., Mabury, S.A., Sibley, P.K., Solomon, K.R., 2009. Effects of pharmaceutical mixtures in aquatic microcosms. *Environ. Toxicol. Chem.* 23, 1035–1042.
- Saaristo, M., Lagesson, A., Bertram, M.G., Fick, J., Klaminder, J., Johnstone, C.P., Wong, B.B.M., Brodin, T., 2019. Behavioural effects of psychoactive pharmaceutical exposure on European perch (*Perca fluviatilis*) in a multi-stressor environment. *Sci. Total Environ.* 655, 1311–1320.
- Salahinejad, A., Attaran, A., Meuthen, D., Chivers, D.P., Niyogi, S., 2022. Review - proximate causes and ultimate effects of common antidepressants, fluoxetine and venlafaxine, on fish behaviour. *Sci. Total Environ.* 807, 150846.
- Santarencchi, E., Sprugnoli, G., Tatti, E., Mencarelli, E., Neri, F., Momi, D., Di Lorenzo, G., Pascual-Leone, A., Rossi, S., Rossi, A., 2018. Brain functional connectivity correlates of coping styles. *Cognit. Affect Behav. Neurosci.* 18, 495–508.
- Sehonova, P., Svobodova, Z., Dolezelova, P., Vosmerova, P., Faggio, C., 2018. Effects of waterborne antidepressants on non-target animals living in the aquatic environment: a review. *Sci. Total Environ.* 631–632, 789–794.
- Sehonova, P., Hodkovicova, N., Urbanova, M., Ůrn, S., Blahova, J., Svobodova, Z., Faldyna, M., Chloupek, P., Briedikova, K., Carlsson, G., 2019. Effects of antidepressants with different modes of action on early life stages of fish and amphibians. *Environ. Pollut.* 254, 112999.
- Silva, L.J.G., Lino, C.M., Meisel, L.M., Pena, A., 2012. Review - selective serotonin reuptake inhibitors (SSRIs) in the aquatic environment: an ecopharmacovigilance approach. *Sci. Total Environ.* 437, 185–195.
- Silva, L.J.G., Pereira, A.M.P.T., Meisel, L.M., Lino, C.M., 2015. Reviewing the serotonin reuptake inhibitors (SSRIs) footprint in the aquatic biota: uptake, bioaccumulation and ecotoxicology. *Environ. Pollut.* 197, 127–143.
- Silva, L.J.G., Pereira, A.M.P.T., Rodrigues, H., Meisel, L.M., Lino, C.M., Pena, A., 2017. SSRIs antidepressants in marine mussels from Atlantic coastal areas and human risk assessment. *Sci. Total Environ.* 603–604, 118–125.
- Silva, M.S.S., Oliveira, M., Valente, P., Figueira, E., Martins, M., Pires, A., 2020. Behavior and biochemical responses of the polychaeta *Hediste diversicolor* to polystyrene nanoplastics. *Sci. Total Environ.* 707, 134434.
- Singh, A., Saidulu, D., Gupta, A.K., Kubsad, V., 2022. Occurrence and fate of antidepressants in the aquatic environment: insights into toxicological effects on the aquatic life, analytical methods, and removal techniques. *J. Environ. Chem. Eng.* 10 (6).
- Sumpter, J.P., Margiotta-Casaluci, L., 2022. Environmental occurrence and predicted pharmacological risk to freshwater fish of over 200 neuroactive pharmaceuticals in widespread use. *Toxics* 10, 233.
- Sumpter, J.P., Donnachie, R.L., Johnson, A.C., 2014. The apparently very variable potency of the anti-depressant fluoxetine. *Aquat. Toxicol.* 151, 57–60.
- Tan, H., Poverino, G., Martin, J.M., Bertram, M.G., Wiles, S.C., Palacios, M.M., Bywater, C.L., White, C.R., Wong, B.B.M., 2020. Chronic exposure to a pervasive pharmaceutical pollutant erodes among-individual phenotypic variation in a fish. *Environ. Pollut.* 263, 114450.
- Tanoue, R., Margiotta-Casaluci, L., Huerta, B., Runnals, T., Eguchi, A., Nomiyama, K., Kunisue, T., Tanabe, S., Sumpter, J., 2019. Protecting the environment from psychoactive drugs: problems for regulators illustrated by the possible effects of tramadol on fish behaviour. *Sci. Total Environ.* 664, 915–926.
- Theodoridi, A., Tsalafouta, A., Pavlidis, M., 2017. Acute exposure to fluoxetine alters aggressive behaviour of zebrafish and expression of genes involved in serotonergic system regulation. *Front. Neurosci.* 11, 223.
- Thibaut, R., Porte, C., 2008. Effects of fibrates, anti-inflammatory drugs and antidepressants in the fish hepatoma cell line PLHC-1: cytotoxicity and interactions with cytochrome P450 1A. *Toxicol. Vitro* 22, 1128–1135.
- Thörnqvist, P.O., McCarrick, S., Ericsson, M., Roman, E., Winberg, S., 2019. Bold zebrafish (*Danio rerio*) express higher levels of delta opioid and dopamine D2 receptors in the brain compared to shy fish. *Behav. Brain Res.* 359, 927–934.
- Tran, S., Nowicki, M., Muraleetharan, A., Chatterjee, D., Gerlai, R., 2016. Neurochemical factors underlying individual differences in locomotor activity and anxiety-like behavioural responses in zebrafish. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 65, 25–33.
- Tudorache, C., Schaaf, M.J., Slabbekoorn, H., 2013. Covariation between behaviour and physiology indicators of coping style in zebrafish (*Danio rerio*). *J. Endocrinol.* 219, 251–258.
- Tudorache, C., Braake, A., Tromp, M., Slabbekoorn, H., Schaaf, M.J., 2015. Behavioural and physiological indicators of stress coping styles in larval zebrafish. *Stress* 18 (1), 121–128.
- Tudorache, C., Slabbekoorn, H., Robbers, Y., Hin, E., Meijer, J.H., Spaik, H.P., Schaaf, M.J.M., 2018. Biological clock function is linked to proactive and reactive personality types. *BMC Biol.* 16, 148.
- Vinterstare, J., Brönmark, C., Nilsson, P.A., Langerhans, R.B., Berglund, O., Örsjes, J., Brodin, T., Fick, J., Hulthén, K., 2021. Antipredator phenotype in crucian carp altered by a psychoactive drug. *Ecol. Evol.* 11, 9435–9446.
- Wei, M., He, S., Chen, A., Fan, Z., Liu, W., Zhang, L., Lv, S., Qiu, C., Liu, H., Hao, M., Yin, Z., Ren, D., 2022. Fluoxetine modifies circadian rhythm by reducing melatonin content in zebrafish. *Biomed. Pharmacother.* 153, 113268.
- Weinberger, J., Klaper, R., 2014. Environmental concentrations of the selective serotonin reuptake inhibitor fluoxetine impact specific behaviours involved in reproduction, feeding and predator avoidance in the fish *Pimephales promelas* (fathead minnow). *Aquat. Toxicol.* 151, 77–83.
- Wilson, A.D.M., Godin, J.G.J., 2009. Boldness and behavioural syndromes in the bluegill sunfish, *Lepomis macrochirus*. *Behav. Ecol.* 20, 231–237.
- Wilson, A.D.M., Stevens, E.D., 2005. Consistency in context-specific measures of shyness and boldness in rainbow trout, *Oncorhynchus mykiss*. *Ethology* 111, 849–862.
- Wilson, V., Guenther, A., Øverli, Ø., Seltmann, M.W., Altschul, D., 2019. Future directions for personality research: contributing new insights to the understanding of animal behaviour. *Animals* 9, 240.
- Wong, R.Y., Perrin, F., Oxendine, S.E., Kezios, Z.D., Sawyer, S., Zhou, L.R., Dereje, S., Godwin, J., 2012. Comparing behavioral responses across assays of stress and anxiety in zebrafish (*Danio rerio*). *Behaviour* 149, 1205–1240.
- Wong, R.Y., Oxendine, S.E., Godwin, J., 2013. Behavioural and neurogenomic transcriptome changes in wild-derived zebrafish with fluoxetine treatment. *BMC Genom.* 14 (1), 1–13.
- Wong, R.Y., French, J., Russ, J.B., 2019. Differences in Stress Reactivity between Zebrafish with Alternative Stress Coping Styles, vol. 6. Royal Society Open Science, pp. 1–12.
- Xie, H., Li, M., Kang, Y., et al., 2022. Zebrafish: an important model for understanding scoliosis. *Cell. Mol. Life Sci.* 79, 506.
- Xin, Y., Wu, J., Yao, Z., Guan, Q., Aleman, A., Luo, Y., 2017. The relationship between personality and the response to acute psychological stress. *Sci. Rep.* 7, 16906.
- Yang, H., Liang, X., Zhao, Y., Gu, X., Mao, Z., Zeng, Q., Chen, H., Martyniuk, C.J., 2021. Molecular and behavioural responses of zebrafish embryos/larvae after sertraline exposure. *Ecotoxicol. Environ. Saf.* 208, 111700.
- Yuan, M., Chen, Y., Huang, Y., Lu, W., 2018. Behavioural and metabolic phenotype indicate personality in zebrafish (*Danio rerio*). *Frontiers in Physiology. Aquatic Physiology* 9, 653.
- Zacarias, C.C., Barocio, M.E., Vázquez, E.H., Hernández, J.E.S., Arroyo, L.P., Pacheco, I. Y.L., Barceló, D., Iqbal, H.N.M., Saldívar, R.P., 2021. Antidepressant drugs as emerging contaminants: occurrence in urban and non-urban waters and analytical methods for their detection (Review). *Sci. Total Environ.* 757, 143722.
- Zozulya, A.A., Gabaeva, M.V., Sokolov, O.Y., Surkina, I.D., Kost, N.V., 2008. Personality, coping style, and constitutional neuroimmunology. *J. Immunol.* 5, 221–225.